

Table 4B
2004 Pediatric Nutrition Surveillance ⁽¹⁾
North Carolina
Comparison of Racial and Ethnic Distribution by County
Children Aged < 5 Years ⁽²⁾

Comparison of Racial and Ethnic Distribution by County							
County	Number Included in Analysis (3)	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaskan Native %	Asian/Pacific Islander %	All Other / Unknown %
Alamance	2,759	39.8	26.7	33.0	0.1	0.5	0.0
Alexander	911	70.1	5.7	20.4	0.1	3.6	0.0
Alleghany	343	76.7	0.6	22.7	0.0	0.0	0.0
Anson	1,009	29.3	66.0	3.1	1.2	0.4	0.0
Ashe	912	84.8	1.4	13.8	0.0	0.0	0.0
Avery	253	88.5	0.4	10.3	0.0	0.8	0.0
Beaufort	1,557	32.9	42.6	24.3	0.0	0.1	0.0
Bertie	798	13.3	84.7	2.0	0.0	0.0	0.0
Bladen	1,230	32.8	44.3	22.3	0.7	0.0	0.0
Brunswick	2,232	64.8	17.9	16.7	0.0	0.5	0.0
Buncombe	4,933	68.0	13.0	18.3	0.2	0.5	0.0
Burke	2,102	65.5	8.0	18.6	0.0	7.8	0.0
Cabarrus	649	3.5	14.3	82.1	0.0	0.0	0.0
Caldwell	2,319	72.1	13.3	14.2	0.0	0.3	0.0
Camden	235	71.5	26.8	0.4	0.0	1.3	0.0
Carteret	1,019	67.0	17.3	15.3	0.0	0.4	0.0
Caswell	596	47.8	46.6	5.5	0.0	0.0	0.0
Catawba	3,103	48.5	14.5	28.9	0.1	8.0	0.0
Chatham	1,558	21.4	12.6	65.7	0.0	0.2	0.0

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with errors.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 4B
2004 Pediatric Nutrition Surveillance ⁽¹⁾
North Carolina
Comparison of Racial and Ethnic Distribution by County
Children Aged < 5 Years ⁽²⁾

Comparison of Racial and Ethnic Distribution by County							
County	Number Included in Analysis (3)	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaskan Native %	Asian/Pacific Islander %	All Other / Unknown %
Cherokee	795	90.1	4.0	3.9	1.1	0.9	0.0
Chowan	537	32.2	62.9	4.8	0.0	0.0	0.0
Clay	283	95.8	1.4	2.5	0.4	0.0	0.0
Cleveland	761	59.4	34.6	5.1	0.0	0.9	0.0
Columbus	1,351	36.4	46.0	14.7	2.7	0.1	0.0
Craven	3,412	48.4	36.8	13.2	0.3	1.2	0.0
Cumberland	13,431	42.7	44.4	10.0	1.1	1.8	0.0
Currituck	464	87.3	9.3	3.2	0.0	0.2	0.0
Dare	613	65.6	5.2	29.0	0.0	0.2	0.0
Davidson	3,745	61.2	12.8	24.6	0.1	1.3	0.0
Davie	842	59.7	9.0	31.1	0.0	0.1	0.0
Duplin	2,084	24.4	20.5	54.6	0.0	0.5	0.0
Durham	6,368	9.3	55.0	33.5	0.0	2.2	0.0
Edgecombe	796	25.1	66.5	8.3	0.1	0.0	0.0
Forsyth	7,959	23.8	31.7	43.7	0.1	0.7	0.0
Franklin	1,500	31.3	40.4	26.9	0.5	1.0	0.0
Gaston	5,071	50.7	27.4	20.2	0.1	1.7	0.0
Gates	229	44.1	53.3	1.3	1.3	0.0	0.0
Graham	314	88.9	1.0	3.8	6.4	0.0	0.0

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with errors.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 4B
2004 Pediatric Nutrition Surveillance ⁽¹⁾
North Carolina
Comparison of Racial and Ethnic Distribution by County
Children Aged < 5 Years ⁽²⁾

Comparison of Racial and Ethnic Distribution by County							
County	Number Included in Analysis (3)	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaskan Native %	Asian/Pacific Islander %	All Other / Unknown %
Granville	1,371	35.0	42.2	22.5	0.0	0.3	0.0
Greene	1,047	17.2	43.5	39.4	0.0	0.0	0.0
Guilford	7,169	22.7	54.2	19.4	0.2	3.6	0.0
Halifax	2,334	17.7	70.0	3.8	7.8	0.7	0.0
Harnett	3,156	42.1	35.0	22.5	0.2	0.1	0.0
Haywood	1,244	88.1	1.6	9.8	0.1	0.4	0.0
Henderson	2,066	51.8	2.7	44.9	0.0	0.6	0.0
Hertford	948	15.6	82.5	1.3	0.2	0.4	0.0
Hoke	1,947	21.4	33.5	32.3	12.8	0.1	0.0
Hyde	166	42.2	33.7	22.9	0.0	1.2	0.0
Iredell	3,166	50.3	24.8	22.8	0.1	2.1	0.0
Jackson	735	87.8	1.9	7.1	2.4	0.8	0.0
Johnston	2,808	44.4	21.4	33.9	0.0	0.3	0.0
Jones	294	44.9	41.8	13.3	0.0	0.0	0.0
Lee	1,785	29.2	25.7	44.6	0.1	0.4	0.0
Lenoir	2,130	25.1	59.5	15.1	0.0	0.3	0.0
Lincoln	1,668	71.6	6.1	22.1	0.0	0.2	0.0
Macon	892	82.4	1.2	16.3	0.0	0.1	0.0
Madison	587	93.2	0.3	6.5	0.0	0.0	0.0

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with errors.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 4B
2004 Pediatric Nutrition Surveillance ⁽¹⁾
North Carolina
Comparison of Racial and Ethnic Distribution by County
Children Aged < 5 Years ⁽²⁾

Comparison of Racial and Ethnic Distribution by County							
County	Number Included in Analysis (3)	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaskan Native %	Asian/Pacific Islander %	All Other / Unknown %
Martin	866	23.1	67.7	9.0	0.0	0.2	0.0
Mcdowell	1,161	74.2	3.9	21.1	0.1	0.8	0.0
Mecklenbur	14,555	10.8	47.6	36.2	0.3	5.0	0.0
Mitchell	514	87.2	0.2	12.6	0.0	0.0	0.0
Montgomery	1,126	28.8	21.7	48.0	0.0	1.6	0.0
Moore	2,171	46.5	29.3	24.0	0.1	0.1	0.0
Nash	2,784	25.6	54.7	19.1	0.3	0.3	0.0
New Hanove	3,417	44.4	40.2	14.8	0.1	0.5	0.0
Northhampt	828	17.1	76.8	5.9	0.0	0.1	0.0
Onslow	7,962	62.1	22.3	14.2	0.2	1.2	0.0
Orange	2,191	23.2	21.1	54.6	0.0	1.1	0.0
Pamlico	378	57.1	30.4	11.9	0.0	0.5	0.0
Pasquotank	1,233	37.2	55.1	6.3	0.0	1.4	0.0
Pender	1,553	41.9	31.2	19.8	1.7	5.3	0.0
Perquimans	322	52.2	44.4	3.1	0.3	0.0	0.0
Person	1,033	46.3	43.6	10.1	0.1	0.0	0.0
Pitt	4,324	21.0	62.7	15.6	0.1	0.7	0.0
Polk	380	64.2	6.3	28.4	1.1	0.0	0.0
Randolph	4,008	58.4	7.4	34.1	0.0	0.1	0.0

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with errors.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 4B
2004 Pediatric Nutrition Surveillance ⁽¹⁾
North Carolina
Comparison of Racial and Ethnic Distribution by County
Children Aged < 5 Years ⁽²⁾

Comparison of Racial and Ethnic Distribution by County							
County	Number Included in Analysis (3)	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaskan Native %	Asian/Pacific Islander %	All Other / Unknown %
Richmond	2,194	43.6	41.5	13.6	0.9	0.4	0.0
Robeson	3,903	13.6	28.4	21.2	34.7	2.2	0.0
Rockingham	2,182	55.6	24.8	18.5	0.1	0.9	0.0
Rowan	3,437	50.1	25.5	23.4	0.0	1.0	0.0
Rutherford	1,570	64.3	22.4	12.6	0.0	0.7	0.0
Sampson	2,687	16.9	25.5	55.4	1.0	1.2	0.0
Scotland	1,590	24.5	60.1	2.5	12.5	0.4	0.0
Stanly	1,507	60.4	19.9	15.4	0.1	4.2	0.0
Stokes	1,019	86.9	6.0	6.8	0.2	0.1	0.0
Surry	1,908	58.7	5.9	34.6	0.1	0.7	0.0
Swain	402	86.3	1.0	2.2	9.5	1.0	0.0
Transylvan	552	81.7	8.9	8.0	0.5	0.9	0.0
Tyrrell	270	31.9	40.7	27.4	0.0	0.0	0.0
Union	3,503	31.3	23.0	45.4	0.1	0.2	0.0
Vance	1,871	19.1	62.1	18.4	0.1	0.3	0.0
Wake	13,300	15.1	39.2	41.3	0.0	4.5	0.0
Warren	874	19.1	69.3	9.8	1.6	0.1	0.0
Washington	678	17.4	75.5	6.9	0.0	0.1	0.0
Watauga	890	87.5	2.4	9.3	0.4	0.3	0.0

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with errors.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 4B
2004 Pediatric Nutrition Surveillance ⁽¹⁾
North Carolina
Comparison of Racial and Ethnic Distribution by County
Children Aged < 5 Years ⁽²⁾

Comparison of Racial and Ethnic Distribution by County							
County	Number Included in Analysis (3)	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaskan Native %	Asian/Pacific Islander %	All Other / Unknown %
Wayne Coun	3,700	31.9	41.5	25.9	0.2	0.6	0.0
Wilkes	1,846	75.8	5.4	18.4	0.1	0.3	0.0
Wilson Cou	2,971	17.6	51.2	30.3	0.1	0.9	0.0
Yadkin	1,075	50.8	4.0	45.1	0.0	0.1	0.0
Yancey	605	76.4	1.3	22.3	0.0	0.0	0.0
North Carolina	219,926	38.8	33.5	25.0	1.2	1.6	0.0
Nation (Prior Year)	5,091,536	38.8	20.4	32.2	1.3	3.6	3.7

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with errors.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 5B
2004 Pediatric Nutrition Surveillance ⁽¹⁾
North Carolina
Comparison of Age Distribution by County
Children Aged < 5 Years ⁽²⁾

Comparison of Age Distribution by County				
County	Number Included in Analysis (3,4)	0 - 11 Months %	12 - 23 Months %	24 - 59 Months %
Alamance	2,759	41.1	22.4	36.5
Alexander	911	34.0	22.6	43.4
Alleghany	343	30.0	19.2	50.7
Anson	1,009	31.3	23.7	45.0
Ashe	912	33.4	20.4	46.2
Avery	253	58.5	6.3	35.2
Beaufort	1,557	34.3	23.4	42.3
Bertie	798	29.3	19.9	50.8
Bladen	1,230	38.4	21.8	39.8
Brunswick	2,232	36.3	21.6	42.2
Buncombe	4,933	35.7	22.9	41.4
Burke	2,102	35.8	23.1	41.1
Cabarrus	649	24.2	28.5	47.3
Caldwell	2,319	38.6	20.8	40.5
Camden	235	27.7	27.7	44.7
Carteret	1,019	37.1	17.5	45.4
Caswell	596	35.1	22.3	42.6
Catawba	3,103	40.4	21.3	38.2

(1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Records rejected if date of birth and date of visit unknown.
* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 5B
2004 Pediatric Nutrition Surveillance ⁽¹⁾
North Carolina
Comparison of Age Distribution by County
Children Aged < 5 Years ⁽²⁾

Comparison of Age Distribution by County				
County	Number Included in Analysis (3,4)	0 - 11 Months %	12 - 23 Months %	24 - 59 Months %
Chatham	1,558	40.1	23.0	36.8
Cherokee	795	35.0	22.1	42.9
Chowan	537	33.0	21.0	46.0
Clay	283	30.7	21.2	48.1
Cleveland	761	84.9	9.1	6.0
Columbus	1,351	23.2	24.9	51.8
Craven	3,412	40.3	20.5	39.2
Cumberland	13,431	37.2	22.7	40.1
Currituck	464	40.9	23.3	35.8
Dare	613	33.4	23.5	43.1
Davidson	3,745	41.1	20.5	38.4
Davie	842	32.1	22.4	45.5
Duplin	2,084	37.8	14.7	47.6
Durham	6,368	31.7	22.4	45.9
Edgecombe	796	91.5	4.1	4.4
Forsyth	7,959	40.7	20.9	38.4
Franklin	1,500	37.1	12.3	50.6
Gaston	5,071	46.6	20.4	33.0

(1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Records rejected if date of birth and date of visit unknown.
* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 5B
2004 Pediatric Nutrition Surveillance ⁽¹⁾
North Carolina
Comparison of Age Distribution by County
Children Aged < 5 Years ⁽²⁾

Comparison of Age Distribution by County				
County	Number Included in Analysis (3,4)	0 - 11 Months %	12 - 23 Months %	24 - 59 Months %
Gates	229	31.9	15.3	52.8
Graham	314	29.0	26.8	44.3
Granville	1,371	33.3	19.8	46.9
Greene	1,047	30.9	20.9	48.1
Guilford	7,169	42.4	7.2	50.4
Halifax	2,334	32.0	26.3	41.6
Harnett	3,156	39.1	18.4	42.5
Haywood	1,244	32.8	23.9	43.3
Henderson	2,066	34.2	22.5	43.3
Hertford	948	36.8	24.5	38.7
Hoke	1,947	34.7	21.2	44.1
Hyde	166	30.1	11.4	58.4
Iredell	3,166	33.9	25.0	41.1
Jackson	735	30.5	21.6	47.9
Johnston	2,808	37.7	24.4	37.9
Jones	294	37.4	14.3	48.3
Lee	1,785	37.6	22.1	40.3
Lenoir	2,130	42.3	18.2	39.5

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Records rejected if date of birth and date of visit unknown.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 5B
2004 Pediatric Nutrition Surveillance ⁽¹⁾
North Carolina
Comparison of Age Distribution by County
Children Aged < 5 Years ⁽²⁾

Comparison of Age Distribution by County				
County	Number Included in Analysis (3,4)	0 - 11 Months %	12 - 23 Months %	24 - 59 Months %
Lincoln	1,668	40.9	18.9	40.2
Macon	892	35.1	20.3	44.6
Madison	587	31.2	20.4	48.4
Martin	866	35.9	23.2	40.9
Mcdowell	1,161	33.0	20.7	46.3
Mecklenbur	14,555	35.0	25.3	39.8
Mitchell	514	30.7	21.2	48.1
Montgomery	1,126	35.8	18.7	45.5
Moore	2,171	33.6	21.0	45.4
Nash	2,784	40.6	22.0	37.5
New Hanove	3,417	46.0	20.7	33.3
Northhampt	828	30.0	21.4	48.7
Onslow	7,962	38.2	22.7	39.1
Orange	2,191	30.0	24.2	45.8
Pamlico	378	38.4	16.1	45.5
Pasquotank	1,233	35.4	26.8	37.8
Pender	1,553	35.5	19.8	44.7
Perquimans	322	34.2	24.5	41.3

(1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Records rejected if date of birth and date of visit unknown.
* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 5B
2004 Pediatric Nutrition Surveillance ⁽¹⁾
North Carolina
Comparison of Age Distribution by County
Children Aged < 5 Years ⁽²⁾

Comparison of Age Distribution by County				
County	Number Included in Analysis (3,4)	0 - 11 Months %	12 - 23 Months %	24 - 59 Months %
Person	1,033	34.8	20.9	44.2
Pitt	4,324	34.0	21.5	44.5
Polk	380	40.0	17.9	42.1
Randolph	4,008	44.7	18.1	37.2
Richmond	2,194	33.3	19.1	47.6
Robeson	3,903	36.6	9.1	54.3
Rockingham	2,182	33.7	23.1	43.2
Rowan	3,437	37.9	20.9	41.2
Rutherford	1,570	36.0	18.5	45.5
Sampson	2,687	32.6	24.9	42.5
Scotland	1,590	28.7	25.7	45.7
Stanly	1,507	36.1	20.9	43.0
Stokes	1,019	33.0	23.1	44.0
Surry	1,908	36.3	22.2	41.5
Swain	402	31.1	21.1	47.8
Transylvan	552	44.2	12.7	43.1
Tyrrell	270	29.3	18.9	51.9
Union	3,503	32.5	25.7	41.9

(1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Records rejected if date of birth and date of visit unknown.
* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 5B
2004 Pediatric Nutrition Surveillance ⁽¹⁾
North Carolina
Comparison of Age Distribution by County
Children Aged < 5 Years ⁽²⁾

Comparison of Age Distribution by County				
County	Number Included in Analysis (3,4)	0 - 11 Months %	12 - 23 Months %	24 - 59 Months %
Vance	1,871	32.3	23.0	44.7
Wake	13,300	32.8	25.3	41.9
Warren	874	28.7	22.9	48.4
Washington	678	29.2	21.2	49.6
Watauga	890	34.2	22.6	43.3
Wayne Coun	3,700	33.9	24.6	41.5
Wilkes	1,846	37.8	18.6	43.6
Wilson Cou	2,971	39.2	20.6	40.2
Yadkin	1,075	34.7	21.1	44.2
Yancey	605	28.1	22.8	49.1
North Carolina	219,926	36.8	21.3	41.8
Nation (Prior Year)	5,091,536	36.5	21.5	41.9

(1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Records rejected if date of birth and date of visit unknown.
* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 6B
2004 Pediatric Nutrition Surveillance (1)
North Carolina
Comparison of Growth and Anemia Indicators by County
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators by County (3,4)												
County	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)		
	Low		High	Short Stature (6)		Under- weight (7)	Over- weight (7)	≥ 2 Yrs Over- weight (7)				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Alamance	815	8.7 (27)	9.0 (85)	2,648	5.4 (45)	7.2 (37)	18.5 (86)	984	18.6 (81)	18.2 (73)	1,913	31.9 (96)
Alexander	265	9.1 (36)	6.8 (38)	592	18.6(100)	8.1 (56)	19.4 (91)	331	16.9 (56)	20.8 (91)	639	15.0 (71)
Alleghany	80	* *	* *	292	7.9 (88)	6.2 (20)	19.9 (94)	161	21.1 (94)	24.8 (95)	239	8.4 (35)
Anson	262	8.0 (20)	4.6 (9)	761	6.3 (62)	7.8 (50)	20.0 (95)	421	11.2 (8)	14.7 (36)	704	12.1 (61)
Ashe	228	6.6 (5)	4.4 (7)	688	6.3 (62)	7.1 (34)	17.0 (72)	386	11.7 (9)	15.5 (47)	606	10.4 (50)
Avery	123	12.2 (76)	9.8 (90)	139	11.5 (99)	7.2 (37)	19.4 (91)	80	* *	* *	90	* *
Beaufort	432	10.2 (50)	6.3 (26)	1,286	5.5 (48)	6.3 (22)	15.6 (53)	627	16.9 (56)	16.7 (63)	1,122	10.4 (50)
Bertie	167	16.8 (91)	3.6 (3)	678	8.6 (93)	6.3 (22)	17.6 (81)	367	13.1 (17)	17.7 (72)	623	26.8 (94)
Bladen	375	10.9 (65)	6.7 (36)	1,181	3.5 (10)	11.7 (85)	12.9 (21)	493	15.2 (35)	13.8 (29)	762	5.9 (21)
Brunswick	659	8.6 (26)	8.8 (79)	1,890	7.4 (83)	6.9 (30)	15.2 (49)	870	17.1 (60)	14.7 (36)	1,400	4.6 (11)
Buncombe	1,409	9.5 (40)	8.7 (77)	3,721	5.8 (54)	4.5 (4)	16.4 (66)	1,882	17.7 (70)	16.6 (62)	3,417	11.7 (60)

- (1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this county's rate to other counties. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6B
2004 Pediatric Nutrition Surveillance (1)
North Carolina
Comparison of Growth and Anemia Indicators by County
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators by County (3,4)												
County	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)	Under- weight (7)	Over- weight (7)		≥ 2 Yrs Over- weight (7)				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Burke	629	7.9 (18)	4.3 (6)	1,741	4.4 (23)	11.1 (83)	11.8 (12)	794	10.7 (5)	10.8 (9)	1,463	7.3 (27)
Cabarrus	137	7.3 (10)	5.1 (16)	625	4.0 (18)	10.1 (79)	16.0 (56)	300	13.3 (19)	16.3 (56)	444	6.8 (25)
Caldwell	761	9.6 (43)	7.6 (57)	1,735	6.9 (75)	7.2 (37)	16.5 (67)	872	19.4 (87)	15.5 (47)	1,545	7.1 (26)
Camden	53	* *	* *	214	6.5 (68)	14.5 (95)	13.1 (23)	99	* *	* *	141	8.5 (36)
Carteret	276	10.5 (56)	9.1 (87)	837	5.1 (33)	11.1 (83)	14.3 (39)	382	15.7 (42)	19.1 (81)	730	9.0 (39)
Caswell	168	12.5 (78)	4.2 (5)	476	5.5 (48)	8.2 (57)	13.4 (29)	233	13.3 (19)	11.2 (11)	370	5.1 (13)
Catawba	927	9.8 (46)	6.9 (40)	1,562	4.7 (26)	18.5 (98)	8.6 (3)	534	19.3 (86)	15.4 (46)	1,979	9.1 (42)
Chatham	498	8.4 (22)	7.0 (44)	1,116	7.1 (77)	6.9 (30)	17.0 (72)	497	15.9 (45)	18.9 (79)	933	10.9 (55)
Cherokee	220	11.4 (68)	7.7 (60)	432	6.3 (62)	12.3 (87)	15.5 (52)	145	17.9 (73)	20.7 (90)	531	10.2 (47)
Chowan	140	12.1 (74)	7.9 (65)	527	5.5 (48)	10.4 (80)	13.1 (23)	248	16.9 (56)	13.7 (28)	386	17.9 (79)
Clay	72	* *	* *	216	7.9 (88)	3.7 (1)	13.4 (29)	115	13.0 (15)	19.1 (81)	204	3.4 (5)

- (1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this county's rate to other counties. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6B
2004 Pediatric Nutrition Surveillance (1)
North Carolina
Comparison of Growth and Anemia Indicators by County
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators by County (3,4)												
County	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)	Under- weight (7)	Over- weight (7)		≥ 2 Yrs Over- weight (7)				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% >95th (Rank)	Number	% 85th- <95th (Rank)	% >95th (Rank)	Number	% (Rank)
Cleveland	618	9.2 (38)	7.6 (57)	664	2.6 (2)	12.8 (90)	13.0 (22)	39	* *	* *	120	8.3 (34)
Columbus	189	14.8 (89)	8.5 (74)	1,282	7.1 (77)	4.6 (6)	18.9 (89)	669	13.8 (23)	16.1 (52)	919	18.9 (82)
Craven	1,027	10.2 (50)	9.2 (88)	2,362	4.1 (19)	7.8 (50)	13.1 (23)	1,136	16.5 (50)	14.2 (32)	2,086	14.1 (66)
Cumberland	3,970	10.1 (49)	7.6 (57)	11,756	4.7 (26)	12.2 (86)	11.8 (12)	4,976	12.9 (14)	9.8 (5)	9,239	19.2 (85)
Currituck	166	7.2 (7)	7.2 (47)	440	7.5 (85)	7.0 (33)	15.7 (54)	167	14.4 (31)	13.2 (23)	283	7.8 (30)
Dare	122	4.9 (2)	7.4 (51)	442	7.2 (80)	4.5 (4)	17.4 (77)	221	19.9 (90)	16.3 (56)	392	10.2 (47)
Davidson	1,131	10.5 (56)	6.5 (29)	2,937	5.2 (36)	7.3 (44)	13.2 (26)	1,308	13.2 (18)	12.7 (21)	2,527	6.5 (22)
Davie	195	5.1 (3)	8.2 (73)	631	8.1 (91)	5.1 (11)	16.2 (60)	328	21.6 (95)	14.3 (34)	640	6.7 (24)
Duplin	709	9.7 (44)	4.9 (11)	1,468	4.1 (19)	15.6 (96)	15.1 (46)	633	13.7 (22)	19.1 (81)	791	27.6 (95)
Durham	1,360	9.5 (40)	6.6 (34)	2,602	7.2 (80)	8.6 (63)	16.1 (58)	1,204	12.0 (10)	13.0 (22)	315	14.6 (70)
Edgecombe	718	12.7 (80)	3.8 (4)	340	6.5 (68)	33.5(100)	6.8 (1)	23	* *	* *	71	* *

- (1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this county's rate to other counties. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6B
2004 Pediatric Nutrition Surveillance (1)
North Carolina
Comparison of Growth and Anemia Indicators by County
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators by County (3,4)												
County	Birthweight (5)			Height and Weight							Anemia Low Hb/Hct (8)	
	Low	High		Short Stature (6)	Under- weight (7)	Over- weight (7)		≥ 2 Yrs Over- weight (7)				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Forsyth	2,483	10.5 (56)	6.1 (24)	6,472	5.5 (48)	10.4 (80)	14.5 (43)	2,959	15.8 (43)	15.0 (39)	5,188	13.5 (65)
Franklin	403	7.9 (18)	8.9 (83)	1,057	3.2 (5)	9.3 (72)	14.7 (44)	598	17.6 (69)	15.7 (49)	749	5.1 (13)
Gaston	1,676	9.5 (40)	6.9 (40)	2,201	6.5 (68)	8.3 (58)	10.9 (6)	516	15.5 (40)	13.2 (23)	2,792	2.7 (4)
Gates	57	* *	* *	190	4.7 (26)	6.8 (27)	11.6 (8)	111	13.5 (21)	12.6 (20)	148	12.8 (62)
Graham	73	* *	* *	237	3.8 (13)	4.2 (3)	17.3 (76)	130	12.3 (12)	18.5 (76)	229	3.9 (7)
Granville	414	8.9 (31)	5.1 (16)	1,074	3.0 (3)	9.6 (76)	15.3 (50)	536	17.5 (65)	18.5 (76)	860	19.1 (83)
Greene	215	14.0 (85)	6.0 (22)	859	7.5 (85)	4.0 (2)	21.2 (97)	457	13.8 (23)	18.2 (73)	738	17.8 (78)
Guilford	2,452	10.7 (60)	6.0 (22)	5,264	5.4 (45)	10.0 (78)	10.9 (6)	3,172	13.9 (25)	12.5 (19)	3,574	5.6 (18)
Halifax	718	13.8 (84)	4.9 (11)	1,643	4.7 (26)	7.4 (47)	21.2 (97)	708	14.1 (30)	19.4 (84)	1,595	21.8 (91)
Harnett	991	9.2 (38)	8.1 (70)	2,602	7.3 (82)	6.9 (30)	15.3 (50)	1,218	16.3 (48)	14.0 (31)	1,970	14.4 (69)
Haywood	356	14.0 (85)	8.1 (70)	450	6.0 (57)	8.4 (59)	14.2 (36)	231	16.5 (50)	19.5 (86)	883	5.4 (17)

- (1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this county's rate to other counties. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6B
2004 Pediatric Nutrition Surveillance (1)
North Carolina
Comparison of Growth and Anemia Indicators by County
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators by County (3,4)												
County	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)	Under- weight (7)	Over- weight (7)		≥ 2 Yrs Over- weight (7)				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Henderson	577	8.7 (27)	8.7 (77)	1,072	5.3 (41)	8.7 (66)	13.4 (29)	468	15.2 (35)	13.2 (23)	1,279	15.7 (72)
Hertford	338	13.0 (83)	2.7 (1)	896	3.9 (15)	9.3 (72)	12.7 (19)	365	18.1 (76)	16.7 (63)	578	9.3 (44)
Hoke	518	12.9 (82)	6.8 (38)	1,827	10.0 (95)	6.7 (25)	22.7(100)	817	18.0 (75)	17.5 (69)	1,134	32.8 (97)
Hyde	48	* *	* *	105	1.9 (1)	6.7 (25)	18.1 (85)	75	* *	* *	114	21.1 (89)
Iredell	873	8.4 (22)	6.3 (26)	3,044	5.2 (36)	7.3 (44)	13.6 (32)	1,256	15.4 (37)	12.0 (17)	2,240	3.8 (6)
Jackson	148	6.8 (6)	8.8 (79)	655	6.1 (58)	5.5 (13)	16.3 (63)	318	19.2 (83)	17.6 (70)	509	5.7 (20)
Johnston	805	10.4 (55)	6.7 (36)	1,651	5.8 (54)	7.1 (34)	15.1 (46)	764	16.1 (47)	16.8 (65)	1,727	10.2 (47)
Jones	102	7.8 (15)	8.8 (79)	169	4.7 (26)	8.9 (68)	18.9 (89)	108	16.7 (54)	19.4 (84)	122	10.7 (52)
Lee	554	8.7 (27)	6.3 (26)	1,638	3.9 (15)	7.9 (53)	16.6 (68)	685	17.4 (64)	17.2 (67)	1,175	26.0 (93)
Lenoir	703	12.1 (74)	5.0 (14)	1,906	3.1 (4)	12.7 (89)	12.0 (14)	783	11.1 (7)	9.2 (4)	1,281	10.8 (54)
Lincoln	530	7.2 (7)	7.5 (55)	393	3.3 (6)	13.7 (94)	15.0 (45)	156	7.7 (1)	9.0 (3)	1,092	2.6 (3)

- (1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this county's rate to other counties. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6B
2004 Pediatric Nutrition Surveillance (1)
North Carolina
Comparison of Growth and Anemia Indicators by County
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators by County (3,4)												
County	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)	Under- weight (7)	Over- weight (7)		≥ 2 Yrs Over- weight (7)				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Macon	266	8.3 (21)	9.0 (85)	598	6.9 (75)	4.8 (10)	19.4 (91)	358	18.4 (78)	18.2 (73)	569	5.6 (18)
Madison	133	10.5 (56)	9.8 (90)	552	7.1 (77)	6.2 (20)	18.8 (87)	272	16.5 (50)	20.2 (89)	431	12.8 (62)
Martin	260	11.2 (67)	7.7 (60)	806	5.2 (36)	8.7 (66)	17.6 (81)	344	14.0 (26)	14.5 (35)	546	18.1 (80)
Mcdowell	310	12.6 (79)	5.8 (21)	989	6.1 (58)	7.2 (37)	12.7 (19)	509	17.5 (65)	11.2 (11)	795	18.7 (81)
Mecklenbur	3,768	8.8 (30)	8.0 (66)	9,633	6.3 (62)	7.9 (53)	15.7 (54)	4,425	17.2 (62)	16.4 (59)	6,111	6.5 (22)
Mitchell	124	4.0 (1)	6.5 (29)	390	10.0 (95)	5.9 (17)	16.2 (60)	223	10.8 (6)	16.1 (52)	373	1.1 (1)
Montgomery	297	10.8 (63)	7.4 (51)	1,060	6.6 (72)	8.6 (63)	18.0 (84)	479	18.4 (78)	18.6 (78)	774	4.4 (9)
Moore	567	7.8 (15)	6.9 (40)	1,735	6.1 (58)	7.8 (50)	18.8 (87)	842	18.5 (80)	16.5 (60)	1,579	10.7 (52)
Nash	944	7.6 (13)	7.3 (48)	2,339	5.3 (41)	9.9 (77)	13.9 (34)	944	14.0 (26)	15.9 (50)	1,416	9.0 (39)
New Hanove	1,286	12.0 (73)	7.3 (48)	2,024	7.9 (88)	4.7 (8)	16.0 (56)	926	16.5 (50)	15.2 (42)	1,336	8.0 (33)
Northhampt	196	14.8 (89)	7.1 (45)	727	4.4 (23)	7.2 (37)	14.4 (41)	365	17.8 (71)	15.1 (40)	555	21.3 (90)

- (1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this county's rate to other counties. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6B
2004 Pediatric Nutrition Surveillance (1)
North Carolina
Comparison of Growth and Anemia Indicators by County
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators by County (3,4)												
County	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)		
	Low		High	Short Stature (6)		Under- weight (7)	Over- weight (7)	≥ 2 Yrs Over- weight (7)				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Onslow	2,406	8.9 (31)	7.8 (63)	6,293	4.2 (21)	8.5 (62)	10.7 (5)	2,617	14.4 (31)	10.2 (6)	4,723	5.0 (12)
Orange	498	10.0 (47)	8.6 (76)	1,839	6.2 (61)	7.2 (37)	16.9 (70)	864	17.8 (71)	20.9 (92)	1,527	16.2 (73)
Pamlico	118	8.5 (25)	5.1 (16)	343	6.4 (66)	9.0 (69)	17.2 (74)	167	19.2 (83)	22.2 (93)	153	11.1 (57)
Pasquotank	372	11.6 (70)	3.0 (2)	1,098	5.1 (33)	7.9 (53)	13.2 (26)	455	12.7 (13)	11.2 (11)	738	9.1 (42)
Pender	393	10.7 (60)	7.4 (51)	1,289	5.3 (41)	9.1 (71)	16.2 (60)	569	12.1 (11)	12.3 (18)	1,004	7.9 (32)
Perquimans	77	* *	* *	311	10.0 (95)	8.4 (59)	21.2 (97)	131	16.8 (55)	17.6 (70)	225	14.2 (67)
Person	307	12.7 (80)	8.8 (79)	913	3.3 (6)	7.4 (47)	13.8 (33)	422	15.4 (37)	13.3 (27)	711	5.2 (15)
Pitt	1,083	11.5 (69)	6.5 (29)	3,873	5.2 (36)	5.1 (11)	16.9 (70)	1,855	17.1 (60)	15.9 (50)	3,153	16.7 (75)
Polk	112	8.9 (31)	8.0 (66)	279	3.9 (15)	6.8 (27)	13.3 (28)	144	16.0 (46)	13.9 (30)	262	19.1 (83)
Randolph	1,262	9.7 (44)	8.9 (83)	3,114	4.9 (32)	9.0 (69)	17.5 (80)	1,315	19.2 (83)	24.6 (94)	2,642	4.0 (8)
Richmond	541	7.8 (15)	6.5 (29)	1,860	3.3 (6)	16.2 (97)	10.3 (4)	974	8.1 (2)	8.3 (1)	1,635	24.3 (92)

- (1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this county's rate to other counties. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6B
2004 Pediatric Nutrition Surveillance (1)
North Carolina
Comparison of Growth and Anemia Indicators by County
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators by County (3,4)												
County	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)	Under- weight (7)	Over- weight (7)		≥ 2 Yrs Over- weight (7)				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Robeson	1,198	10.8 (63)	7.8 (63)	2,931	6.4 (66)	8.6 (63)	16.1 (58)	1,980	14.7 (33)	15.1 (40)	2,367	5.2 (15)
Rockingham	552	10.7 (60)	7.1 (45)	2,076	3.3 (6)	13.6 (92)	11.7 (9)	923	10.5 (4)	10.5 (8)	1,620	8.8 (38)
Rowan	1,040	11.9 (72)	6.5 (29)	2,613	3.5 (10)	6.1 (19)	16.6 (68)	1,250	18.2 (77)	16.9 (66)	2,205	11.2 (58)
Rutherford	460	10.0 (47)	6.1 (24)	1,293	5.1 (33)	5.5 (13)	17.2 (74)	666	14.0 (26)	14.9 (38)	1,076	17.4 (77)
Sampson	710	7.5 (12)	9.6 (89)	2,204	6.8 (73)	10.4 (80)	17.4 (77)	1,069	15.6 (41)	16.1 (52)	1,905	20.4 (88)
Scotland	407	9.1 (36)	7.4 (51)	1,199	4.3 (22)	6.8 (27)	16.3 (63)	667	15.4 (37)	14.2 (32)	1,154	17.3 (76)
Stanly	438	12.3 (77)	7.5 (55)	1,061	5.9 (56)	7.3 (44)	15.1 (46)	577	13.0 (15)	11.6 (15)	1,010	42.4 (98)
Stokes	281	14.2 (87)	5.0 (14)	924	5.6 (52)	12.4 (88)	11.7 (9)	417	17.5 (65)	16.5 (60)	605	4.5 (10)
Surry	491	9.0 (35)	7.7 (60)	558	8.2 (92)	22.4 (99)	7.5 (2)	118	19.5 (89)	11.9 (16)	1,385	2.5 (2)
Swain	88	* *	* *	307	6.5 (68)	4.6 (6)	12.1 (15)	176	14.8 (34)	10.2 (6)	290	9.0 (39)
Transylvan	213	6.1 (4)	8.0 (66)	477	3.8 (13)	12.8 (90)	11.7 (9)	220	19.1 (82)	13.2 (23)	330	8.5 (36)

- (1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this county's rate to other counties. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6B
2004 Pediatric Nutrition Surveillance (1)
North Carolina
Comparison of Growth and Anemia Indicators by County
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators by County (3,4)												
County	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)	Under- weight (7)	Over- weight (7)		≥ 2 Yrs Over- weight (7)				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Tyrrell	64	* *	* *	232	10.8 (98)	4.7 (8)	14.2 (36)	129	14.0 (26)	16.3 (56)	160	20.0 (86)
Union	966	7.7 (14)	7.3 (48)	3,010	6.8 (73)	5.9 (17)	14.3 (39)	1,379	16.9 (56)	15.2 (42)	2,427	11.0 (56)
Vance	578	11.1 (66)	4.8 (10)	1,531	5.2 (36)	9.5 (74)	14.4 (41)	780	16.4 (49)	15.3 (44)	1,157	16.2 (73)
Wake	3,501	8.4 (22)	8.0 (66)	12,868	5.3 (41)	7.7 (49)	14.2 (36)	5,453	17.9 (73)	16.2 (55)	8,627	7.6 (28)
Warren	180	14.4 (88)	4.4 (7)	817	7.6 (87)	5.8 (16)	12.6 (18)	406	15.8 (43)	11.1 (10)	669	10.0 (46)
Washington	165	7.3 (10)	5.5 (20)	546	4.6 (25)	9.5 (74)	13.9 (34)	303	10.2 (3)	8.6 (2)	466	20.0 (86)
Watauga	247	11.7 (71)	4.9 (11)	770	3.6 (12)	8.4 (59)	12.3 (16)	376	17.3 (63)	11.4 (14)	559	11.6 (59)
Wayne Coun	1,176	10.3 (54)	8.1 (70)	3,174	4.8 (31)	13.6 (92)	12.5 (17)	1,446	19.4 (87)	15.3 (44)	2,419	13.1 (64)
Wilkes	597	8.9 (31)	6.9 (40)	1,286	5.4 (45)	6.5 (24)	16.3 (63)	742	20.1 (91)	17.4 (68)	1,142	7.8 (30)
Wilson Cou	888	10.2 (50)	5.2 (19)	2,674	5.6 (52)	7.1 (34)	20.4 (96)	1,128	17.5 (65)	20.0 (88)	2,025	9.4 (45)
Yadkin	293	10.2 (50)	8.5 (74)	755	7.4 (83)	7.2 (37)	17.7 (83)	381	20.5 (92)	18.9 (79)	691	14.3 (68)

- (1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this county's rate to other counties. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6B
2004 Pediatric Nutrition Surveillance (1)
North Carolina
Comparison of Growth and Anemia Indicators by County
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators by County (3,4)												
County	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)	Under- weight (7)	Over- weight (7)	≥ 2 Yrs Over- weight (7)					
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- (Rank)	% <95th (Rank)	% ≥95th (Rank)	Number
Yancey	152	7.2 (7)	6.6 (34)	470	9.1 (94)	5.7 (15)	17.4 (77)	278	20.9 (93)	19.8 (87)	421	7.6 (28)
North Carolina	63,940	9.8	7.1	169,527	5.5	8.7	14.7	78,020	15.8	14.9	133,569	11.7
Nation (Prior Year)	1,172,866	9.1	7.3	4,838,257	6.2	5.2	13.6	2,044,705	15.7	14.7	3,118,041	12.8

- (1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this county's rate to other counties. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7B
2004 Pediatric Nutrition Surveillance (1)
North Carolina

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by County (2)
Children Aged < 5 Years (3)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by County (4,5)						
County	Breastfeeding (6)					
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)
Alamance	815	46.0 (56)	328	22.0 (13)	421	19.2 (13)
Alexander	265	45.3 (58)	97	* *	140	15.0 (21)
Alleghany	81	* *	27	* *	35	* *
Anson	262	35.5 (79)	85	* *	156	4.5 (67)
Ashe	230	48.3 (52)	66	* *	96	* *
Avery	123	71.5 (13)	6	* *	7	* *
Beaufort	432	44.2 (61)	165	13.3 (37)	244	11.1 (41)
Bertie	168	34.5 (80)	75	* *	102	2.9 (72)
Bladen	376	49.5 (49)	99	* *	166	13.9 (27)
Brunswick	658	40.0 (69)	186	19.9 (16)	309	12.6 (31)
Buncombe	1,415	76.0 (8)	515	27.0 (8)	746	16.6 (17)
Burke	629	52.0 (44)	216	11.6 (43)	318	9.1 (50)
Cabarrus	137	74.5 (10)	60	* *	131	27.5 (5)

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this county's rate to other counties. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7B
2004 Pediatric Nutrition Surveillance (1)
North Carolina

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by County (2)
Children Aged < 5 Years (3)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by County (4,5)						
County	Breastfeeding (6)					
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)
Caldwell	761	63.1 (21)	213	11.7 (42)	299	8.0 (58)
Camden	53	* *	26	* *	48	* *
Carteret	276	43.1 (62)	96	* *	102	8.8 (54)
Caswell	167	47.9 (53)	53	* *	91	* *
Catawba	929	56.2 (29)	348	19.0 (19)	435	14.9 (23)
Chatham	500	74.6 (9)	136	33.1 (3)	197	29.9 (4)
Cherokee	221	54.3 (36)	75	* *	113	17.7 (16)
Chowan	140	30.7 (85)	40	* *	74	* *
Clay	72	* *	24	* *	37	* *
Cleveland	623	26.5 (88)	30	* *	58	* *
Columbus	192	51.0 (46)	133	11.3 (44)	224	8.9 (53)
Craven	1,033	55.0 (33)	335	20.0 (15)	458	16.6 (17)
Cumberland	4,002	55.0 (33)	1,404	13.2 (38)	1,955	12.0 (34)

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this county's rate to other counties. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7B
2004 Pediatric Nutrition Surveillance (1)
North Carolina

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by County (2)
Children Aged < 5 Years (3)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by County (4,5)						
County	Breastfeeding (6)					
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)
Currituck	166	62.0 (22)	47	* *	76	* *
Dare	128	96.9 (1)	60	* *	83	* *
Davidson	1,139	52.1 (42)	415	16.4 (30)	475	13.3 (29)
Davie	195	63.6 (19)	89	* *	129	20.2 (11)
Duplin	711	52.5 (41)	119	19.3 (18)	180	11.7 (38)
Durham	1,371	49.1 (51)	699	26.6 (9)	963	23.1 (8)
Edgecombe	719	25.9 (89)	12	* *	24	* *
Forsyth	2,499	68.7 (15)	833	30.3 (5)	1,095	27.4 (6)
Franklin	405	37.3 (75)	28	* *	37	* *
Gaston	1,682	36.7 (77)	388	14.7 (33)	644	9.6 (49)
Gates	57	* *	12	* *	17	* *
Graham	73	* *	27	* *	53	* *
Granville	419	30.5 (86)	93	* *	190	12.6 (31)

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this county's rate to other counties. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7B
2004 Pediatric Nutrition Surveillance (1)
North Carolina

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by County (2)
Children Aged < 5 Years (3)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by County (4,5)						
County	Breastfeeding (6)					
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)
Greene	216	51.4 (45)	99	* *	125	8.8 (54)
Guilford	2,433	55.8 (31)	217	17.1 (26)	362	13.8 (28)
Halifax	718	31.1 (83)	213	3.8 (50)	417	3.1 (71)
Harnett	990	37.3 (75)	234	16.7 (29)	356	7.3 (61)
Haywood	356	46.6 (54)	118	27.1 (7)	198	11.1 (41)
Henderson	582	73.5 (11)	175	37.1 (1)	291	23.4 (7)
Hertford	338	29.9 (87)	86	* *	172	1.2 (73)
Hoke	518	52.1 (42)	119	17.6 (23)	265	8.3 (56)
Hyde	48	* *	5	* *	12	* *
Iredell	875	41.5 (66)	351	18.2 (22)	541	15.0 (21)
Jackson	150	63.3 (20)	73	* *	92	* *
Johnston	857	35.7 (78)	275	14.5 (35)	463	12.7 (30)
Jones	102	43.1 (62)	2	* *	10	* *

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this county's rate to other counties. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7B
2004 Pediatric Nutrition Surveillance (1)
North Carolina

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by County (2)
Children Aged < 5 Years (3)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by County (4,5)						
County	Breastfeeding (6)					
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)
Lee	557	56.4 (27)	172	14.0 (36)	249	12.0 (34)
Lenoir	703	31.0 (84)	207	11.1 (45)	253	3.6 (69)
Lincoln	531	50.8 (47)	171	9.4 (47)	230	7.8 (60)
Macon	267	67.4 (16)	57	* *	126	7.1 (62)
Madison	133	94.0 (3)	55	* *	82	* *
Martin	261	32.6 (82)	77	* *	140	7.1 (62)
Mcdowell	310	53.5 (38)	118	19.5 (17)	163	12.3 (33)
Mecklenbur	3,913	54.7 (35)	1,595	21.4 (14)	2,471	18.4 (14)
Mitchell	126	66.7 (18)	38	* *	80	* *
Montgomery	299	55.9 (30)	104	22.1 (12)	121	10.7 (45)
Moore	567	45.3 (58)	194	15.5 (31)	293	14.0 (25)
Nash	944	38.7 (74)	206	12.6 (39)	419	9.8 (47)
New Hanove	1,289	42.6 (64)	341	17.3 (25)	505	11.9 (36)

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this county's rate to other counties. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7B
2004 Pediatric Nutrition Surveillance (1)
North Carolina

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by County (2)
Children Aged < 5 Years (3)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by County (4,5)						
County	Breastfeeding (6)					
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)
Northhampt	196	14.3 (91)	62	* *	115	4.3 (68)
Onslow	2,409	53.5 (38)	783	17.1 (26)	1,134	9.1 (50)
Orange	498	67.1 (17)	196	31.6 (4)	324	19.8 (12)
Pamlico	118	72.9 (12)	24	* *	45	* *
Pasquotank	374	23.5 (90)	124	9.7 (46)	222	9.0 (52)
Pender	394	34.5 (80)	132	3.0 (51)	189	1.1 (74)
Perquimans	77	* *	33	* *	42	* *
Person	307	39.4 (73)	92	* *	125	8.0 (58)
Pitt	1,089	46.2 (55)	424	12.3 (41)	560	9.8 (47)
Polk	113	53.1 (40)	39	* *	39	* *
Randolph	1,304	60.7 (25)	412	18.4 (21)	488	11.1 (41)
Richmond	541	41.2 (67)	217	9.2 (48)	303	8.3 (56)
Robeson	1,202	40.3 (68)	82	* *	146	11.6 (39)

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this county's rate to other counties. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7B
2004 Pediatric Nutrition Surveillance (1)
North Carolina

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by County (2)
Children Aged < 5 Years (3)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by County (4,5)						
County	Breastfeeding (6)					
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)
Rockingham	552	44.4 (60)	218	15.1 (32)	332	15.1 (20)
Rowan	1,047	53.6 (37)	357	17.6 (23)	513	11.3 (40)
Rutherford	461	49.5 (49)	66	* *	118	17.8 (15)
Sampson	716	42.3 (65)	295	22.7 (11)	485	15.7 (19)
Scotland	407	45.5 (57)	157	1.9 (52)	288	3.5 (70)
Stanly	438	71.5 (13)	132	5.3 (49)	213	6.1 (66)
Stokes	284	57.4 (26)	73	* *	152	9.9 (46)
Surry	491	55.8 (31)	226	23.0 (10)	287	22.0 (10)
Swain	92	* *	37	* *	55	* *
Transylvan	213	93.0 (4)	25	* *	31	* *
Tyrrell	64	* *	13	* *	37	* *
Union	982	80.8 (5)	299	18.7 (20)	620	14.0 (25)
Vance	579	61.0 (24)	135	35.6 (2)	291	38.5 (1)

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this county's rate to other counties. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7B
2004 Pediatric Nutrition Surveillance (1)
North Carolina

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by County (2)
Children Aged < 5 Years (3)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by County (4,5)						
County	Breastfeeding (6)					
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)
Wake	3,538	79.0 (6)	1,310	29.3 (6)	2,248	30.6 (3)
Warren	180	40.0 (69)	93	* *	146	6.2 (65)
Washington	165	40.0 (69)	58	* *	109	11.0 (44)
Watauga	247	61.1 (23)	66	* *	125	33.6 (2)
Wayne Coun	1,179	56.3 (28)	320	14.7 (33)	656	7.0 (64)
Wilkes	597	49.7 (48)	136	16.9 (28)	204	11.8 (37)
Wilson Cou	892	39.8 (72)	258	12.4 (40)	388	14.2 (24)
Yadkin	300	77.0 (7)	89	* *	154	22.1 (9)
Yancey	153	94.1 (2)	43	* *	92	* *
North Carolina	64,406	53.0	19,988	18.7	30,640	15.2
Nation (Prior Year)	928,626	53.2	490,960	21.5	504,421	13.6

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this county's rate to other counties. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6B
2002 - 2004 Pediatric Nutrition Surveillance (1)
North Carolina
Comparison of Growth and Anemia Indicators by County
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators by County (3,4)												
County	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)	Under- weight (7)	Over- weight (7)		≥ 2 Yrs Over- weight (7)				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Alamance	2,127	8.2 (9)	8.4 (83)	6,996	7.1 (84)	7.6 (46)	18.5 (93)	2,532	17.9 (83)	16.2 (67)	5,057	28.5 (99)
Alexander	706	8.4 (13)	7.4 (63)	1,750	16.6(100)	6.6 (23)	19.8 (97)	991	17.9 (83)	21.8 (97)	1,830	19.7 (88)
Alleghany	220	11.8 (85)	5.0 (9)	912	6.7 (75)	6.6 (23)	16.9 (75)	482	16.0 (51)	19.1 (90)	735	10.1 (46)
Anson	735	9.5 (37)	6.1 (20)	2,172	6.1 (67)	7.5 (45)	18.0 (88)	1,208	11.3 (7)	13.0 (29)	2,058	11.7 (56)
Ashe	501	9.2 (28)	5.4 (11)	1,977	5.8 (62)	6.8 (29)	15.3 (56)	1,145	11.9 (10)	12.8 (25)	1,809	11.3 (53)
Avery	235	10.6 (72)	7.7 (67)	591	9.0 (95)	5.9 (16)	17.4 (80)	387	18.3 (90)	17.8 (80)	457	4.2 (5)
Beaufort	1,038	9.6 (41)	6.5 (34)	3,505	4.9 (31)	6.8 (29)	15.0 (49)	1,779	15.7 (45)	17.0 (74)	3,225	12.5 (61)
Bertie	439	12.8 (95)	3.0 (2)	1,993	8.2 (91)	6.6 (23)	17.4 (80)	1,139	14.7 (27)	17.1 (77)	1,912	27.7 (98)
Bladen	882	10.4 (65)	6.3 (27)	3,153	3.6 (4)	10.5 (82)	14.1 (35)	1,352	15.2 (36)	14.1 (38)	2,187	7.0 (27)
Brunswick	1,587	9.3 (33)	7.0 (47)	5,229	6.9 (82)	7.2 (37)	14.1 (35)	2,505	16.0 (51)	14.9 (48)	4,034	5.4 (13)
Buncombe	3,399	9.7 (46)	8.1 (73)	10,034	6.2 (68)	5.2 (7)	15.6 (63)	5,143	16.6 (66)	14.9 (48)	9,358	11.8 (58)

- (1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this county's rate to other counties. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6B
2002 - 2004 Pediatric Nutrition Surveillance (1)
North Carolina
Comparison of Growth and Anemia Indicators by County
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators by County (3,4)												
County	Birthweight (5)			Height and Weight							Anemia Low Hb/Hct (8)	
	Low	High		Short Stature (6)	Under- weight (7)	Over- weight (7)		≥ 2 Yrs Over- weight (7)				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Burke	1,538	8.6 (17)	4.9 (8)	4,859	5.0 (34)	10.5 (82)	13.2 (28)	2,214	13.0 (15)	12.7 (24)	4,103	5.4 (13)
Cabarrus	1,827	9.5 (37)	7.0 (47)	6,343	5.3 (47)	12.6 (92)	12.9 (21)	2,669	16.1 (57)	13.9 (36)	3,859	6.6 (21)
Caldwell	1,808	10.9 (75)	7.1 (53)	4,925	7.1 (84)	8.3 (55)	15.1 (52)	2,461	14.8 (30)	13.5 (34)	4,494	6.1 (16)
Camden	147	11.6 (83)	4.1 (5)	579	6.7 (75)	14.3 (97)	10.5 (3)	285	6.0 (1)	4.9 (1)	368	12.5 (61)
Carteret	824	9.8 (51)	9.5 (98)	2,434	4.8 (27)	10.3 (81)	14.5 (42)	1,154	15.8 (48)	18.2 (85)	2,161	6.8 (25)
Caswell	449	11.6 (83)	6.7 (42)	1,387	5.6 (58)	8.6 (62)	12.7 (19)	690	12.2 (11)	9.9 (6)	1,090	6.6 (21)
Catawba	2,454	10.2 (58)	6.9 (43)	4,529	5.1 (38)	17.6 (99)	9.6 (2)	1,623	16.7 (68)	15.0 (51)	5,466	8.1 (36)
Chatham	1,106	8.5 (14)	7.0 (47)	2,937	6.8 (80)	6.6 (23)	18.1 (90)	1,280	15.9 (49)	19.7 (94)	2,463	12.6 (63)
Cherokee	607	9.9 (53)	6.6 (39)	1,166	5.3 (47)	12.3 (88)	14.4 (40)	413	18.4 (92)	19.6 (92)	1,576	8.4 (39)
Chowan	347	10.4 (65)	6.3 (27)	1,440	4.7 (23)	9.6 (76)	13.3 (30)	661	17.9 (83)	13.0 (29)	1,095	17.4 (79)
Clay	187	6.4 (2)	10.7 (100)	583	6.7 (75)	5.7 (13)	15.1 (52)	306	16.0 (51)	18.6 (88)	525	7.4 (30)

- (1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this county's rate to other counties. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6B
2002 - 2004 Pediatric Nutrition Surveillance ⁽¹⁾
North Carolina
Comparison of Growth and Anemia Indicators by County
Children Aged < 5 Years ⁽²⁾

Comparison of Growth and Anemia Indicators by County ^(3,4)												
County	Birthweight ⁽⁵⁾			Height and Weight						Anemia Low Hb/Hct ⁽⁸⁾		
	Low	High		Short Stature ⁽⁶⁾	Under- weight ⁽⁷⁾	Over- weight ⁽⁷⁾		≥ 2 Yrs Over- weight ⁽⁷⁾			Anemia Low Hb/Hct ⁽⁸⁾	
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Cleveland	1,632	9.2 (28)	8.1 (73)	1,814	2.8 (1)	13.6 (93)	14.8 (47)	163	15.3 (39)	12.3 (18)	352	7.7 (33)
Columbus	765	11.9 (86)	8.1 (73)	4,714	6.5 (72)	5.6 (11)	17.2 (78)	2,554	13.5 (21)	13.2 (31)	3,707	17.6 (80)
Craven	2,550	9.4 (35)	9.2 (96)	6,960	3.7 (7)	8.8 (67)	12.0 (14)	3,400	14.3 (24)	12.2 (17)	6,209	14.5 (72)
Cumberland	9,698	9.7 (46)	7.7 (67)	32,883	4.7 (23)	11.1 (86)	11.7 (9)	14,651	13.3 (18)	9.4 (3)	26,665	18.8 (83)
Currituck	367	8.7 (20)	8.7 (89)	1,273	5.5 (56)	6.9 (32)	15.4 (61)	542	15.3 (39)	14.4 (41)	902	9.8 (42)
Dare	362	9.1 (27)	7.2 (57)	1,137	7.1 (84)	4.9 (5)	17.7 (85)	585	17.9 (83)	16.6 (71)	1,003	11.3 (53)
Davidson	2,790	11.1 (76)	6.6 (39)	7,831	5.2 (42)	7.2 (37)	12.9 (21)	3,656	14.5 (25)	12.3 (18)	6,604	8.7 (40)
Davie	525	10.3 (61)	9.1 (94)	1,701	7.4 (87)	4.4 (2)	18.3 (92)	907	19.1 (96)	16.6 (71)	1,749	8.1 (36)
Duplin	1,786	9.3 (33)	5.0 (9)	3,887	4.0 (9)	14.7 (98)	15.3 (56)	1,669	14.5 (25)	18.0 (83)	1,780	27.2 (96)
Durham	3,715	9.5 (37)	6.4 (30)	8,313	6.7 (75)	8.7 (65)	14.4 (40)	3,700	11.7 (9)	10.5 (9)	1,175	12.8 (65)
Edgecombe	1,573	12.2 (90)	3.9 (4)	723	5.0 (34)	28.9(100)	7.1 (1)	65	* *	* *	195	11.8 (58)

- (1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this county's rate to other counties. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6B
2002 - 2004 Pediatric Nutrition Surveillance (1)
North Carolina
Comparison of Growth and Anemia Indicators by County
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators by County (3,4)												
County	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)	Under- weight (7)	Over- weight (7)		≥ 2 Yrs Over- weight (7)				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Forsyth	6,291	11.1 (76)	6.4 (30)	17,559	5.8 (62)	9.5 (74)	14.6 (44)	8,284	16.0 (51)	14.4 (41)	14,185	14.0 (70)
Franklin	968	9.0 (25)	8.4 (83)	2,808	4.7 (23)	7.7 (50)	15.1 (52)	1,629	18.0 (88)	15.7 (61)	2,338	6.4 (20)
Gaston	3,998	9.9 (53)	7.9 (70)	5,654	9.4 (96)	7.6 (46)	11.4 (8)	1,528	15.0 (34)	14.4 (41)	8,191	3.7 (3)
Gates	171	14.0 (98)	8.2 (76)	683	4.8 (27)	9.1 (70)	12.9 (21)	357	17.6 (81)	12.9 (26)	520	12.9 (66)
Graham	217	9.2 (28)	8.8 (90)	736	4.5 (19)	10.1 (80)	12.2 (16)	418	9.6 (4)	12.9 (26)	705	4.4 (7)
Granville	1,013	8.5 (14)	5.8 (14)	2,819	3.6 (4)	8.3 (55)	16.2 (71)	1,489	17.5 (78)	18.1 (84)	2,442	16.8 (77)
Greene	561	10.7 (73)	6.2 (25)	2,414	6.9 (82)	3.6 (1)	18.8 (95)	1,334	15.7 (45)	15.6 (59)	2,143	19.5 (86)
Guilford	5,356	10.4 (65)	6.1 (20)	12,679	5.4 (49)	8.9 (68)	11.7 (9)	8,374	13.4 (20)	12.9 (26)	9,722	6.2 (18)
Halifax	1,688	14.2 (100)	4.6 (7)	4,503	4.4 (14)	7.6 (46)	18.9 (96)	2,116	14.9 (32)	17.9 (82)	4,732	23.3 (92)
Harnett	2,440	9.7 (46)	7.8 (69)	7,507	5.8 (62)	7.0 (33)	15.6 (63)	3,762	16.7 (68)	14.3 (40)	6,253	13.0 (67)
Haywood	924	12.2 (90)	7.9 (70)	1,406	6.3 (70)	9.4 (73)	12.8 (20)	733	16.4 (62)	15.4 (55)	2,470	4.9 (9)

- (1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this county's rate to other counties. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6B
2002 - 2004 Pediatric Nutrition Surveillance ⁽¹⁾
North Carolina
Comparison of Growth and Anemia Indicators by County
Children Aged < 5 Years ⁽²⁾

Comparison of Growth and Anemia Indicators by County ^(3,4)												
County	Birthweight ⁽⁵⁾			Height and Weight							Anemia Low Hb/Hct ⁽⁸⁾	
	Low	High		Short Stature ⁽⁶⁾	Under- weight ⁽⁷⁾	Over- weight ⁽⁷⁾		≥ 2 Yrs Over- weight ⁽⁷⁾				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Henderson	1,541	8.8 (21)	7.6 (65)	3,357	5.0 (34)	8.6 (62)	13.7 (33)	1,578	16.0 (51)	14.8 (46)	3,955	19.0 (84)
Hertford	747	13.0 (96)	3.6 (3)	2,376	4.4 (14)	9.7 (77)	13.0 (25)	1,049	17.1 (72)	16.4 (69)	1,647	9.2 (41)
Hoke	1,284	11.4 (79)	7.2 (57)	5,294	10.3 (99)	7.4 (41)	20.1 (98)	2,538	16.5 (65)	16.1 (66)	3,515	26.9 (95)
Hyde	115	9.6 (41)	1.7 (1)	296	4.4 (14)	6.8 (29)	17.9 (87)	233	13.3 (18)	17.2 (78)	311	20.3 (89)
Iredell	2,215	9.8 (51)	6.6 (39)	8,037	5.4 (49)	7.4 (41)	14.3 (37)	3,388	15.1 (35)	12.5 (20)	6,042	3.7 (3)
Jackson	471	8.3 (10)	8.3 (79)	1,976	6.0 (65)	6.6 (23)	17.4 (80)	948	20.3 (99)	17.8 (80)	1,487	6.7 (23)
Johnston	1,888	10.1 (56)	6.1 (20)	4,840	5.7 (61)	7.3 (39)	14.9 (48)	2,238	16.2 (58)	16.0 (64)	5,050	10.3 (49)
Jones	237	12.2 (90)	8.9 (93)	636	3.6 (4)	8.6 (62)	14.6 (44)	388	15.5 (43)	16.0 (64)	572	11.2 (52)
Lee	1,437	8.6 (17)	5.8 (14)	4,478	5.1 (38)	6.2 (18)	20.7 (99)	1,980	20.2 (98)	22.3 (98)	3,420	27.6 (97)
Lenoir	1,633	10.4 (65)	5.9 (17)	5,241	3.9 (8)	11.2 (87)	13.0 (25)	2,321	13.1 (16)	11.2 (12)	3,862	10.4 (50)
Lincoln	1,235	7.9 (6)	6.9 (43)	1,240	4.8 (27)	9.8 (78)	13.5 (31)	545	14.9 (32)	10.5 (9)	2,720	4.3 (6)

- (1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this county's rate to other counties. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6B
2002 - 2004 Pediatric Nutrition Surveillance (1)
North Carolina
Comparison of Growth and Anemia Indicators by County
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators by County (3,4)												
County	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)	Under- weight (7)	Over- weight (7)		≥ 2 Yrs Over- weight (7)				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Macon	656	7.9 (6)	9.8 (99)	1,714	6.8 (80)	4.6 (3)	16.0 (70)	995	16.6 (66)	14.6 (44)	1,657	6.2 (18)
Madison	387	14.0 (98)	8.5 (85)	1,586	8.2 (91)	6.2 (18)	17.8 (86)	769	17.0 (71)	18.6 (88)	1,228	10.9 (51)
Martin	659	12.7 (94)	6.4 (30)	2,323	5.1 (38)	8.2 (54)	15.8 (68)	1,060	15.2 (36)	15.0 (51)	1,697	20.5 (90)
Mcdowell	810	10.4 (65)	6.9 (43)	3,152	4.8 (27)	7.6 (46)	13.8 (34)	1,594	14.8 (30)	13.9 (36)	2,442	15.1 (74)
Mecklenbur	9,938	9.6 (41)	7.2 (57)	26,348	6.7 (75)	8.1 (52)	15.0 (49)	11,850	16.2 (58)	16.2 (67)	17,551	6.9 (26)
Mitchell	327	6.4 (2)	7.0 (47)	1,182	10.0 (98)	6.3 (20)	12.9 (21)	682	12.9 (14)	12.6 (22)	1,107	3.3 (2)
Montgomery	803	10.2 (58)	7.1 (53)	3,064	7.5 (90)	9.1 (70)	18.0 (88)	1,376	17.2 (74)	18.3 (86)	2,315	5.9 (15)
Moore	1,415	9.0 (25)	7.3 (62)	4,949	5.4 (49)	7.0 (33)	15.8 (68)	2,531	17.1 (72)	14.2 (39)	4,549	10.1 (46)
Nash	2,330	8.8 (21)	7.1 (53)	6,420	4.9 (31)	10.5 (82)	13.2 (28)	2,611	15.4 (42)	15.7 (61)	4,323	8.2 (38)
New Hanove	2,932	12.1 (89)	7.1 (53)	5,635	7.4 (87)	5.3 (8)	15.3 (56)	2,538	16.3 (60)	14.9 (48)	4,355	7.1 (29)
Northhampt	520	13.7 (97)	5.8 (14)	2,193	4.5 (19)	7.4 (41)	15.6 (63)	1,186	16.4 (62)	15.5 (57)	1,751	23.6 (93)

- (1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this county's rate to other counties. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6B
2002 - 2004 Pediatric Nutrition Surveillance ⁽¹⁾
North Carolina
Comparison of Growth and Anemia Indicators by County
Children Aged < 5 Years ⁽²⁾

Comparison of Growth and Anemia Indicators by County ^(3,4)												
County	Birthweight ⁽⁵⁾			Height and Weight							Anemia Low Hb/Hct ⁽⁸⁾	
	Low	High		Short Stature ⁽⁶⁾	Under- weight ⁽⁷⁾	Over- weight ⁽⁷⁾		≥ 2 Yrs Over- weight ⁽⁷⁾			Anemia Low Hb/Hct ⁽⁸⁾	
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Onslow	6,298	8.5 (14)	7.9 (70)	18,824	4.4 (14)	9.0 (69)	10.5 (3)	8,107	13.1 (16)	9.5 (5)	14,308	4.9 (9)
Orange	1,190	9.7 (46)	9.2 (96)	4,379	6.6 (73)	7.1 (36)	17.5 (83)	2,148	17.4 (77)	20.0 (95)	3,630	14.2 (71)
Pamlico	255	5.9 (1)	7.5 (64)	997	4.4 (14)	8.3 (55)	15.5 (62)	501	17.2 (74)	20.4 (96)	549	17.3 (78)
Pasquotank	941	9.6 (41)	5.4 (11)	2,945	4.2 (11)	8.5 (60)	13.0 (25)	1,322	12.3 (12)	10.4 (8)	2,007	9.9 (45)
Pender	969	9.7 (46)	8.3 (79)	3,810	5.4 (49)	8.7 (65)	16.2 (71)	1,796	11.4 (8)	12.5 (20)	2,789	9.8 (42)
Perquimans	209	12.0 (88)	7.2 (57)	846	8.9 (94)	8.5 (60)	21.3(100)	361	14.7 (27)	15.2 (53)	621	13.0 (67)
Person	792	12.2 (90)	9.1 (94)	2,648	4.0 (9)	7.4 (41)	15.3 (56)	1,224	15.2 (36)	15.4 (55)	2,105	7.4 (30)
Pitt	2,807	11.4 (79)	5.9 (17)	10,590	5.2 (42)	5.4 (10)	18.1 (90)	5,390	18.6 (94)	19.6 (92)	8,883	19.1 (85)
Polk	258	7.4 (4)	8.5 (85)	779	4.2 (11)	5.3 (8)	16.7 (74)	412	17.5 (78)	17.0 (74)	756	19.6 (87)
Randolph	3,011	9.2 (28)	8.6 (88)	7,174	5.4 (49)	8.3 (55)	17.3 (79)	3,228	19.4 (97)	22.9 (99)	7,404	5.1 (12)
Richmond	1,341	10.0 (55)	7.6 (65)	5,372	3.4 (2)	14.2 (96)	11.0 (7)	2,933	8.0 (2)	8.5 (2)	4,917	33.1(100)

- (1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this county's rate to other counties. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6B
2002 - 2004 Pediatric Nutrition Surveillance (1)
North Carolina
Comparison of Growth and Anemia Indicators by County
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators by County (3,4)												
County	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)	Under- weight (7)	Over- weight (7)		≥ 2 Yrs Over- weight (7)				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Robeson	3,171	10.3 (61)	7.0 (47)	7,966	5.4 (49)	9.5 (74)	14.6 (44)	5,499	13.5 (21)	13.3 (32)	6,700	4.9 (9)
Rockingham	1,434	10.1 (56)	6.9 (43)	5,502	4.2 (11)	12.4 (90)	11.8 (13)	2,427	11.1 (6)	10.3 (7)	4,338	9.8 (42)
Rowan	2,662	11.4 (79)	5.7 (13)	7,155	4.5 (19)	5.9 (16)	17.0 (76)	3,511	17.2 (74)	17.4 (79)	6,222	11.4 (55)
Rutherford	1,177	10.3 (61)	6.1 (20)	4,341	5.2 (42)	5.7 (13)	15.7 (66)	2,191	16.0 (51)	13.3 (32)	3,561	13.7 (69)
Sampson	1,819	8.9 (24)	8.2 (76)	6,199	5.4 (49)	10.7 (85)	16.2 (71)	2,985	15.9 (49)	15.5 (57)	5,268	18.3 (82)
Scotland	1,156	11.2 (78)	6.5 (34)	3,619	4.7 (23)	6.7 (28)	14.3 (37)	2,101	15.6 (44)	13.7 (35)	3,425	15.3 (75)
Stanly	1,147	11.9 (86)	6.5 (34)	2,934	5.5 (56)	7.3 (39)	14.3 (37)	1,573	12.6 (13)	11.3 (13)	2,864	25.2 (94)
Stokes	730	10.7 (73)	5.9 (17)	2,782	5.2 (42)	12.5 (91)	10.6 (5)	1,329	14.7 (27)	12.6 (22)	1,822	4.6 (8)
Surry	1,336	8.8 (21)	8.3 (79)	2,958	6.0 (65)	13.9 (94)	10.6 (5)	1,099	9.1 (3)	9.4 (3)	3,838	2.6 (1)
Swain	206	9.2 (28)	8.3 (79)	918	5.6 (58)	4.8 (4)	11.7 (9)	552	13.9 (23)	11.4 (15)	884	7.7 (33)
Transylvan	561	8.0 (8)	7.0 (47)	1,463	4.6 (22)	12.3 (88)	12.2 (16)	754	17.6 (81)	14.7 (45)	1,050	6.1 (16)

- (1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this county's rate to other counties. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6B
2002 - 2004 Pediatric Nutrition Surveillance ⁽¹⁾
North Carolina
Comparison of Growth and Anemia Indicators by County
Children Aged < 5 Years ⁽²⁾

Comparison of Growth and Anemia Indicators by County ^(3,4)												
County	Birthweight ⁽⁵⁾			Height and Weight						Anemia Low Hb/Hct ⁽⁸⁾		
	Low	High		Short Stature ⁽⁶⁾	Under- weight ⁽⁷⁾	Over- weight ⁽⁷⁾		≥ 2 Yrs Over- weight ⁽⁷⁾			Anemia Low Hb/Hct ⁽⁸⁾	
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Tyrrell	153	7.8 (5)	8.5 (85)	656	8.7 (93)	4.9 (5)	15.7 (66)	359	16.4 (62)	15.9 (63)	463	17.7 (81)
Union	2,236	8.3 (10)	6.4 (30)	7,280	6.6 (73)	6.3 (20)	15.1 (52)	3,599	16.9 (70)	15.2 (53)	6,456	12.7 (64)
Vance	1,432	10.5 (70)	6.1 (20)	4,469	5.1 (38)	8.4 (59)	15.0 (49)	2,290	16.3 (60)	16.6 (71)	3,459	16.1 (76)
Wake	8,945	8.3 (10)	8.2 (76)	33,066	5.2 (42)	9.1 (70)	13.6 (32)	13,957	15.7 (45)	14.8 (46)	22,841	6.7 (23)
Warren	507	11.4 (79)	4.3 (6)	2,420	6.2 (68)	8.0 (51)	12.2 (16)	1,240	15.3 (39)	11.5 (16)	1,952	10.1 (46)
Washington	384	9.4 (35)	6.5 (34)	1,588	5.0 (34)	9.9 (79)	14.5 (42)	905	10.2 (5)	10.9 (11)	1,378	21.8 (91)
Watauga	540	10.2 (58)	6.3 (27)	1,993	3.5 (3)	8.1 (52)	12.1 (15)	1,047	17.9 (83)	11.3 (13)	1,482	11.9 (60)
Wayne Coun	2,867	10.3 (61)	8.8 (90)	8,086	4.9 (31)	14.0 (95)	11.7 (9)	3,643	18.3 (90)	15.6 (59)	6,986	11.7 (56)
Wilkes	1,548	9.6 (41)	7.2 (57)	3,560	6.3 (70)	6.3 (20)	15.3 (56)	1,990	18.7 (95)	16.4 (69)	3,236	7.4 (30)
Wilson Cou	2,127	10.5 (70)	6.2 (25)	7,527	5.6 (58)	7.0 (33)	18.5 (93)	3,349	18.0 (88)	18.5 (87)	6,011	7.7 (33)
Yadkin	703	9.5 (37)	8.8 (90)	2,040	7.4 (87)	5.8 (15)	17.1 (77)	1,036	17.5 (78)	17.0 (74)	2,052	14.5 (72)

- (1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this county's rate to other counties. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6B
2002 - 2004 Pediatric Nutrition Surveillance (1)
North Carolina
Comparison of Growth and Anemia Indicators by County
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators by County (3,4)												
County	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)	Under- weight (7)	Over- weight (7)		≥ 2 Yrs Over- weight (7)				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- (Rank)	% <95th (Rank)	% ≥95th (Rank)	Number
Yancey	337	8.6 (17)	6.5 (34)	1,369	9.5 (97)	5.6 (11)	17.6 (84)	835	18.4 (92)	19.2 (91)	1,290	7.0 (27)
North Carolina	161,382	9.9	7.1	474,090	5.6	8.7	14.4	224,426	15.3	14.4	385,007	11.7

- (1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this county's rate to other counties. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7B
2002 - 2004 Pediatric Nutrition Surveillance ⁽¹⁾
North Carolina

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by County ⁽²⁾
Children Aged < 5 Years ⁽³⁾

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by County ^(4,5)						
County	Breastfeeding ⁽⁶⁾					
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)
Alamance	3,275	50.3 (54)	1,904	21.7 (23)	1,601	17.7 (20)
Alexander	1,025	43.6 (66)	531	15.4 (50)	535	8.8 (70)
Alleghany	312	64.7 (21)	181	6.1 (87)	187	4.3 (85)
Anson	1,004	35.9 (84)	555	7.4 (82)	605	4.5 (84)
Ashe	771	52.1 (47)	477	14.9 (53)	445	9.9 (61)
Avery	288	68.1 (17)	52	* *	50	* *
Beaufort	1,589	46.8 (60)	992	12.8 (71)	973	8.7 (74)
Bertie	704	29.4 (91)	514	6.2 (86)	493	4.3 (85)
Bladen	1,275	45.4 (64)	733	16.5 (43)	685	16.6 (25)
Brunswick	2,265	43.0 (69)	1,166	15.6 (49)	1,179	12.6 (44)
Buncombe	4,969	82.3 (6)	2,865	24.4 (13)	2,829	15.8 (26)
Burke	2,338	50.7 (52)	1,318	13.7 (68)	1,309	9.0 (67)
Cabarrus	2,589	61.5 (23)	1,496	29.1 (9)	1,721	20.7 (13)

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this county's rate to other counties. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7B
2002 - 2004 Pediatric Nutrition Surveillance ⁽¹⁾
North Carolina

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by County ⁽²⁾
Children Aged < 5 Years ⁽³⁾

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by County ^(4,5)						
County	Breastfeeding ⁽⁶⁾					
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)
Caldwell	2,556	58.6 (26)	1,289	12.7 (72)	1,316	9.0 (67)
Camden	222	19.8 (99)	124	1.6 (95)	138	2.2 (95)
Carteret	1,150	50.8 (51)	536	14.0 (65)	514	8.4 (75)
Caswell	604	43.2 (68)	314	7.6 (81)	354	4.0 (89)
Catawba	3,390	56.9 (30)	1,957	18.5 (35)	1,733	14.1 (35)
Chatham	1,701	73.8 (11)	884	36.8 (2)	823	25.2 (6)
Cherokee	836	52.9 (44)	479	18.2 (37)	448	12.1 (48)
Chowan	530	22.3 (98)	307	4.2 (91)	310	2.6 (92)
Clay	280	51.8 (48)	151	17.2 (39)	165	17.0 (21)
Cleveland	1,683	25.0 (96)	166	14.5 (59)	157	8.9 (69)
Columbus	1,427	22.4 (97)	1,157	3.6 (93)	1,127	2.8 (91)
Craven	3,547	59.4 (24)	1,981	19.6 (29)	1,790	14.8 (29)
Cumberland	14,906	53.1 (43)	8,192	14.0 (65)	7,881	10.5 (57)

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this county's rate to other counties. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7B
2002 - 2004 Pediatric Nutrition Surveillance ⁽¹⁾
North Carolina

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by County ⁽²⁾
Children Aged < 5 Years ⁽³⁾

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by County ^(4,5)						
County	Breastfeeding ⁽⁶⁾					
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)
Currituck	528	75.8 (7)	293	1.4 (96)	289	1.0 (97)
Dare	551	89.8 (2)	308	26.0 (11)	282	19.1 (16)
Davidson	4,066	49.9 (55)	2,185	15.9 (47)	1,934	12.6 (44)
Davie	817	58.4 (27)	517	22.8 (16)	480	21.9 (12)
Duplin	2,195	53.9 (39)	691	16.9 (40)	658	14.4 (31)
Durham	5,819	54.9 (37)	4,190	25.5 (12)	3,898	20.4 (14)
Edgecombe	1,609	25.5 (93)	92	* *	91	* *
Forsyth	9,161	67.1 (19)	5,018	30.6 (8)	4,505	24.0 (8)
Franklin	1,100	36.2 (83)	228	18.4 (36)	208	16.8 (23)
Gaston	5,445	39.4 (77)	2,469	14.2 (62)	2,437	10.1 (60)
Gates	259	25.1 (95)	129	4.7 (89)	152	3.9 (90)
Graham	317	47.9 (58)	191	20.4 (27)	180	22.2 (11)
Granville	1,273	35.8 (85)	519	13.7 (68)	563	12.1 (48)

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this county's rate to other counties. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7B
2002 - 2004 Pediatric Nutrition Surveillance ⁽¹⁾
North Carolina

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by County ⁽²⁾
Children Aged < 5 Years ⁽³⁾

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by County ^(4,5)						
County	Breastfeeding ⁽⁶⁾					
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)
Greene	849	51.1 (50)	601	14.0 (65)	568	6.3 (80)
Guilford	6,414	54.7 (38)	1,577	21.4 (24)	1,411	14.6 (30)
Halifax	2,571	32.4 (89)	1,382	7.1 (83)	1,557	4.3 (85)
Harnett	3,361	36.4 (82)	1,672	14.6 (57)	1,640	8.8 (70)
Haywood	1,306	53.9 (39)	652	21.9 (20)	672	12.4 (47)
Henderson	2,322	75.6 (8)	1,288	32.2 (6)	1,229	26.4 (4)
Hertford	1,024	31.3 (90)	511	4.3 (90)	571	2.5 (93)
Hoke	1,815	50.5 (53)	958	16.6 (42)	1,130	8.8 (70)
Hyde	134	53.7 (41)	36	* *	40	* *
Iredell	3,263	42.7 (71)	2,000	16.1 (46)	1,948	13.1 (41)
Jackson	684	57.5 (29)	447	22.1 (18)	424	14.9 (28)
Johnston	3,051	40.5 (75)	1,732	15.4 (50)	1,677	13.7 (37)
Jones	275	43.6 (66)	89	* *	109	11.9 (50)

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this county's rate to other counties. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7B
2002 - 2004 Pediatric Nutrition Surveillance ⁽¹⁾
North Carolina

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by County ⁽²⁾
Children Aged < 5 Years ⁽³⁾

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by County ^(4,5)						
County	Breastfeeding ⁽⁶⁾					
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)
Lee	2,037	52.5 (45)	1,081	18.7 (34)	1,058	13.7 (37)
Lenoir	2,372	34.1 (87)	1,225	10.4 (79)	1,095	5.8 (82)
Lincoln	1,716	49.8 (56)	795	10.9 (76)	732	7.5 (76)
Macon	931	70.1 (16)	475	22.1 (18)	497	12.5 (46)
Madison	589	89.3 (3)	362	23.5 (15)	347	18.4 (18)
Martin	945	37.1 (81)	528	11.6 (74)	539	7.1 (78)
Mcdowell	1,200	55.4 (33)	691	19.0 (32)	695	14.1 (35)
Mecklenbur	15,541	62.2 (22)	9,435	21.2 (25)	8,915	16.9 (22)
Mitchell	482	66.2 (20)	280	27.9 (10)	294	22.8 (10)
Montgomery	1,142	55.2 (35)	604	19.5 (30)	591	13.7 (37)
Moore	2,129	45.7 (63)	1,187	14.1 (63)	1,178	11.9 (50)
Nash	3,076	42.3 (72)	1,381	13.0 (70)	1,524	9.3 (63)
New Hanove	4,216	43.0 (69)	1,979	16.5 (43)	1,895	10.6 (56)

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this county's rate to other counties. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7B
2002 - 2004 Pediatric Nutrition Surveillance ⁽¹⁾
North Carolina

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by County ⁽²⁾
Children Aged < 5 Years ⁽³⁾

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by County ^(4,5)						
County	Breastfeeding ⁽⁶⁾					
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)
Northhampt	749	16.6(100)	396	4.0 (92)	451	2.4 (94)
Onslow	8,879	53.6 (42)	4,866	14.7 (56)	4,614	9.3 (63)
Orange	1,985	70.9 (15)	1,165	33.6 (4)	1,141	25.3 (5)
Pamlico	368	73.6 (12)	205	14.6 (57)	197	9.6 (62)
Pasquotank	1,354	27.0 (92)	742	7.0 (84)	783	6.0 (81)
Pender	1,445	33.8 (88)	873	2.9 (94)	845	1.5 (96)
Perquimans	329	25.2 (94)	202	11.9 (73)	194	10.3 (58)
Person	1,059	39.4 (77)	558	10.9 (76)	587	7.2 (77)
Pitt	3,999	46.6 (61)	2,473	14.4 (61)	2,370	10.8 (55)
Polk	379	55.1 (36)	228	23.7 (14)	186	16.7 (24)
Randolph	4,358	67.3 (18)	2,348	18.8 (33)	1,909	10.3 (58)
Richmond	2,072	42.2 (73)	1,266	11.1 (75)	1,205	8.8 (70)
Robeson	3,429	38.1 (79)	536	16.4 (45)	489	11.5 (54)

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this county's rate to other counties. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7B
2002 - 2004 Pediatric Nutrition Surveillance ⁽¹⁾
North Carolina

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by County ⁽²⁾
Children Aged < 5 Years ⁽³⁾

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by County ^(4,5)						
County	Breastfeeding ⁽⁶⁾					
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)
Rockingham	2,135	45.3 (65)	1,411	16.9 (40)	1,298	12.8 (43)
Rowan	3,718	52.3 (46)	1,979	17.7 (38)	1,910	11.6 (53)
Rutherford	1,635	47.9 (58)	767	15.0 (52)	748	11.9 (50)
Sampson	2,641	45.9 (62)	1,612	20.8 (26)	1,646	18.0 (19)
Scotland	1,648	41.0 (74)	927	5.5 (88)	996	5.1 (83)
Stanly	1,660	72.3 (14)	928	6.3 (85)	907	4.2 (88)
Stokes	1,036	55.3 (34)	540	14.1 (63)	596	9.1 (66)
Surry	1,980	58.0 (28)	1,230	20.2 (28)	1,144	18.5 (17)
Swain	366	98.6 (1)	225	21.8 (21)	226	23.9 (9)
Transylvan	675	86.4 (4)	197	19.3 (31)	180	14.4 (31)
Tyrrell	212	56.6 (32)	113	10.6 (78)	132	12.9 (42)
Union	3,448	75.2 (9)	1,945	21.8 (21)	2,048	19.4 (15)
Vance	1,982	56.8 (31)	983	41.4 (1)	1,139	39.3 (1)

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this county's rate to other counties. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7B
2002 - 2004 Pediatric Nutrition Surveillance ⁽¹⁾
North Carolina

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by County ⁽²⁾
Children Aged < 5 Years ⁽³⁾

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by County ^(4,5)						
County	Breastfeeding ⁽⁶⁾					
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)
Wake	12,920	74.7 (10)	7,580	30.7 (7)	7,673	29.7 (3)
Warren	758	35.1 (86)	499	14.8 (55)	513	14.2 (34)
Washington	562	39.5 (76)	337	8.3 (80)	395	6.6 (79)
Watauga	805	59.0 (25)	452	35.2 (3)	461	35.6 (2)
Wayne Coun	4,030	51.2 (49)	1,973	14.5 (59)	2,174	9.3 (63)
Wilkes	2,027	48.1 (57)	926	14.9 (53)	882	13.3 (40)
Wilson Cou	3,066	38.0 (80)	1,766	15.8 (48)	1,695	14.3 (33)
Yadkin	1,007	72.9 (13)	558	22.2 (17)	598	15.1 (27)
Yancey	486	84.2 (5)	297	32.3 (5)	351	24.5 (7)
North Carolina	231,023	53.7	123,222	18.7	120,064	14.5

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this county's rate to other counties. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.