

Trend Data for the WIC Agreement Addenda SFY 2013-2014

The overall purpose of trend data is to track patterns or changes in data over time. For purposes of the WIC Agreement Addenda, trend data is being used to track the performance of the State and Local WIC Programs with respect to seventeen identified program indicators and sub-indicators. The goal for both the State and Local WIC Programs is to implement or enhance activities which improve performance over time for each indicator.

Attached are the reports of WIC Program Quality Assurance Deliverables data for each of the WIC Local Agency along with the North Carolina state wide data. It may be useful to compare an agency's performance with respect to the state as a whole. The counties representing the health districts are presented starting on page 88 of this document.

To compare one local agency data with other agencies' data, please refer to the indicator-specific reports by local agency posted on the website <http://www.nutritionnc.com/wic/wicLAR.htm>

Each report includes the following information.

❖ Process Outcome Objectives

This column includes a description of the Process Outcome Objectives (POO) data indicators of the WIC Program Quality Assurance Deliverable being assessed.

❖ Trend Data

The data displayed in these columns represent nine years of data divided into 3-year time periods (2003 to 2005, 2006 to 2008, and 2009 to 2011*) with the average of each three year time period shown.

- POO 1A, 1B, 1C, 4A, and 4B are calendar year (Jan – Dec) data.
- POO 2A, 2B, 3A, 3B, and 5A, are fiscal year (July – June) data, six months more recent than calendar year data.
- POO 5B is fiscal year (July – June) data which is twelve months prior to other fiscal year data.

❖ % Change from 2003-2011

This column displays the percent change from the first three year period to the most recent three-year time period along with the direction of the change (i.e., increase, decrease or no change).

❖ CY2011 or FY2012 Data

This column displays the most recent year's data including the numerator (Num) and denominator (Den) counts and the percent. Percentages calculated based on numbers less than 20 can show wide fluctuations from one year to the next. Therefore numbers below 20 in the denominator should be used with caution as the computed rates using small numbers are very unreliable.

❖ Note for POO 3A and 3B (Child Health data)

The FY2012 data used for Child Health for POO indicators (3A and 3B) are not currently available because of changes in the North Carolina's Health Information System (HIS). However, data for preceding fiscal years (2004 to 2009) were available for analysis from the old HSIS and have been presented in three year time intervals. *Please note that the FY2010 to FY2012 trend does not include the FY2011 and FY2012 data for these two indicators and as such rates were derived by using only the first six months of FY2010 time period (7/1/09 to 12/31/09).

SFY 2013-2014 AGREEMENT ADDENDA DATA
ATTACHMENT A: WIC QUALITY ASSURANCE DELIVERABLES
North Carolina

Process Outcome Objectives	2003 to 2005	2006 to 2008	2009 to 2011	% Change from 2003 to 2011	2011 Data		
					Num	Den	Percent
1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2)	71.8%	71.0%	70.7%	-1.6%	38,242	55,427	69.0%
1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services.	71.2%	71.6%	71.2%	0.0%	52,675	73,962	71.2%
1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services.	54.9%	52.8%	55.6%	1.2%	179,911	312,180	57.6%
2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy.	31.2%	29.5%	29.3%	-6.0%	39,840	134,973	29.5%
2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum.	66.3%	68.8%	70.6%	6.4%	47,119	65,205	72.3%
3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1)	86.4%	85.5%	87.8%	1.6%			
3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1)	79.4%	78.2%	80.4%	1.2%			
4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were:							
a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1)	65.7%	65.1%	64.0%	-2.5%	66,293	103,716	63.9%
b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile)	15.4%	15.9%	16.1%	4.4%	16,853	103,716	16.2%
c. Obese (BMI at or greater than the 95th percentile).	14.6%	15.3%	15.6%	7.0%	16,240	103,716	15.7%
d. Underweight (BMI less than the 5th percentile).	4.3%	3.7%	4.3%	-0.9%	4,330	103,716	4.2%
4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3)							
a. Recommended prenatal weight gain	31.7%	31.5%	30.1%	-5.2%	17,266	61,090	28.3%
b. Excessive prenatal weight gain	46.4%	46.0%	47.0%	1.4%	28,848	61,090	47.2%
c. Inadequate prenatal weight gain	21.9%	22.5%	22.9%	4.5%	14,976	61,090	24.5%
5A. Percentage of women who participated in WIC who initiated breastfeeding.	49.7%	53.9%	55.8%	12.3%	47,333	82,139	57.6%
5B. Percentage of infants who participated in WIC who were:							
a. Breastfeeding at 6 weeks of age (1)	28.5%	32.0%	32.8%	15.3%	27,714	82,385	33.6%
b. Breastfeeding at 6 months of age	15.0%	16.7%	17.7%	18.5%	15,210	82,385	18.5%

(1) Item appears also on other WCH Section Agreement Addenda.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI >=30.0): Adapted from the National Academy of Sciences. Weight Gain During Pregnancy: Reexamining the Guidelines. Washington, DC : National Academy Press, 2009.

SFY 2013-2014 AGREEMENT ADDENDA DATA

ATTACHMENT A: WIC QUALITY ASSURANCE DELIVERABLES

Alamance County

Process Outcome Objectives	2003 to 2005	2006 to 2008	2009 to 2011	% Change from 2003 to 2011	2011 Data		
					Num	Den	Percent
1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2)	76.0%	76.6%	75.4%	-0.7%	636	872	72.9%
1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services.	73.9%	76.0%	73.2%	-0.9%	842	1,158	72.7%
1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services.	51.9%	54.0%	58.4%	12.5%	3,092	5,207	59.4%
2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy.	39.3%	42.6%	38.6%	-1.8%	722	2,042	35.4%
2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum.	37.3%	63.6%	77.9%	109.0%	842	1,090	77.2%
3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1)	79.5%	48.1%					
3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1)	64.3%	55.8%	41.7%	-35.2%			
4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were:							
a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1)	62.5%	64.1%	62.7%	0.4%	1,187	1,954	60.7%
b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile)	18.6%	17.9%	17.8%	-4.8%	372	1,954	19.0%
c. Obese (BMI at or greater than the 95th percentile).	16.7%	15.9%	16.7%	-0.2%	331	1,954	16.9%
d. Underweight (BMI less than the 5th percentile).	2.2%	2.1%	2.9%	32.7%	64	1,954	3.3%
4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3)							
a. Recommended prenatal weight gain	33.6%	32.7%	31.6%	-6.0%	272	978	27.8%
b. Excessive prenatal weight gain	47.6%	45.5%	45.5%	-4.4%	478	978	48.9%
c. Inadequate prenatal weight gain	18.9%	21.8%	23.0%	21.7%	228	978	23.3%
5A. Percentage of women who participated in WIC who initiated breastfeeding.	46.6%	48.4%	55.9%	19.8%	762	1,278	59.6%
5B. Percentage of infants who participated in WIC who were:							
a. Breastfeeding at 6 weeks of age (1)	30.6%	31.9%	34.7%	13.6%	456	1,244	36.7%
b. Breastfeeding at 6 months of age	16.2%	17.9%	19.6%	21.2%	250	1,244	20.1%

(1) Item appears also on other WCH Section Agreement Addenda.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI >=30.0): Adapted from the National Academy of Sciences. Weight Gain During Pregnancy: Reexamining the Guidelines. Washington, DC : National Academy Press, 2009.

SFY 2013-2014 AGREEMENT ADDENDA DATA

ATTACHMENT A: WIC QUALITY ASSURANCE DELIVERABLES

Albemarle Regional Health Services District

Process Outcome Objectives	2003 to 2005	2006 to 2008	2009 to 2011	% Change from 2003 to 2011	2011 Data		
					Num	Den	Percent
1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2)	80.9%	72.3%	69.4%	-14.2%	495	780	63.5%
1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services.	75.3%	72.8%	73.9%	-1.8%	691	922	74.9%
1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services.	61.4%	56.7%	56.9%	-7.3%	2,345	3,961	59.2%
2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy.	38.8%	33.7%	28.0%	-27.9%	524	2,039	25.7%
2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum.	37.6%	37.5%	46.5%	23.7%	397	833	47.7%
3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1)	72.9%	67.5%					
3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1)	72.5%	69.8%					
4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were:							
a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1)	65.2%	64.8%	63.9%	-2.1%	940	1,496	62.8%
b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile)	14.8%	14.6%	14.5%	-2.2%	211	1,496	14.1%
c. Obese (BMI at or greater than the 95th percentile).	14.5%	16.5%	15.6%	7.0%	239	1,496	16.0%
d. Underweight (BMI less than the 5th percentile).	5.4%	4.1%	6.1%	11.9%	106	1,496	7.1%
4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3)							
a. Recommended prenatal weight gain	32.2%	31.8%	28.3%	-12.1%	151	632	23.9%
b. Excessive prenatal weight gain	43.7%	44.6%	41.3%	-5.5%	260	632	41.1%
c. Inadequate prenatal weight gain	24.1%	23.5%	30.4%	26.2%	221	632	35.0%
5A. Percentage of women who participated in WIC who initiated breastfeeding.	25.3%	20.4%	28.1%	10.8%	372	1,073	34.7%
5B. Percentage of infants who participated in WIC who were:							
a. Breastfeeding at 6 weeks of age (1)	9.6%	9.4%	12.4%	28.4%	160	1,021	15.7%
b. Breastfeeding at 6 months of age	3.8%	3.3%	5.1%	33.9%	90	1,021	8.8%

(1) Item appears also on other WCH Section Agreement Addenda.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI >=30.0): Adapted from the National Academy of Sciences. Weight Gain During Pregnancy: Reexamining the Guidelines. Washington, DC : National Academy Press, 2009.

SFY 2013-2014 AGREEMENT ADDENDA DATA

ATTACHMENT A: WIC QUALITY ASSURANCE DELIVERABLES

Alexander County

Process Outcome Objectives	2003 to 2005	2006 to 2008	2009 to 2011	% Change from 2003 to 2011	2011 Data		
					Num	Den	Percent
1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2)	74.3%	71.5%	75.1%	1.1%	138	186	74.2%
1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services.	79.1%	76.4%	75.1%	-5.1%	160	214	74.8%
1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services.	63.5%	59.6%	56.5%	-11.0%	614	1,044	58.8%
2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy.	44.5%	37.0%	37.7%	-15.2%	145	370	39.2%
2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum.	28.8%	76.7%	78.6%	172.8%	169	218	77.5%
3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1)	96.0%	98.3%	86.7%	-9.7%			
3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1)	82.8%	90.0%	95.1%	14.8%			
4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were:							
a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1)	61.2%	68.1%	65.3%	6.8%	143	224	63.8%
b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile)	17.3%	15.1%	15.8%	-8.7%	39	224	17.4%
c. Obese (BMI at or greater than the 95th percentile).	17.1%	13.8%	13.7%	-20.0%	29	224	12.9%
d. Underweight (BMI less than the 5th percentile).	4.5%	3.1%	5.3%	16.9%	13	224	5.8%
4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3)							
a. Recommended prenatal weight gain	27.0%	31.5%	31.0%	14.7%	68	189	36.0%
b. Excessive prenatal weight gain	47.7%	47.4%	48.8%	2.4%	87	189	46.0%
c. Inadequate prenatal weight gain	25.4%	21.2%	20.3%	-20.1%	34	189	18.0%
5A. Percentage of women who participated in WIC who initiated breastfeeding.	30.3%	46.9%	85.5%	182.5%	240	276	87.0%
5B. Percentage of infants who participated in WIC who were:							
a. Breastfeeding at 6 weeks of age (1)	26.9%	28.1%	29.0%	7.6%	89	289	30.8%
b. Breastfeeding at 6 months of age	15.7%	16.6%	17.3%	10.7%	55	289	19.0%

(1) Item appears also on other WCH Section Agreement Addenda.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI >=30.0): Adapted from the National Academy of Sciences. Weight Gain During Pregnancy: Reexamining the Guidelines. Washington, DC : National Academy Press, 2009.

SFY 2013-2014 AGREEMENT ADDENDA DATA

ATTACHMENT A: WIC QUALITY ASSURANCE DELIVERABLES

Anson County

Process Outcome Objectives	2003 to 2005	2006 to 2008	2009 to 2011	% Change from 2003 to 2011	2011 Data		
					Num	Den	Percent
1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2)	81.7%	83.3%	83.3%	2.0%	186	223	83.4%
1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services.	75.6%	77.8%	76.2%	0.9%	172	230	74.8%
1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services.	61.9%	59.2%	61.0%	-1.5%	540	866	62.4%
2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy.	50.8%	51.3%	44.6%	-12.3%	166	420	39.5%
2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum.	81.9%	80.5%	82.4%	0.6%	180	211	85.3%
3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1)	94.7%	97.6%	91.7%	-3.2%			
3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1)	87.4%	90.7%	94.1%	7.7%			
4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were:							
a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1)	66.9%	66.8%	59.6%	-10.9%	202	395	51.1%
b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile)	12.0%	10.6%	15.7%	31.0%	80	395	20.3%
c. Obese (BMI at or greater than the 95th percentile).	13.3%	13.5%	17.3%	29.9%	95	395	24.1%
d. Underweight (BMI less than the 5th percentile).	7.8%	9.1%	7.4%	-5.4%	18	395	4.6%
4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3)							
a. Recommended prenatal weight gain	27.7%	28.5%	25.1%	-9.3%	52	220	23.6%
b. Excessive prenatal weight gain	48.8%	48.1%	49.3%	0.9%	110	220	50.0%
c. Inadequate prenatal weight gain	23.5%	23.3%	25.6%	9.1%	58	220	26.4%
5A. Percentage of women who participated in WIC who initiated breastfeeding.	31.2%	33.6%	39.4%	26.3%	98	249	39.4%
5B. Percentage of infants who participated in WIC who were:							
a. Breastfeeding at 6 weeks of age (1)	16.4%	18.3%	15.3%	-6.5%	37	256	14.5%
b. Breastfeeding at 6 months of age	6.1%	5.6%	5.9%	-2.9%	12	256	4.7%

(1) Item appears also on other WCH Section Agreement Addenda.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI >=30.0): Adapted from the National Academy of Sciences. Weight Gain During Pregnancy: Reexamining the Guidelines. Washington, DC : National Academy Press, 2009.

SFY 2013-2014 AGREEMENT ADDENDA DATA

ATTACHMENT A: WIC QUALITY ASSURANCE DELIVERABLES

Appalachian District

Process Outcome Objectives	2003 to 2005	2006 to 2008	2009 to 2011	% Change from 2003 to 2011	2011 Data		
					Num	Den	Percent
1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2)	79.5%	64.5%	68.2%	-14.2%	243	370	65.7%
1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services.	68.1%	70.8%	67.4%	-1.0%	297	455	65.3%
1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services.	68.7%	62.4%	63.4%	-7.7%	1,241	1,947	63.7%
2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy.	51.9%	47.8%	40.7%	-21.5%	344	891	38.6%
2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum.	66.2%	73.3%	73.6%	11.1%	307	420	73.1%
3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1)	86.7%	92.1%					
3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1)	83.5%	85.7%					
4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were:							
a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1)	67.1%	66.4%	64.2%	-4.3%	425	679	62.6%
b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile)	15.1%	14.8%	17.4%	15.5%	120	679	17.7%
c. Obese (BMI at or greater than the 95th percentile).	13.4%	15.4%	14.5%	8.5%	115	679	16.9%
d. Underweight (BMI less than the 5th percentile).	4.5%	3.4%	3.9%	-13.2%	19	679	2.8%
4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3)							
a. Recommended prenatal weight gain	31.5%	31.9%	31.3%	-0.8%	117	357	32.8%
b. Excessive prenatal weight gain	47.9%	48.6%	48.3%	1.0%	174	357	48.7%
c. Inadequate prenatal weight gain	20.6%	19.5%	20.4%	-1.2%	66	357	18.5%
5A. Percentage of women who participated in WIC who initiated breastfeeding.	52.0%	58.7%	60.7%	16.7%	329	515	63.9%
5B. Percentage of infants who participated in WIC who were:							
a. Breastfeeding at 6 weeks of age (1)	33.7%	36.3%	36.2%	7.5%	204	524	38.9%
b. Breastfeeding at 6 months of age	18.4%	19.7%	21.4%	16.0%	123	524	23.5%

(1) Item appears also on other WCH Section Agreement Addenda.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI >=30.0): Adapted from the National Academy of Sciences. *Weight Gain During Pregnancy: Reexamining the Guidelines*. Washington, DC : National Academy Press, 2009.

SFY 2013-2014 AGREEMENT ADDENDA DATA

ATTACHMENT A: WIC QUALITY ASSURANCE DELIVERABLES

Beaufort County

Process Outcome Objectives	2003 to 2005	2006 to 2008	2009 to 2011	% Change from 2003 to 2011	2011 Data		
					Num	Den	Percent
1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2)	83.3%	80.0%	80.7%	-3.0%	235	285	82.5%
1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services.	75.5%	75.2%	74.0%	-1.9%	254	362	70.2%
1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services.	58.8%	58.5%	61.2%	4.1%	1,000	1,571	63.7%
2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy.	38.7%	37.5%	35.7%	-7.7%	333	964	34.5%
2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum.	74.5%	70.4%	79.0%	5.9%	312	391	79.8%
3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1)							
3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1)							
4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were:							
a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1)	64.6%	65.5%	64.0%	-1.0%	521	817	63.8%
b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile)	15.8%	15.6%	16.2%	2.6%	138	817	16.9%
c. Obese (BMI at or greater than the 95th percentile).	16.9%	16.3%	16.1%	-4.5%	129	817	15.8%
d. Underweight (BMI less than the 5th percentile).	2.7%	2.6%	3.7%	35.8%	29	817	3.5%
4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3)							
a. Recommended prenatal weight gain	33.0%	33.5%	31.6%	-4.1%	83	321	25.9%
b. Excessive prenatal weight gain	44.8%	45.3%	44.4%	-0.8%	145	321	45.2%
c. Inadequate prenatal weight gain	22.2%	21.2%	23.9%	7.8%	93	321	29.0%
5A. Percentage of women who participated in WIC who initiated breastfeeding.	43.7%	42.8%	48.0%	9.9%	227	462	49.1%
5B. Percentage of infants who participated in WIC who were:							
a. Breastfeeding at 6 weeks of age (1)	23.0%	27.4%	29.4%	27.7%	162	438	37.0%
b. Breastfeeding at 6 months of age	11.7%	13.5%	13.5%	14.9%	79	438	18.0%

(1) Item appears also on other WCH Section Agreement Addenda.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI < 18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI ≥ 30.0): Adapted from the National Academy of Sciences. Weight Gain During Pregnancy: Reexamining the Guidelines. Washington, DC : National Academy Press, 2009.

SFY 2013-2014 AGREEMENT ADDENDA DATA

ATTACHMENT A: WIC QUALITY ASSURANCE DELIVERABLES

Bladen County

Process Outcome Objectives	2003 to 2005	2006 to 2008	2009 to 2011	% Change from 2003 to 2011	2011 Data		
					Num	Den	Percent
1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2)	82.6%	82.5%	83.5%	1.2%	154	198	77.8%
1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services.	81.5%	83.2%	79.6%	-2.3%	204	263	77.6%
1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services.	62.5%	63.2%	65.9%	5.4%	813	1,214	67.0%
2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy.	42.5%	46.9%	38.8%	-8.7%	275	787	34.9%
2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum.	74.8%	72.8%	73.0%	-2.4%	232	321	72.3%
3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1)	90.7%	94.1%	96.2%	6.0%			
3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1)	92.2%	90.2%	93.8%	1.8%			
4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were:							
a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1)	66.6%	63.4%	62.9%	-5.5%	341	535	63.7%
b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile)	14.4%	15.9%	16.8%	16.2%	91	535	17.0%
c. Obese (BMI at or greater than the 95th percentile).	12.8%	16.0%	16.9%	31.6%	85	535	15.9%
d. Underweight (BMI less than the 5th percentile).	6.1%	4.7%	3.4%	-44.5%	18	535	3.4%
4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3)							
a. Recommended prenatal weight gain	29.1%	30.1%	25.8%	-11.3%	65	250	26.0%
b. Excessive prenatal weight gain	44.5%	42.9%	47.9%	7.6%	109	250	43.6%
c. Inadequate prenatal weight gain	26.4%	27.1%	26.3%	-0.4%	76	250	30.4%
5A. Percentage of women who participated in WIC who initiated breastfeeding.	45.8%	58.7%	52.6%	15.0%	209	383	54.6%
5B. Percentage of infants who participated in WIC who were:							
a. Breastfeeding at 6 weeks of age (1)	28.1%	46.9%	41.7%	48.4%	147	342	43.0%
b. Breastfeeding at 6 months of age	16.3%	33.7%	30.6%	87.5%	112	342	32.7%

(1) Item appears also on other WCH Section Agreement Addenda.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI >=30.0): Adapted from the National Academy of Sciences. Weight Gain During Pregnancy: Reexamining the Guidelines. Washington, DC : National Academy Press, 2009.

SFY 2013-2014 AGREEMENT ADDENDA DATA

ATTACHMENT A: WIC QUALITY ASSURANCE DELIVERABLES

Brunswick County

Process Outcome Objectives	2003 to 2005	2006 to 2008	2009 to 2011	% Change from 2003 to 2011	2011 Data		
					Num	Den	Percent
1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2)	74.0%	75.4%	73.6%	-0.6%	379	542	69.9%
1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services.	71.5%	72.6%	72.6%	1.5%	470	633	74.2%
1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services.	52.3%	48.8%	55.8%	6.7%	1,706	2,824	60.4%
2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy.	34.5%	32.4%	29.2%	-15.3%	376	1,339	28.1%
2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum.	69.9%	73.8%	74.7%	6.8%	425	569	74.7%
3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1)	92.4%	90.3%	85.7%	-7.2%			
3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1)	80.4%	77.4%	84.2%	4.8%			
4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were:							
a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1)	66.9%	64.6%	62.4%	-6.8%	664	1,085	61.2%
b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile)	15.6%	17.5%	17.0%	8.8%	192	1,085	17.7%
c. Obese (BMI at or greater than the 95th percentile).	14.9%	14.8%	17.4%	17.2%	190	1,085	17.5%
d. Underweight (BMI less than the 5th percentile).	2.6%	3.0%	3.2%	22.5%	39	1,085	3.6%
4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3)							
a. Recommended prenatal weight gain	31.8%	31.2%	29.4%	-7.7%	140	529	26.5%
b. Excessive prenatal weight gain	50.2%	51.2%	51.0%	1.6%	297	529	56.1%
c. Inadequate prenatal weight gain	18.1%	17.5%	19.7%	9.0%	92	529	17.4%
5A. Percentage of women who participated in WIC who initiated breastfeeding.	40.6%	37.4%	36.7%	-9.6%	313	738	42.4%
5B. Percentage of infants who participated in WIC who were:							
a. Breastfeeding at 6 weeks of age (1)	25.6%	25.0%	24.5%	-4.5%	189	758	24.9%
b. Breastfeeding at 6 months of age	14.6%	13.3%	13.2%	-9.8%	103	758	13.6%

(1) Item appears also on other WCH Section Agreement Addenda.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI >=30.0): Adapted from the National Academy of Sciences. Weight Gain During Pregnancy: Reexamining the Guidelines. Washington, DC : National Academy Press, 2009.

SFY 2013-2014 AGREEMENT ADDENDA DATA

ATTACHMENT A: WIC QUALITY ASSURANCE DELIVERABLES

Buncombe County

Process Outcome Objectives	2003 to 2005	2006 to 2008	2009 to 2011	% Change from 2003 to 2011	2011 Data		
					Num	Den	Percent
1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2)	72.2%	70.4%	70.9%	-1.8%	959	1,385	69.2%
1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services.	65.9%	66.5%	62.1%	-5.8%	1,030	1,683	61.2%
1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services.	54.3%	51.4%	52.1%	-4.1%	3,800	6,968	54.5%
2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy.	42.8%	35.9%	34.8%	-18.6%	1,012	2,950	34.3%
2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum.	76.4%	75.8%	72.9%	-4.6%	1,082	1,426	75.9%
3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1)	94.0%	92.9%	88.8%	-5.6%			
3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1)	87.9%	87.7%	90.2%	2.6%			
4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were:							
a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1)	63.6%	64.5%	65.4%	2.9%	1,258	1,906	66.0%
b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile)	16.5%	16.6%	16.1%	-2.8%	303	1,906	15.9%
c. Obese (BMI at or greater than the 95th percentile).	16.0%	16.5%	15.3%	-4.5%	287	1,906	15.1%
d. Underweight (BMI less than the 5th percentile).	3.8%	2.5%	3.2%	-17.6%	58	1,906	3.0%
4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3)							
a. Recommended prenatal weight gain	34.3%	33.5%	30.6%	-10.7%	351	1,310	26.8%
b. Excessive prenatal weight gain	45.2%	47.4%	48.5%	7.4%	631	1,310	48.2%
c. Inadequate prenatal weight gain	20.5%	19.1%	20.8%	1.4%	328	1,310	25.0%
5A. Percentage of women who participated in WIC who initiated breastfeeding.	68.9%	69.1%	72.5%	5.2%	1,254	1,728	72.6%
5B. Percentage of infants who participated in WIC who were:							
a. Breastfeeding at 6 weeks of age (1)	34.9%	35.9%	39.0%	11.9%	650	1,659	39.2%
b. Breastfeeding at 6 months of age	19.2%	20.6%	22.4%	16.4%	389	1,659	23.4%

(1) Item appears also on other WCH Section Agreement Addenda.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI < 18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI ≥ 30.0): Adapted from the National Academy of Sciences. *Weight Gain During Pregnancy: Reexamining the Guidelines*. Washington, DC : National Academy Press, 2009.

SFY 2013-2014 AGREEMENT ADDENDA DATA

ATTACHMENT A: WIC QUALITY ASSURANCE DELIVERABLES

Burke County

Process Outcome Objectives	2003 to 2005	2006 to 2008	2009 to 2011	% Change from 2003 to 2011	2011 Data		
					Num	Den	Percent
1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2)	65.1%	69.9%	71.3%	9.5%	387	569	68.0%
1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services.	74.4%	75.6%	77.7%	4.4%	520	664	78.3%
1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services.	55.9%	55.8%	59.6%	6.7%	1,754	2,805	62.5%
2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy.	28.1%	20.1%	25.5%	-9.3%	362	1,204	30.1%
2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum.	83.0%	84.3%	83.0%	-0.1%	504	596	84.6%
3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1)	100.0%	83.3%	66.7%	-33.3%			
3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1)	71.8%	71.5%	79.2%	10.3%			
4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were:							
a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1)	66.6%	63.7%	63.2%	-5.2%	689	1,078	63.9%
b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile)	16.0%	17.8%	17.2%	7.8%	189	1,078	17.5%
c. Obese (BMI at or greater than the 95th percentile).	13.3%	16.2%	16.6%	25.1%	162	1,078	15.0%
d. Underweight (BMI less than the 5th percentile).	4.1%	2.4%	3.0%	-27.0%	38	1,078	3.5%
4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3)							
a. Recommended prenatal weight gain	34.0%	33.2%	29.8%	-12.3%	145	520	27.9%
b. Excessive prenatal weight gain	42.5%	42.2%	44.4%	4.3%	242	520	46.5%
c. Inadequate prenatal weight gain	23.5%	24.6%	25.8%	10.0%	133	520	25.6%
5A. Percentage of women who participated in WIC who initiated breastfeeding.	56.5%	60.9%	64.6%	14.3%	511	730	70.0%
5B. Percentage of infants who participated in WIC who were:							
a. Breastfeeding at 6 weeks of age (1)	22.8%	23.3%	26.6%	16.9%	197	695	28.3%
b. Breastfeeding at 6 months of age	9.4%	10.9%	12.2%	29.5%	94	695	13.5%

(1) Item appears also on other WCH Section Agreement Addenda.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI >=30.0): Adapted from the National Academy of Sciences. Weight Gain During Pregnancy: Reexamining the Guidelines. Washington, DC : National Academy Press, 2009.

SFY 2013-2014 AGREEMENT ADDENDA DATA

ATTACHMENT A: WIC QUALITY ASSURANCE DELIVERABLES

Cabarrus County

Process Outcome Objectives	2003 to 2005	2006 to 2008	2009 to 2011	% Change from 2003 to 2011	2011 Data		
					Num	Den	Percent
1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2)	67.3%	71.8%	66.5%	-1.1%	673	1,001	67.2%
1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services.	71.4%	73.1%	69.6%	-2.5%	929	1,314	70.7%
1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services.	54.9%	55.2%	53.4%	-2.7%	2,949	5,417	54.4%
2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy.	29.2%	22.5%	24.9%	-14.7%	679	2,346	28.9%
2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum.	77.9%	82.0%	78.6%	0.9%	884	1,117	79.1%
3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1)	79.9%	86.1%	94.2%	17.9%			
3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1)	66.6%	72.6%	75.2%	12.9%			
4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were:							
a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1)	64.8%	62.9%	62.2%	-4.0%	1,211	1,899	63.8%
b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile)	16.6%	17.0%	18.0%	8.4%	325	1,899	17.1%
c. Obese (BMI at or greater than the 95th percentile).	14.0%	17.6%	16.8%	20.2%	300	1,899	15.8%
d. Underweight (BMI less than the 5th percentile).	4.6%	2.5%	3.0%	-35.4%	63	1,899	3.3%
4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3)							
a. Recommended prenatal weight gain	30.2%	30.1%	30.0%	-0.7%	317	1,132	28.0%
b. Excessive prenatal weight gain	45.6%	46.8%	46.0%	1.0%	545	1,132	48.1%
c. Inadequate prenatal weight gain	24.3%	23.2%	24.0%	-1.1%	270	1,132	23.9%
5A. Percentage of women who participated in WIC who initiated breastfeeding.	55.8%	64.5%	62.7%	12.4%	917	1,436	63.9%
5B. Percentage of infants who participated in WIC who were:							
a. Breastfeeding at 6 weeks of age (1)	35.3%	39.0%	37.1%	5.2%	562	1,451	38.7%
b. Breastfeeding at 6 months of age	20.4%	21.9%	20.2%	-1.1%	286	1,451	19.7%

(1) Item appears also on other WCH Section Agreement Addenda.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI >=30.0): Adapted from the National Academy of Sciences. Weight Gain During Pregnancy: Reexamining the Guidelines. Washington, DC : National Academy Press, 2009.

SFY 2013-2014 AGREEMENT ADDENDA DATA

ATTACHMENT A: WIC QUALITY ASSURANCE DELIVERABLES

Caldwell County

Process Outcome Objectives	2003 to 2005	2006 to 2008	2009 to 2011	% Change from 2003 to 2011	2011 Data		
					Num	Den	Percent
1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2)	78.8%	72.1%	78.8%	-0.1%	409	535	76.4%
1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services.	77.4%	78.6%	79.1%	2.2%	499	624	80.0%
1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services.	59.2%	57.8%	59.6%	0.7%	1,496	2,492	60.0%
2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy.	34.7%	32.5%	29.6%	-14.6%	457	1,670	27.4%
2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum.	78.9%	77.6%	79.0%	0.2%	475	579	82.0%
3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1)	77.8%	75.0%					
3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1)	61.5%	65.3%	68.4%	11.3%			
4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were:							
a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1)	63.9%	58.8%	62.0%	-3.1%	528	848	62.3%
b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile)	17.7%	19.7%	17.6%	-0.5%	150	848	17.7%
c. Obese (BMI at or greater than the 95th percentile).	15.2%	19.1%	17.3%	14.2%	139	848	16.4%
d. Underweight (BMI less than the 5th percentile).	3.2%	2.4%	3.1%	-2.7%	31	848	3.7%
4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3)							
a. Recommended prenatal weight gain	27.5%	30.2%	30.2%	9.9%	179	554	32.3%
b. Excessive prenatal weight gain	53.3%	51.5%	49.3%	-7.5%	258	554	46.6%
c. Inadequate prenatal weight gain	19.2%	18.4%	20.5%	6.5%	117	554	21.1%
5A. Percentage of women who participated in WIC who initiated breastfeeding.	51.2%	47.0%	54.6%	6.6%	425	698	60.9%
5B. Percentage of infants who participated in WIC who were:							
a. Breastfeeding at 6 weeks of age (1)	24.7%	26.8%	27.7%	12.1%	181	662	27.3%
b. Breastfeeding at 6 months of age	11.3%	12.6%	11.1%	-1.7%	78	662	11.8%

(1) Item appears also on other WCH Section Agreement Addenda.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI >=30.0): Adapted from the National Academy of Sciences. Weight Gain During Pregnancy: Reexamining the Guidelines. Washington, DC : National Academy Press, 2009.

SFY 2013-2014 AGREEMENT ADDENDA DATA

ATTACHMENT A: WIC QUALITY ASSURANCE DELIVERABLES

Carteret County

Process Outcome Objectives	2003 to 2005	2006 to 2008	2009 to 2011	% Change from 2003 to 2011	2011 Data		
					Num	Den	Percent
1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2)	60.6%	69.1%	67.4%	11.1%	215	299	71.9%
1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services.	73.6%	75.5%	75.0%	2.0%	277	368	75.3%
1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services.	55.9%	53.6%	57.1%	2.1%	834	1,409	59.2%
2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy.	35.7%	28.5%	29.1%	-18.5%	229	757	30.3%
2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum.	80.5%	71.8%	69.5%	-13.7%	276	359	76.9%
3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1)	50.0%						
3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1)	58.6%	100.0%	33.3%	-43.1%			
4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were:							
a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1)	63.4%	65.1%	64.1%	1.1%	354	548	64.6%
b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile)	15.9%	16.4%	16.7%	5.2%	96	548	17.5%
c. Obese (BMI at or greater than the 95th percentile).	17.6%	15.9%	14.0%	-20.6%	60	548	10.9%
d. Underweight (BMI less than the 5th percentile).	3.1%	2.6%	5.2%	69.1%	38	548	6.9%
4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3)							
a. Recommended prenatal weight gain	32.9%	32.6%	31.1%	-5.5%	110	348	31.6%
b. Excessive prenatal weight gain	52.2%	52.2%	48.3%	-7.4%	162	348	46.6%
c. Inadequate prenatal weight gain	14.9%	15.3%	20.5%	37.9%	76	348	21.8%
5A. Percentage of women who participated in WIC who initiated breastfeeding.	52.0%	57.7%	63.3%	21.7%	310	479	64.7%
5B. Percentage of infants who participated in WIC who were:							
a. Breastfeeding at 6 weeks of age (1)	27.4%	30.1%	31.6%	15.5%	146	451	32.4%
b. Breastfeeding at 6 months of age	11.1%	13.1%	14.6%	31.7%	74	451	16.4%

(1) Item appears also on other WCH Section Agreement Addenda.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI >=30.0): Adapted from the National Academy of Sciences. Weight Gain During Pregnancy: Reexamining the Guidelines. Washington, DC : National Academy Press, 2009.

SFY 2013-2014 AGREEMENT ADDENDA DATA

ATTACHMENT A: WIC QUALITY ASSURANCE DELIVERABLES

Caswell County

Process Outcome Objectives	2003 to 2005	2006 to 2008	2009 to 2011	% Change from 2003 to 2011	2011 Data		
					Num	Den	Percent
1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2)	82.6%	80.7%	76.5%	-7.4%	97	151	64.2%
1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services.	74.3%	77.0%	73.4%	-1.2%	135	184	73.4%
1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services.	57.4%	60.0%	59.7%	3.9%	408	666	61.3%
2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy.	40.5%	41.6%	37.8%	-6.8%	110	318	34.6%
2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum.	77.7%	77.3%	79.6%	2.4%	106	137	77.4%
3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1)	95.8%	97.4%	100.0%	4.4%			
3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1)	86.4%	91.2%	92.8%	7.4%			
4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were:							
a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1)	69.7%	68.7%	69.4%	-0.3%	136	194	70.1%
b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile)	12.4%	15.2%	16.2%	30.4%	29	194	14.9%
c. Obese (BMI at or greater than the 95th percentile).	13.8%	12.5%	11.1%	-19.6%	24	194	12.4%
d. Underweight (BMI less than the 5th percentile).	4.1%	3.6%	3.2%	-20.8%	5	194	2.6%
4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3)							
a. Recommended prenatal weight gain	34.0%	36.5%	27.4%	-19.6%	22	121	18.2%
b. Excessive prenatal weight gain	41.3%	45.8%	45.0%	8.9%	51	121	42.1%
c. Inadequate prenatal weight gain	24.7%	17.7%	27.6%	12.1%	48	121	39.7%
5A. Percentage of women who participated in WIC who initiated breastfeeding.	40.6%	52.9%	54.3%	33.6%	97	177	54.8%
5B. Percentage of infants who participated in WIC who were:							
a. Breastfeeding at 6 weeks of age (1)	21.9%	23.5%	30.8%	40.6%	63	180	35.0%
b. Breastfeeding at 6 months of age	7.2%	10.0%	10.9%	50.6%	26	180	14.4%

(1) Item appears also on other WCH Section Agreement Addenda.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI >=30.0): Adapted from the National Academy of Sciences. Weight Gain During Pregnancy: Reexamining the Guidelines. Washington, DC : National Academy Press, 2009.

SFY 2013-2014 AGREEMENT ADDENDA DATA

ATTACHMENT A: WIC QUALITY ASSURANCE DELIVERABLES

Catawba County

Process Outcome Objectives	2003 to 2005	2006 to 2008	2009 to 2011	% Change from 2003 to 2011	2011 Data		
					Num	Den	Percent
1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2)	70.5%	75.4%	68.8%	-2.3%	655	1,017	64.4%
1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services.	76.6%	76.4%	75.7%	-1.1%	988	1,279	77.2%
1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services.	51.8%	51.7%	53.0%	2.3%	2,992	5,277	56.7%
2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy.	28.9%	23.6%	34.1%	18.1%	696	1,962	35.5%
2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum.	58.2%	56.5%	68.0%	16.8%	752	1,080	69.6%
3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1)	68.8%	68.8%	50.0%	-27.3%			
3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1)	50.5%	41.3%	42.6%	-15.7%			
4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were:							
a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1)	61.7%	61.3%	63.7%	3.3%	499	773	64.6%
b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile)	18.7%	19.2%	16.9%	-9.6%	122	773	15.8%
c. Obese (BMI at or greater than the 95th percentile).	16.9%	17.4%	16.0%	-5.2%	123	773	15.9%
d. Underweight (BMI less than the 5th percentile).	2.8%	2.2%	3.4%	23.4%	29	773	3.8%
4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3)							
a. Recommended prenatal weight gain	32.9%	31.4%	31.3%	-4.8%	313	997	31.4%
b. Excessive prenatal weight gain	43.1%	32.8%	41.9%	-2.9%	435	997	43.6%
c. Inadequate prenatal weight gain	24.0%	35.8%	26.8%	11.8%	249	997	25.0%
5A. Percentage of women who participated in WIC who initiated breastfeeding.	53.0%	60.2%	62.0%	17.0%	836	1,358	61.6%
5B. Percentage of infants who participated in WIC who were:							
a. Breastfeeding at 6 weeks of age (1)	28.5%	30.6%	32.1%	12.6%	433	1,308	33.1%
b. Breastfeeding at 6 months of age	13.2%	14.4%	15.8%	19.7%	234	1,308	17.9%

(1) Item appears also on other WCH Section Agreement Addenda.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI >=30.0): Adapted from the National Academy of Sciences. Weight Gain During Pregnancy: Reexamining the Guidelines. Washington, DC : National Academy Press, 2009.

SFY 2013-2014 AGREEMENT ADDENDA DATA

ATTACHMENT A: WIC QUALITY ASSURANCE DELIVERABLES

Chatham County

Process Outcome Objectives	2003 to 2005	2006 to 2008	2009 to 2011	% Change from 2003 to 2011	2011 Data		
					Num	Den	Percent
1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2)	64.4%	69.6%	69.9%	8.5%	127	188	67.6%
1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services.	48.9%	48.3%	57.2%	17.1%	262	385	68.1%
1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services.	45.4%	41.7%	48.0%	5.7%	828	1,567	52.8%
2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy.	28.3%	28.2%	27.3%	-3.6%	639	2,405	26.6%
2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum.	60.5%	61.0%	67.7%	11.9%	787	1,116	70.5%
3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1)	66.5%	68.4%	66.7%	0.2%			
3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1)	63.3%	71.3%	71.4%	12.8%			
4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were:							
a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1)	62.2%	61.1%	61.7%	-0.7%	690	1,153	59.8%
b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile)	17.0%	17.2%	16.9%	-0.6%	212	1,153	18.4%
c. Obese (BMI at or greater than the 95th percentile).	18.1%	19.3%	17.7%	-2.0%	206	1,153	17.9%
d. Underweight (BMI less than the 5th percentile).	2.8%	2.4%	3.7%	32.4%	45	1,153	3.9%
4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3)							
a. Recommended prenatal weight gain	30.1%	36.1%	31.7%	5.6%	88	292	30.1%
b. Excessive prenatal weight gain	48.4%	43.1%	45.0%	-7.2%	128	292	43.8%
c. Inadequate prenatal weight gain	21.5%	20.9%	23.3%	8.3%	76	292	26.0%
5A. Percentage of women who participated in WIC who initiated breastfeeding.	61.3%	68.4%	68.8%	12.3%	1,012	1,407	71.9%
5B. Percentage of infants who participated in WIC who were:							
a. Breastfeeding at 6 weeks of age (1)	49.8%	52.4%	52.8%	6.0%	782	1,435	54.5%
b. Breastfeeding at 6 months of age	29.7%	32.6%	32.8%	10.5%	484	1,435	33.7%

(1) Item appears also on other WCH Section Agreement Addenda.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI >=30.0): Adapted from the National Academy of Sciences. Weight Gain During Pregnancy: Reexamining the Guidelines. Washington, DC : National Academy Press, 2009.

SFY 2013-2014 AGREEMENT ADDENDA DATA

ATTACHMENT A: WIC QUALITY ASSURANCE DELIVERABLES

Cherokee County

Process Outcome Objectives	2003 to 2005	2006 to 2008	2009 to 2011	% Change from 2003 to 2011	2011 Data		
					Num	Den	Percent
1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2)	77.4%	78.0%	79.3%	2.5%	123	169	72.8%
1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services.	76.1%	74.2%	72.9%	-4.2%	145	200	72.5%
1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services.	66.0%	62.8%	65.3%	-1.0%	545	818	66.6%
2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy.	43.3%	42.7%	47.2%	9.0%	158	345	45.8%
2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum.	82.8%	79.2%	81.4%	-1.7%	125	155	80.6%
3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1)	73.3%	70.0%	100.0%	36.4%			
3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1)	83.2%	75.7%	81.4%	-2.2%			
4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were:							
a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1)	65.6%	71.3%	64.1%	-2.2%	167	266	62.8%
b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile)	14.8%	14.4%	15.4%	3.8%	39	266	14.7%
c. Obese (BMI at or greater than the 95th percentile).	15.7%	11.2%	16.5%	5.3%	47	266	17.7%
d. Underweight (BMI less than the 5th percentile).	3.9%	3.0%	4.0%	2.2%	13	266	4.9%
4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3)							
a. Recommended prenatal weight gain	29.9%	27.4%	31.4%	4.8%	47	160	29.4%
b. Excessive prenatal weight gain	48.5%	50.4%	44.1%	-9.1%	77	160	48.1%
c. Inadequate prenatal weight gain	21.6%	22.2%	24.6%	13.9%	36	160	22.5%
5A. Percentage of women who participated in WIC who initiated breastfeeding.	44.4%	50.1%	54.6%	22.9%	99	187	52.9%
5B. Percentage of infants who participated in WIC who were:							
a. Breastfeeding at 6 weeks of age (1)	28.5%	32.4%	32.0%	12.3%	57	201	28.4%
b. Breastfeeding at 6 months of age	13.4%	15.3%	15.2%	13.3%	27	201	13.4%

(1) Item appears also on other WCH Section Agreement Addenda.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI >=30.0): Adapted from the National Academy of Sciences. Weight Gain During Pregnancy: Reexamining the Guidelines. Washington, DC : National Academy Press, 2009.

SFY 2013-2014 AGREEMENT ADDENDA DATA

ATTACHMENT A: WIC QUALITY ASSURANCE DELIVERABLES

Clay County

Process Outcome Objectives	2003 to 2005	2006 to 2008	2009 to 2011	% Change from 2003 to 2011	2011 Data		
					Num	Den	Percent
1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2)	80.0%	83.8%	77.3%	-3.3%	37	49	75.5%
1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services.	77.0%	73.3%	76.9%	0.0%	47	64	73.4%
1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services.	69.5%	67.2%	65.9%	-5.3%	224	341	65.7%
2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy.	47.2%	54.1%	46.4%	-1.8%	80	183	43.7%
2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum.	77.0%	69.0%	72.3%	-6.1%	49	67	73.1%
3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1)	73.3%	86.2%	100.0%	36.4%			
3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1)	82.5%	81.6%	84.2%	2.0%			
4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were:							
a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1)	60.2%	65.6%	66.8%	11.1%	97	142	68.3%
b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile)	15.5%	13.9%	15.1%	-2.5%	20	142	14.1%
c. Obese (BMI at or greater than the 95th percentile).	19.8%	16.6%	15.9%	-19.5%	23	142	16.2%
d. Underweight (BMI less than the 5th percentile).	4.5%	3.9%	2.1%	-53.8%	2	142	1.4%
4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3)							
a. Recommended prenatal weight gain	26.3%	24.3%	30.6%	16.2%	19	55	34.5%
b. Excessive prenatal weight gain	54.1%	54.4%	49.4%	-8.7%	29	55	52.7%
c. Inadequate prenatal weight gain	19.5%	21.4%	20.0%	2.3%	7	55	12.7%
5A. Percentage of women who participated in WIC who initiated breastfeeding.	43.8%	57.1%	61.3%	40.0%	49	81	60.5%
5B. Percentage of infants who participated in WIC who were:							
a. Breastfeeding at 6 weeks of age (1)	28.3%	37.5%	40.1%	41.8%	38	82	46.3%
b. Breastfeeding at 6 months of age	14.3%	22.7%	22.1%	54.3%	20	82	24.4%

(1) Item appears also on other WCH Section Agreement Addenda.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI >=30.0): Adapted from the National Academy of Sciences. Weight Gain During Pregnancy: Reexamining the Guidelines. Washington, DC : National Academy Press, 2009.

SFY 2013-2014 AGREEMENT ADDENDA DATA

ATTACHMENT A: WIC QUALITY ASSURANCE DELIVERABLES

Cleveland County

Process Outcome Objectives	2003 to 2005	2006 to 2008	2009 to 2011	% Change from 2003 to 2011	2011 Data		
					Num	Den	Percent
1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2)	81.1%	83.9%	84.2%	3.8%	616	743	82.9%
1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services.	77.7%	76.8%	75.2%	-3.2%	604	801	75.4%
1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services.	50.5%	48.5%	52.3%	3.5%	1,922	3,533	54.4%
2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy.	34.6%	34.8%	35.0%	1.0%	608	1,822	33.4%
2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum.	16.3%	19.6%	53.4%	228.2%	456	806	56.6%
3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1)	75.1%	78.7%	85.4%	13.8%			
3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1)	32.9%	40.0%	51.2%	55.3%			
4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were:							
a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1)	69.3%	69.8%	64.3%	-7.2%	653	1,008	64.8%
b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile)	12.0%	11.8%	15.9%	32.2%	161	1,008	16.0%
c. Obese (BMI at or greater than the 95th percentile).	11.2%	11.2%	15.4%	37.4%	155	1,008	15.4%
d. Underweight (BMI less than the 5th percentile).	7.4%	7.3%	4.4%	-41.2%	39	1,008	3.9%
4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3)							
a. Recommended prenatal weight gain	28.7%	28.7%	25.4%	-11.5%	173	752	23.0%
b. Excessive prenatal weight gain	48.8%	42.9%	37.6%	-22.9%	256	752	34.0%
c. Inadequate prenatal weight gain	22.5%	28.4%	37.0%	64.3%	323	752	43.0%
5A. Percentage of women who participated in WIC who initiated breastfeeding.	19.8%	22.6%	40.9%	106.3%	434	920	47.2%
5B. Percentage of infants who participated in WIC who were:							
a. Breastfeeding at 6 weeks of age (1)	12.5%	11.2%	22.8%	82.8%	353	896	39.4%
b. Breastfeeding at 6 months of age	6.6%	6.9%	16.9%	157.4%	290	896	32.4%

(1) Item appears also on other WCH Section Agreement Addenda.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI >=30.0): Adapted from the National Academy of Sciences. Weight Gain During Pregnancy: Reexamining the Guidelines. Washington, DC : National Academy Press, 2009.

SFY 2013-2014 AGREEMENT ADDENDA DATA

ATTACHMENT A: WIC QUALITY ASSURANCE DELIVERABLES

Columbus County

Process Outcome Objectives	2003 to 2005	2006 to 2008	2009 to 2011	% Change from 2003 to 2011	2011 Data		
					Num	Den	Percent
1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2)	74.1%	69.6%	75.9%	2.3%	351	475	73.9%
1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services.	79.6%	79.2%	78.9%	-0.9%	450	579	77.7%
1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services.	61.3%	59.0%	61.3%	0.0%	1,542	2,412	63.9%
2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy.	30.1%	31.8%	30.3%	0.6%	280	951	29.4%
2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum.	80.6%	83.1%	85.7%	6.3%	386	458	84.3%
3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1)	93.3%	93.2%	100.0%	7.1%			
3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1)	91.9%	92.4%	94.4%	2.8%			
4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were:							
a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1)	66.7%	64.9%	59.6%	-10.8%	547	964	56.7%
b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile)	15.4%	15.3%	17.3%	11.7%	166	964	17.2%
c. Obese (BMI at or greater than the 95th percentile).	13.1%	15.1%	19.2%	46.3%	223	964	23.1%
d. Underweight (BMI less than the 5th percentile).	4.7%	4.7%	4.0%	-15.0%	28	964	2.9%
4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3)							
a. Recommended prenatal weight gain	28.4%	26.0%	28.4%	-0.2%	133	470	28.3%
b. Excessive prenatal weight gain	44.7%	45.5%	45.7%	2.3%	200	470	42.6%
c. Inadequate prenatal weight gain	26.8%	28.5%	25.9%	-3.6%	137	470	29.1%
5A. Percentage of women who participated in WIC who initiated breastfeeding.	42.4%	36.8%	33.7%	-20.4%	226	595	38.0%
5B. Percentage of infants who participated in WIC who were:							
a. Breastfeeding at 6 weeks of age (1)	17.9%	22.6%	22.9%	27.8%	138	637	21.7%
b. Breastfeeding at 6 months of age	8.3%	13.3%	17.2%	107.0%	115	637	18.1%

(1) Item appears also on other WCH Section Agreement Addenda.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI >=30.0): Adapted from the National Academy of Sciences. *Weight Gain During Pregnancy: Reexamining the Guidelines*. Washington, DC : National Academy Press, 2009.

SFY 2013-2014 AGREEMENT ADDENDA DATA

ATTACHMENT A: WIC QUALITY ASSURANCE DELIVERABLES

Craven County

Process Outcome Objectives	2003 to 2005	2006 to 2008	2009 to 2011	% Change from 2003 to 2011	2011 Data		
					Num	Den	Percent
1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2)	78.8%	79.6%	82.8%	5.2%	484	597	81.1%
1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services.	77.8%	76.2%	76.9%	-1.1%	520	688	75.6%
1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services.	55.3%	53.5%	59.7%	7.9%	1,813	2,906	62.4%
2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy.	30.2%	31.3%	31.7%	4.9%	796	2,513	31.7%
2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum.	36.0%	31.9%	41.3%	14.7%	639	1,052	60.7%
3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1)	85.0%	82.7%	82.5%	-2.9%			
3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1)	87.9%	82.3%	82.5%	-6.1%			
4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were:							
a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1)	66.4%	67.2%	66.4%	0.0%	973	1,452	67.0%
b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile)	15.8%	15.5%	15.2%	-3.7%	223	1,452	15.4%
c. Obese (BMI at or greater than the 95th percentile).	13.4%	13.5%	13.1%	-2.2%	181	1,452	12.5%
d. Underweight (BMI less than the 5th percentile).	4.5%	3.8%	5.3%	20.2%	75	1,452	5.2%
4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3)							
a. Recommended prenatal weight gain	30.0%	34.2%	29.6%	-1.4%	255	953	26.8%
b. Excessive prenatal weight gain	49.2%	46.1%	43.6%	-11.4%	388	953	40.7%
c. Inadequate prenatal weight gain	20.8%	19.6%	26.8%	28.8%	310	953	32.5%
5A. Percentage of women who participated in WIC who initiated breastfeeding.	49.7%	46.2%	34.8%	-30.0%	533	1,277	41.7%
5B. Percentage of infants who participated in WIC who were:							
a. Breastfeeding at 6 weeks of age (1)	25.9%	24.7%	21.6%	-16.5%	259	1,306	19.8%
b. Breastfeeding at 6 months of age	13.2%	13.0%	11.9%	-9.5%	164	1,306	12.6%

(1) Item appears also on other WCH Section Agreement Addenda.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI >=30.0): Adapted from the National Academy of Sciences. *Weight Gain During Pregnancy: Reexamining the Guidelines*. Washington, DC : National Academy Press, 2009.

SFY 2013-2014 AGREEMENT ADDENDA DATA

ATTACHMENT A: WIC QUALITY ASSURANCE DELIVERABLES

Cumberland County

Process Outcome Objectives	2003 to 2005	2006 to 2008	2009 to 2011	% Change from 2003 to 2011	2011 Data		
					Num	Den	Percent
1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2)	74.4%	74.8%	73.6%	-1.1%	1,663	2,311	72.0%
1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services.	78.6%	77.2%	73.8%	-6.2%	1,936	2,661	72.8%
1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services.	56.7%	53.9%	55.1%	-2.8%	6,251	10,934	57.2%
2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy.	31.9%	30.1%	30.2%	-5.4%	2,413	7,813	30.9%
2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum.	75.4%	76.1%	75.3%	-0.1%	2,773	3,730	74.3%
3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1)	72.4%	61.8%	69.1%	-4.6%			
3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1)	75.8%	68.6%	69.0%	-8.9%			
4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were:							
a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1)	72.7%	74.6%	70.8%	-2.6%	4,365	6,259	69.7%
b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile)	12.4%	10.0%	11.1%	-10.7%	751	6,259	12.0%
c. Obese (BMI at or greater than the 95th percentile).	8.8%	7.4%	8.4%	-4.5%	574	6,259	9.2%
d. Underweight (BMI less than the 5th percentile).	6.1%	8.0%	9.7%	60.0%	569	6,259	9.1%
4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3)							
a. Recommended prenatal weight gain	32.3%	28.9%	28.3%	-12.4%	964	3,644	26.5%
b. Excessive prenatal weight gain	50.6%	52.9%	51.5%	1.9%	1,902	3,644	52.2%
c. Inadequate prenatal weight gain	17.1%	18.2%	20.2%	17.7%	778	3,644	21.4%
5A. Percentage of women who participated in WIC who initiated breastfeeding.	53.5%	56.5%	64.4%	20.4%	3,293	4,864	67.7%
5B. Percentage of infants who participated in WIC who were:							
a. Breastfeeding at 6 weeks of age (1)	22.9%	26.5%	28.4%	24.1%	1,532	5,231	29.3%
b. Breastfeeding at 6 months of age	10.1%	11.3%	12.8%	27.0%	695	5,231	13.3%

(1) Item appears also on other WCH Section Agreement Addenda.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI >=30.0): Adapted from the National Academy of Sciences. Weight Gain During Pregnancy: Reexamining the Guidelines. Washington, DC : National Academy Press, 2009.

SFY 2013-2014 AGREEMENT ADDENDA DATA

ATTACHMENT A: WIC QUALITY ASSURANCE DELIVERABLES

Dare County

Process Outcome Objectives	2003 to 2005	2006 to 2008	2009 to 2011	% Change from 2003 to 2011	2011 Data		
					Num	Den	Percent
1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2)	61.2%	64.1%	66.0%	7.7%	111	167	66.5%
1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services.	59.0%	65.0%	65.8%	11.6%	161	241	66.8%
1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services.	56.1%	51.8%	61.0%	8.7%	590	929	63.5%
2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy.	26.3%	31.7%	29.6%	12.5%	159	575	27.7%
2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum.	65.0%	71.9%	71.7%	10.4%	181	242	74.8%
3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1)	83.6%	81.0%	71.4%	-14.6%			
3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1)	78.3%	78.6%	91.2%	16.5%			
4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were:							
a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1)	65.8%	63.5%	62.7%	-4.7%	218	363	60.1%
b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile)	18.0%	16.1%	18.7%	3.9%	73	363	20.1%
c. Obese (BMI at or greater than the 95th percentile).	14.6%	17.9%	15.9%	8.6%	62	363	17.1%
d. Underweight (BMI less than the 5th percentile).	1.5%	2.5%	2.6%	71.0%	10	363	2.8%
4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3)							
a. Recommended prenatal weight gain	37.5%	36.1%	35.9%	-4.3%	49	175	28.0%
b. Excessive prenatal weight gain	43.4%	39.5%	41.6%	-4.2%	86	175	49.1%
c. Inadequate prenatal weight gain	19.0%	24.4%	22.4%	18.0%	40	175	22.9%
5A. Percentage of women who participated in WIC who initiated breastfeeding.	59.1%	64.3%	71.9%	21.6%	202	296	68.2%
5B. Percentage of infants who participated in WIC who were:							
a. Breastfeeding at 6 weeks of age (1)	50.4%	60.4%	57.8%	14.6%	141	237	59.5%
b. Breastfeeding at 6 months of age	25.5%	32.4%	36.6%	43.6%	99	237	41.8%

(1) Item appears also on other WCH Section Agreement Addenda.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI >=30.0): Adapted from the National Academy of Sciences. Weight Gain During Pregnancy: Reexamining the Guidelines. Washington, DC : National Academy Press, 2009.

SFY 2013-2014 AGREEMENT ADDENDA DATA

ATTACHMENT A: WIC QUALITY ASSURANCE DELIVERABLES

Davidson County

Process Outcome Objectives	2003 to 2005	2006 to 2008	2009 to 2011	% Change from 2003 to 2011	2011 Data		
					Num	Den	Percent
1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2)	68.5%	70.7%	71.5%	4.4%	712	995	71.6%
1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services.	69.9%	73.3%	73.4%	5.1%	893	1,209	73.9%
1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services.	52.3%	50.2%	55.0%	5.1%	2,847	4,972	57.3%
2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy.	33.0%	30.4%	31.6%	-4.3%	631	1,938	32.6%
2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum.	74.2%	76.5%	77.0%	3.8%	784	1,051	74.6%
3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1)	44.4%						
3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1)	77.1%	70.7%	66.7%	-13.5%			
4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were:							
a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1)	70.3%	67.9%	65.5%	-6.9%	1,202	1,840	65.3%
b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile)	14.2%	14.7%	16.1%	13.3%	283	1,840	15.4%
c. Obese (BMI at or greater than the 95th percentile).	11.8%	14.0%	15.3%	29.7%	302	1,840	16.4%
d. Underweight (BMI less than the 5th percentile).	3.7%	3.4%	3.2%	-14.0%	53	1,840	2.9%
4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3)							
a. Recommended prenatal weight gain	30.8%	33.6%	29.7%	-3.5%	299	1,042	28.7%
b. Excessive prenatal weight gain	47.8%	42.5%	48.3%	0.8%	517	1,042	49.6%
c. Inadequate prenatal weight gain	21.4%	23.9%	22.1%	3.2%	226	1,042	21.7%
5A. Percentage of women who participated in WIC who initiated breastfeeding.	52.4%	58.3%	71.7%	36.9%	952	1,340	71.0%
5B. Percentage of infants who participated in WIC who were:							
a. Breastfeeding at 6 weeks of age (1)	27.7%	27.9%	29.5%	6.6%	370	1,275	29.0%
b. Breastfeeding at 6 months of age	14.2%	14.0%	15.8%	11.1%	208	1,275	16.3%

(1) Item appears also on other WCH Section Agreement Addenda.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI >=30.0): Adapted from the National Academy of Sciences. Weight Gain During Pregnancy: Reexamining the Guidelines. Washington, DC : National Academy Press, 2009.

SFY 2013-2014 AGREEMENT ADDENDA DATA

ATTACHMENT A: WIC QUALITY ASSURANCE DELIVERABLES

Davie County

Process Outcome Objectives	2003 to 2005	2006 to 2008	2009 to 2011	% Change from 2003 to 2011	2011 Data		
					Num	Den	Percent
1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2)	72.2%	72.2%	70.3%	-2.6%	122	181	67.4%
1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services.	71.3%	70.9%	71.8%	0.7%	161	225	71.6%
1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services.	62.0%	57.1%	58.5%	-5.7%	636	1,039	61.2%
2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy.	27.4%	24.3%	26.8%	-2.3%	162	553	29.3%
2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum.	70.3%	67.1%	69.3%	-1.5%	154	225	68.4%
3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1)	89.8%	89.9%	92.5%	3.0%			
3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1)	86.0%	88.6%	91.7%	6.6%			
4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were:							
a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1)	64.0%	66.7%	66.1%	3.3%	254	400	63.5%
b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile)	16.5%	15.7%	16.0%	-2.6%	74	400	18.5%
c. Obese (BMI at or greater than the 95th percentile).	17.8%	14.9%	15.4%	-13.7%	61	400	15.3%
d. Underweight (BMI less than the 5th percentile).	1.8%	2.7%	2.5%	43.9%	11	400	2.8%
4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3)							
a. Recommended prenatal weight gain	34.8%	29.7%	28.9%	-16.8%	39	205	19.0%
b. Excessive prenatal weight gain	45.9%	50.2%	49.1%	6.9%	118	205	57.6%
c. Inadequate prenatal weight gain	19.3%	20.2%	22.0%	13.9%	48	205	23.4%
5A. Percentage of women who participated in WIC who initiated breastfeeding.	50.1%	54.8%	42.8%	-14.6%	106	300	35.3%
5B. Percentage of infants who participated in WIC who were:							
a. Breastfeeding at 6 weeks of age (1)	32.9%	32.9%	31.8%	-3.3%	59	240	24.6%
b. Breastfeeding at 6 months of age	19.2%	17.1%	18.4%	-4.1%	33	240	13.8%

(1) Item appears also on other WCH Section Agreement Addenda.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI >=30.0): Adapted from the National Academy of Sciences. Weight Gain During Pregnancy: Reexamining the Guidelines. Washington, DC : National Academy Press, 2009.

SFY 2013-2014 AGREEMENT ADDENDA DATA

ATTACHMENT A: WIC QUALITY ASSURANCE DELIVERABLES

Duplin County

Process Outcome Objectives	2003 to 2005	2006 to 2008	2009 to 2011	% Change from 2003 to 2011	2011 Data		
					Num	Den	Percent
1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2)	73.3%	73.3%	71.8%	-2.1%	335	478	70.1%
1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services.	73.3%	78.0%	79.3%	8.2%	507	629	80.6%
1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services.	63.0%	63.6%	64.6%	2.7%	1,598	2,452	65.2%
2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy.	31.5%	30.6%	31.2%	-0.8%	329	1,087	30.3%
2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum.	83.1%	85.1%	78.7%	-5.3%	431	538	80.1%
3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1)	81.2%	88.3%	89.6%	10.3%			
3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1)	82.9%	83.6%	83.4%	0.6%			
4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were:							
a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1)	57.6%	57.9%	58.4%	1.5%	470	837	56.2%
b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile)	14.3%	16.4%	16.6%	16.1%	160	837	19.1%
c. Obese (BMI at or greater than the 95th percentile).	16.8%	19.0%	21.6%	29.2%	179	837	21.4%
d. Underweight (BMI less than the 5th percentile).	11.4%	6.7%	3.4%	-70.5%	28	837	3.3%
4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3)							
a. Recommended prenatal weight gain	34.2%	31.1%	27.5%	-19.5%	125	520	24.0%
b. Excessive prenatal weight gain	41.6%	40.7%	42.2%	1.5%	218	520	41.9%
c. Inadequate prenatal weight gain	24.2%	28.2%	30.3%	25.0%	177	520	34.0%
5A. Percentage of women who participated in WIC who initiated breastfeeding.	55.1%	51.6%	50.3%	-8.6%	354	718	49.3%
5B. Percentage of infants who participated in WIC who were:							
a. Breastfeeding at 6 weeks of age (1)	26.1%	25.2%	30.3%	16.1%	205	677	30.3%
b. Breastfeeding at 6 months of age	13.0%	11.8%	17.5%	34.6%	120	677	17.7%

(1) Item appears also on other WCH Section Agreement Addenda.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI >=30.0): Adapted from the National Academy of Sciences. Weight Gain During Pregnancy: Reexamining the Guidelines. Washington, DC : National Academy Press, 2009.

SFY 2013-2014 AGREEMENT ADDENDA DATA

ATTACHMENT A: WIC QUALITY ASSURANCE DELIVERABLES

Durham County

Process Outcome Objectives	2003 to 2005	2006 to 2008	2009 to 2011	% Change from 2003 to 2011	2011 Data		
					Num	Den	Percent
1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2)	67.4%	60.4%	63.4%	-6.0%	911	1,438	63.4%
1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services.	66.7%	65.7%	64.4%	-3.5%	1,501	2,406	62.4%
1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services.	56.3%	52.3%	56.0%	-0.6%	5,741	9,848	58.3%
2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy.	21.2%	21.0%	23.8%	12.0%	699	2,794	25.0%
2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum.	71.4%	75.4%	74.9%	5.0%	1,275	1,770	72.0%
3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1)							
3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1)							
4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were:							
a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1)	63.7%	56.5%	60.6%	-5.0%	1,494	2,500	59.8%
b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile)	12.7%	17.1%	16.2%	27.8%	444	2,500	17.8%
c. Obese (BMI at or greater than the 95th percentile).	17.6%	23.3%	19.5%	11.2%	478	2,500	19.1%
d. Underweight (BMI less than the 5th percentile).	6.0%	3.2%	3.7%	-39.0%	84	2,500	3.4%
4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3)							
a. Recommended prenatal weight gain	36.4%	33.7%	30.1%	-17.3%	499	1,885	26.5%
b. Excessive prenatal weight gain	38.2%	42.7%	46.8%	22.5%	898	1,885	47.6%
c. Inadequate prenatal weight gain	25.4%	23.5%	23.1%	-9.1%	488	1,885	25.9%
5A. Percentage of women who participated in WIC who initiated breastfeeding.	54.0%	61.4%	63.2%	17.2%	1,613	2,407	67.0%
5B. Percentage of infants who participated in WIC who were:							
a. Breastfeeding at 6 weeks of age (1)	37.1%	42.9%	40.1%	8.1%	1,010	2,412	41.9%
b. Breastfeeding at 6 months of age	20.3%	23.7%	23.1%	13.5%	579	2,412	24.0%

(1) Item appears also on other WCH Section Agreement Addenda.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI >=30.0): Adapted from the National Academy of Sciences. Weight Gain During Pregnancy: Reexamining the Guidelines. Washington, DC : National Academy Press, 2009.

SFY 2013-2014 AGREEMENT ADDENDA DATA

ATTACHMENT A: WIC QUALITY ASSURANCE DELIVERABLES

Edgecombe County

Process Outcome Objectives	2003 to 2005	2006 to 2008	2009 to 2011	% Change from 2003 to 2011	2011 Data		
					Num	Den	Percent
1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2)	82.9%	84.7%	84.4%	1.7%	414	508	81.5%
1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services.	76.5%	72.6%	73.7%	-3.6%	434	588	73.8%
1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services.	58.9%	53.9%	52.6%	-10.8%	1,371	2,568	53.4%
2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy.	27.6%	28.1%	25.2%	-8.8%	284	1,171	24.3%
2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum.	71.6%	70.1%	69.8%	-2.6%	368	516	71.3%
3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1)	90.7%	72.5%	70.7%	-22.0%			
3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1)	86.6%	70.5%	61.1%	-29.5%			
4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were:							
a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1)	65.3%	66.1%	67.5%	3.3%	220	319	69.0%
b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile)	15.7%	15.0%	13.8%	-11.9%	45	319	14.1%
c. Obese (BMI at or greater than the 95th percentile).	16.0%	15.5%	13.4%	-16.2%	39	319	12.2%
d. Underweight (BMI less than the 5th percentile).	3.0%	3.4%	5.3%	77.6%	15	319	4.7%
4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3)							
a. Recommended prenatal weight gain	34.2%	28.8%	31.1%	-9.1%	133	491	27.1%
b. Excessive prenatal weight gain	39.4%	43.7%	41.3%	4.8%	214	491	43.6%
c. Inadequate prenatal weight gain	26.4%	27.5%	27.6%	4.5%	144	491	29.3%
5A. Percentage of women who participated in WIC who initiated breastfeeding.	30.6%	29.1%	34.7%	13.2%	242	611	39.6%
5B. Percentage of infants who participated in WIC who were:							
a. Breastfeeding at 6 weeks of age (1)	16.3%	16.0%	13.0%	-20.7%	73	591	12.4%
b. Breastfeeding at 6 months of age	8.8%	6.4%	6.4%	-27.4%	29	591	4.9%

(1) Item appears also on other WCH Section Agreement Addenda.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI >=30.0): Adapted from the National Academy of Sciences. *Weight Gain During Pregnancy: Reexamining the Guidelines*. Washington, DC : National Academy Press, 2009.

SFY 2013-2014 AGREEMENT ADDENDA DATA

ATTACHMENT A: WIC QUALITY ASSURANCE DELIVERABLES

Forsyth County

Process Outcome Objectives	2003 to 2005	2006 to 2008	2009 to 2011	% Change from 2003 to 2011	2011 Data		
					Num	Den	Percent
1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2)	67.8%	62.6%	72.7%	7.2%	1,468	2,109	69.6%
1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services.	66.8%	69.5%	67.3%	0.6%	2,007	3,067	65.4%
1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services.	52.4%	53.0%	58.3%	11.1%	7,569	12,484	60.6%
2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy.	23.0%	29.5%	34.8%	51.7%	1,467	4,462	32.9%
2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum.	81.2%	84.2%	81.0%	-0.3%	2,089	2,609	80.1%
3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1)	100.0%		100.0%	0.0%			
3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1)	59.0%	41.9%	47.1%	-20.2%			
4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were:							
a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1)	66.2%	65.1%	65.5%	-1.1%	3,314	5,054	65.6%
b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile)	15.8%	16.8%	16.6%	4.9%	844	5,054	16.7%
c. Obese (BMI at or greater than the 95th percentile).	14.5%	15.3%	14.5%	0.3%	713	5,054	14.1%
d. Underweight (BMI less than the 5th percentile).	3.5%	2.8%	3.5%	-1.9%	183	5,054	3.6%
4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3)							
a. Recommended prenatal weight gain	32.9%	32.4%	29.5%	-10.4%	691	2,475	27.9%
b. Excessive prenatal weight gain	42.6%	44.3%	47.0%	10.5%	1,198	2,475	48.4%
c. Inadequate prenatal weight gain	24.5%	23.3%	23.5%	-4.3%	586	2,475	23.7%
5A. Percentage of women who participated in WIC who initiated breastfeeding.	67.2%	71.3%	70.7%	5.2%	2,295	3,237	70.9%
5B. Percentage of infants who participated in WIC who were:							
a. Breastfeeding at 6 weeks of age (1)	40.6%	45.0%	47.0%	15.6%	1,521	3,241	46.9%
b. Breastfeeding at 6 months of age	25.7%	24.7%	27.3%	5.9%	890	3,241	27.5%

(1) Item appears also on other WCH Section Agreement Addenda.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI >=30.0): Adapted from the National Academy of Sciences. Weight Gain During Pregnancy: Reexamining the Guidelines. Washington, DC : National Academy Press, 2009.

SFY 2013-2014 AGREEMENT ADDENDA DATA

ATTACHMENT A: WIC QUALITY ASSURANCE DELIVERABLES

Franklin County

Process Outcome Objectives	2003 to 2005	2006 to 2008	2009 to 2011	% Change from 2003 to 2011	2011 Data		
					Num	Den	Percent
1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2)	74.6%	71.0%	66.8%	-10.5%	202	323	62.5%
1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services.	72.5%	75.0%	68.2%	-6.0%	282	420	67.1%
1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services.	55.3%	54.3%	54.4%	-1.5%	1,014	1,856	54.6%
2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy.	25.6%	23.5%	24.0%	-6.3%	164	837	19.6%
2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum.	78.2%	78.8%	78.2%	0.1%	259	318	81.4%
3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1)	94.1%	94.0%	94.1%	-0.1%			
3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1)	88.3%	91.6%	90.0%	2.0%			
4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were:							
a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1)	65.5%	57.8%	60.8%	-7.2%	367	601	61.1%
b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile)	16.4%	20.0%	17.7%	8.1%	111	601	18.5%
c. Obese (BMI at or greater than the 95th percentile).	14.2%	19.6%	17.2%	21.5%	102	601	17.0%
d. Underweight (BMI less than the 5th percentile).	4.0%	2.6%	4.3%	8.1%	21	601	3.5%
4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3)							
a. Recommended prenatal weight gain	34.9%	31.4%	31.0%	-11.2%	101	317	31.9%
b. Excessive prenatal weight gain	40.7%	42.7%	44.9%	10.3%	132	317	41.6%
c. Inadequate prenatal weight gain	24.4%	25.9%	24.1%	-1.2%	84	317	26.5%
5A. Percentage of women who participated in WIC who initiated breastfeeding.	40.0%	41.6%	43.9%	9.8%	207	412	50.2%
5B. Percentage of infants who participated in WIC who were:							
a. Breastfeeding at 6 weeks of age (1)	26.0%	25.4%	25.9%	-0.2%	121	470	25.7%
b. Breastfeeding at 6 months of age	12.8%	13.3%	12.6%	-1.7%	59	470	12.6%

(1) Item appears also on other WCH Section Agreement Addenda.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI >=30.0): Adapted from the National Academy of Sciences. Weight Gain During Pregnancy: Reexamining the Guidelines. Washington, DC : National Academy Press, 2009.

SFY 2013-2014 AGREEMENT ADDENDA DATA

ATTACHMENT A: WIC QUALITY ASSURANCE DELIVERABLES

Gaston County

Process Outcome Objectives	2003 to 2005	2006 to 2008	2009 to 2011	% Change from 2003 to 2011	2011 Data		
					Num	Den	Percent
1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2)	72.1%	71.5%	71.5%	-0.8%	1,023	1,492	68.6%
1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services.	74.6%	74.2%	75.8%	1.6%	1,408	1,829	77.0%
1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services.	50.5%	49.4%	51.2%	1.4%	3,836	7,276	52.7%
2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy.	19.2%	24.6%	32.8%	70.3%	810	2,561	31.6%
2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum.	69.0%	81.3%	82.8%	20.0%	1,173	1,428	82.1%
3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1)	86.8%	80.8%	78.1%	-10.1%			
3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1)	89.4%	86.3%	84.2%	-5.8%			
4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were:							
a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1)	67.3%	66.6%	66.0%	-2.1%	932	1,438	64.8%
b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile)	14.5%	15.9%	16.1%	10.5%	219	1,438	15.2%
c. Obese (BMI at or greater than the 95th percentile).	14.5%	13.8%	13.9%	-4.6%	229	1,438	15.9%
d. Underweight (BMI less than the 5th percentile).	3.6%	3.8%	4.1%	14.9%	58	1,438	4.0%
4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3)							
a. Recommended prenatal weight gain	29.8%	29.0%	28.4%	-4.7%	440	1,498	29.4%
b. Excessive prenatal weight gain	47.3%	45.1%	44.4%	-6.1%	663	1,498	44.3%
c. Inadequate prenatal weight gain	22.9%	25.9%	27.2%	18.8%	395	1,498	26.4%
5A. Percentage of women who participated in WIC who initiated breastfeeding.	39.3%	49.6%	50.2%	27.5%	913	1,878	48.6%
5B. Percentage of infants who participated in WIC who were:							
a. Breastfeeding at 6 weeks of age (1)	22.5%	24.1%	26.1%	15.9%	553	1,893	29.2%
b. Breastfeeding at 6 months of age	10.2%	11.8%	12.4%	21.7%	269	1,893	14.2%

(1) Item appears also on other WCH Section Agreement Addenda.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI >=30.0): Adapted from the National Academy of Sciences. *Weight Gain During Pregnancy: Reexamining the Guidelines*. Washington, DC : National Academy Press, 2009.

SFY 2013-2014 AGREEMENT ADDENDA DATA

ATTACHMENT A: WIC QUALITY ASSURANCE DELIVERABLES

Graham County

Process Outcome Objectives	2003 to 2005	2006 to 2008	2009 to 2011	% Change from 2003 to 2011	2011 Data		
					Num	Den	Percent
1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2)	80.1%	77.6%	84.2%	5.1%	67	81	82.7%
1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services.	76.2%	76.9%	75.7%	-0.6%	66	91	72.5%
1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services.	69.7%	67.7%	77.3%	11.0%	263	323	81.4%
2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy.	62.0%	65.8%	62.5%	0.9%	87	146	59.6%
2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum.	81.7%	73.0%	78.9%	-3.4%	65	84	77.4%
3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1)	91.4%	93.8%	100.0%	9.4%			
3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1)	90.8%	94.7%	97.4%	7.2%			
4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were:							
a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1)	68.8%	65.0%	59.1%	-14.2%	103	175	58.9%
b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile)	11.8%	15.7%	23.2%	95.9%	43	175	24.6%
c. Obese (BMI at or greater than the 95th percentile).	13.8%	16.6%	14.1%	2.3%	23	175	13.1%
d. Underweight (BMI less than the 5th percentile).	5.6%	2.7%	3.6%	-34.5%	6	175	3.4%
4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3)							
a. Recommended prenatal weight gain	30.6%	29.9%	21.3%	-30.5%	13	76	17.1%
b. Excessive prenatal weight gain	51.1%	56.9%	57.9%	13.4%	47	76	61.8%
c. Inadequate prenatal weight gain	18.3%	13.2%	20.8%	13.7%	16	76	21.1%
5A. Percentage of women who participated in WIC who initiated breastfeeding.	26.2%	22.1%	38.9%	48.7%	47	96	49.0%
5B. Percentage of infants who participated in WIC who were:							
a. Breastfeeding at 6 weeks of age (1)	23.8%	12.6%	26.3%	10.6%	11	66	16.7%
b. Breastfeeding at 6 months of age	16.6%	56.8%	16.6%	-0.1%	1	66	1.5%

(1) Item appears also on other WCH Section Agreement Addenda.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI >=30.0): Adapted from the National Academy of Sciences. *Weight Gain During Pregnancy: Reexamining the Guidelines*. Washington, DC : National Academy Press, 2009.

SFY 2013-2014 AGREEMENT ADDENDA DATA

ATTACHMENT A: WIC QUALITY ASSURANCE DELIVERABLES

Granville-Vance District

Process Outcome Objectives	2003 to 2005	2006 to 2008	2009 to 2011	% Change from 2003 to 2011	2011 Data		
					Num	Den	Percent
1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2)	72.4%	75.4%	73.5%	1.5%	485	662	73.3%
1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services.	78.3%	74.8%	76.1%	-2.8%	674	881	76.5%
1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services.	58.7%	56.8%	57.7%	-1.8%	2,302	3,880	59.3%
2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy.	32.6%	32.7%	29.0%	-11.1%	408	1,500	27.2%
2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum.	81.7%	77.8%	73.9%	-9.5%	558	735	75.9%
3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1)	91.6%	93.0%	96.4%	5.2%			
3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1)	92.9%	90.6%	85.4%	-8.1%			
4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were:							
a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1)	64.3%	65.9%	65.8%	2.3%	901	1,374	65.6%
b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile)	16.1%	15.5%	12.7%	-20.8%	172	1,374	12.5%
c. Obese (BMI at or greater than the 95th percentile).	15.7%	13.9%	13.8%	-12.0%	193	1,374	14.0%
d. Underweight (BMI less than the 5th percentile).	3.9%	4.7%	7.6%	96.5%	108	1,374	7.9%
4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3)							
a. Recommended prenatal weight gain	29.0%	30.4%	31.5%	8.6%	221	721	30.7%
b. Excessive prenatal weight gain	39.8%	43.2%	42.3%	6.3%	309	721	42.9%
c. Inadequate prenatal weight gain	31.2%	26.4%	26.2%	-16.0%	191	721	26.5%
5A. Percentage of women who participated in WIC who initiated breastfeeding.	43.3%	52.9%	41.4%	-4.3%	408	922	44.3%
5B. Percentage of infants who participated in WIC who were:							
a. Breastfeeding at 6 weeks of age (1)	28.7%	34.5%	36.3%	26.6%	320	959	33.4%
b. Breastfeeding at 6 months of age	20.9%	22.5%	26.8%	28.5%	238	959	24.8%

(1) Item appears also on other WCH Section Agreement Addenda.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI >=30.0): Adapted from the National Academy of Sciences. *Weight Gain During Pregnancy: Reexamining the Guidelines*. Washington, DC : National Academy Press, 2009.

SFY 2013-2014 AGREEMENT ADDENDA DATA

ATTACHMENT A: WIC QUALITY ASSURANCE DELIVERABLES

Greene County

Process Outcome Objectives	2003 to 2005	2006 to 2008	2009 to 2011	% Change from 2003 to 2011	2011 Data		
					Num	Den	Percent
1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2)	83.3%	79.5%	81.6%	-2.0%	104	127	81.9%
1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services.	80.0%	77.3%	73.5%	-8.1%	130	181	71.8%
1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services.	65.6%	61.4%	62.8%	-4.3%	528	832	63.5%
2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy.	33.7%	29.0%	30.6%	-9.1%	128	437	29.3%
2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum.	76.3%	84.3%	84.3%	10.5%	150	178	84.3%
3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1)	96.0%	96.6%	98.9%	3.1%			
3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1)	93.2%	94.6%	96.5%	3.6%			
4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were:							
a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1)	65.4%	65.7%	62.0%	-5.1%	287	462	62.1%
b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile)	16.2%	16.2%	17.9%	10.3%	83	462	18.0%
c. Obese (BMI at or greater than the 95th percentile).	14.9%	14.9%	17.0%	13.6%	79	462	17.1%
d. Underweight (BMI less than the 5th percentile).	3.5%	3.2%	3.1%	-10.0%	13	462	2.8%
4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3)							
a. Recommended prenatal weight gain	31.3%	32.8%	32.1%	2.6%	49	166	29.5%
b. Excessive prenatal weight gain	42.9%	39.8%	42.6%	-0.5%	76	166	45.8%
c. Inadequate prenatal weight gain	25.9%	27.4%	25.3%	-2.3%	41	166	24.7%
5A. Percentage of women who participated in WIC who initiated breastfeeding.	46.8%	53.6%	59.8%	27.6%	138	215	64.2%
5B. Percentage of infants who participated in WIC who were:							
a. Breastfeeding at 6 weeks of age (1)	31.3%	35.0%	33.2%	6.2%	99	266	37.2%
b. Breastfeeding at 6 months of age	13.6%	17.5%	13.7%	1.1%	31	266	11.7%

(1) Item appears also on other WCH Section Agreement Addenda.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI >=30.0): Adapted from the National Academy of Sciences. Weight Gain During Pregnancy: Reexamining the Guidelines. Washington, DC : National Academy Press, 2009.

SFY 2013-2014 AGREEMENT ADDENDA DATA

ATTACHMENT A: WIC QUALITY ASSURANCE DELIVERABLES

Guilford County

Process Outcome Objectives	2003 to 2005	2006 to 2008	2009 to 2011	% Change from 2003 to 2011	2011 Data		
					Num	Den	Percent
1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2)	68.4%	65.6%	57.1%	-16.6%	1,604	2,873	55.8%
1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services.	70.7%	69.4%	66.9%	-5.4%	2,629	3,958	66.4%
1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services.	49.9%	49.7%	53.8%	7.8%	8,861	15,611	56.8%
2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy.	29.7%	28.2%	33.9%	14.2%	2,086	6,185	33.7%
2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum.	57.5%	78.5%	79.9%	39.1%	2,686	3,317	81.0%
3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1)							
3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1)							
4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were:							
a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1)	70.0%	66.1%	64.4%	-7.9%	2,676	4,155	64.4%
b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile)	13.8%	16.3%	15.8%	15.0%	624	4,155	15.0%
c. Obese (BMI at or greater than the 95th percentile).	12.5%	14.4%	15.5%	24.1%	667	4,155	16.1%
d. Underweight (BMI less than the 5th percentile).	3.8%	3.3%	4.3%	12.7%	188	4,155	4.5%
4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3)							
a. Recommended prenatal weight gain	30.5%	31.2%	29.1%	-4.5%	724	2,578	28.1%
b. Excessive prenatal weight gain	50.4%	48.7%	51.7%	2.6%	1,333	2,578	51.7%
c. Inadequate prenatal weight gain	19.2%	20.1%	19.2%	0.3%	521	2,578	20.2%
5A. Percentage of women who participated in WIC who initiated breastfeeding.	55.6%	65.0%	62.3%	12.1%	2,713	4,273	63.5%
5B. Percentage of infants who participated in WIC who were:							
a. Breastfeeding at 6 weeks of age (1)	32.9%	37.5%	39.7%	20.5%	1,688	4,083	41.3%
b. Breastfeeding at 6 months of age	17.1%	22.2%	23.9%	39.6%	1,052	4,083	25.8%

(1) Item appears also on other WCH Section Agreement Addenda.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI >=30.0): Adapted from the National Academy of Sciences. *Weight Gain During Pregnancy: Reexamining the Guidelines*. Washington, DC : National Academy Press, 2009.

SFY 2013-2014 AGREEMENT ADDENDA DATA

ATTACHMENT A: WIC QUALITY ASSURANCE DELIVERABLES

Halifax County

Process Outcome Objectives	2003 to 2005	2006 to 2008	2009 to 2011	% Change from 2003 to 2011	2011 Data		
					Num	Den	Percent
1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2)	81.4%	81.9%	82.3%	1.1%	381	464	82.1%
1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services.	74.3%	74.4%	71.2%	-4.1%	333	487	68.4%
1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services.	62.0%	60.6%	58.0%	-6.6%	1,219	2,102	58.0%
2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy.	29.0%	29.2%	25.9%	-10.9%	269	954	28.2%
2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum.	70.8%	75.6%	65.1%	-8.1%	314	480	65.4%
3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1)	69.1%	68.2%	57.6%	-16.7%			
3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1)	76.9%	65.5%	46.2%	-40.0%			
4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were:							
a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1)	65.4%	66.2%	64.9%	-0.8%	532	838	63.5%
b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile)	14.2%	15.0%	16.0%	12.7%	144	838	17.2%
c. Obese (BMI at or greater than the 95th percentile).	16.7%	14.4%	14.9%	-10.8%	129	838	15.4%
d. Underweight (BMI less than the 5th percentile).	3.7%	4.5%	4.3%	13.9%	33	838	3.9%
4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3)							
a. Recommended prenatal weight gain	30.1%	30.7%	28.4%	-5.7%	115	425	27.1%
b. Excessive prenatal weight gain	46.0%	47.5%	44.1%	-4.1%	178	425	41.9%
c. Inadequate prenatal weight gain	23.9%	21.8%	27.5%	15.0%	132	425	31.1%
5A. Percentage of women who participated in WIC who initiated breastfeeding.	33.5%	32.4%	34.5%	3.2%	200	570	35.1%
5B. Percentage of infants who participated in WIC who were:							
a. Breastfeeding at 6 weeks of age (1)	13.6%	13.7%	12.8%	-6.0%	59	545	10.8%
b. Breastfeeding at 6 months of age	4.7%	5.5%	5.2%	11.0%	25	545	4.6%

(1) Item appears also on other WCH Section Agreement Addenda.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI >=30.0): Adapted from the National Academy of Sciences. *Weight Gain During Pregnancy: Reexamining the Guidelines*. Washington, DC : National Academy Press, 2009.

SFY 2013-2014 AGREEMENT ADDENDA DATA

ATTACHMENT A: WIC QUALITY ASSURANCE DELIVERABLES

Harnett County

Process Outcome Objectives	2003 to 2005	2006 to 2008	2009 to 2011	% Change from 2003 to 2011	2011 Data		
					Num	Den	Percent
1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2)	72.1%	70.3%	68.6%	-4.9%	447	704	63.5%
1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services.	76.5%	74.8%	75.2%	-1.8%	695	916	75.9%
1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services.	57.4%	56.6%	57.7%	0.4%	2,284	3,855	59.2%
2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy.	35.3%	35.3%	32.3%	-8.6%	421	1,404	30.0%
2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum.	60.3%	69.8%	77.2%	28.0%	557	720	77.4%
3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1)	84.8%	86.2%	84.0%	-0.9%			
3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1)	80.2%	80.9%	80.4%	0.2%			
4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were:							
a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1)	66.5%	66.4%	67.7%	1.8%	781	1,136	68.8%
b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile)	16.6%	15.7%	12.9%	-21.9%	141	1,136	12.4%
c. Obese (BMI at or greater than the 95th percentile).	13.4%	14.3%	13.8%	2.7%	148	1,136	13.0%
d. Underweight (BMI less than the 5th percentile).	3.5%	3.7%	5.6%	60.2%	66	1,136	5.8%
4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3)							
a. Recommended prenatal weight gain	31.0%	30.0%	30.3%	-2.2%	239	829	28.8%
b. Excessive prenatal weight gain	47.3%	50.1%	48.7%	3.0%	401	829	48.4%
c. Inadequate prenatal weight gain	21.7%	19.9%	20.9%	-3.5%	189	829	22.8%
5A. Percentage of women who participated in WIC who initiated breastfeeding.	38.6%	46.7%	49.6%	28.4%	459	945	48.6%
5B. Percentage of infants who participated in WIC who were:							
a. Breastfeeding at 6 weeks of age (1)	23.4%	26.8%	25.5%	9.0%	236	996	23.7%
b. Breastfeeding at 6 months of age	10.8%	11.6%	11.2%	3.5%	114	996	11.4%

(1) Item appears also on other WCH Section Agreement Addenda.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI >=30.0): Adapted from the National Academy of Sciences. Weight Gain During Pregnancy: Reexamining the Guidelines. Washington, DC : National Academy Press, 2009.

SFY 2013-2014 AGREEMENT ADDENDA DATA

ATTACHMENT A: WIC QUALITY ASSURANCE DELIVERABLES

Haywood County

Process Outcome Objectives	2003 to 2005	2006 to 2008	2009 to 2011	% Change from 2003 to 2011	2011 Data		
					Num	Den	Percent
1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2)	73.8%	68.1%	77.1%	4.5%	278	360	77.2%
1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services.	68.0%	71.2%	72.2%	6.1%	312	427	73.1%
1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services.	55.2%	47.8%	56.6%	2.4%	1,013	1,664	60.9%
2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy.	47.2%	43.2%	49.4%	4.7%	348	714	48.7%
2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum.	58.1%	66.1%	76.2%	31.3%	301	381	79.0%
3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1)	68.0%	75.0%	100.0%	47.1%			
3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1)	77.7%	74.6%	63.4%	-18.4%			
4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were:							
a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1)	64.5%	65.5%	64.3%	-0.4%	357	540	66.1%
b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile)	18.0%	16.6%	17.3%	-4.2%	103	540	19.1%
c. Obese (BMI at or greater than the 95th percentile).	13.7%	14.2%	14.7%	7.7%	67	540	12.4%
d. Underweight (BMI less than the 5th percentile).	3.8%	3.7%	3.7%	-0.9%	13	540	2.4%
4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3)							
a. Recommended prenatal weight gain	30.4%	31.6%	30.4%	0.0%	110	359	30.6%
b. Excessive prenatal weight gain	50.9%	48.6%	50.5%	-0.8%	170	359	47.4%
c. Inadequate prenatal weight gain	18.7%	19.8%	19.1%	2.1%	79	359	22.0%
5A. Percentage of women who participated in WIC who initiated breastfeeding.	36.6%	43.0%	55.9%	52.6%	294	457	64.3%
5B. Percentage of infants who participated in WIC who were:							
a. Breastfeeding at 6 weeks of age (1)	25.7%	27.3%	19.2%	-25.5%	87	409	21.3%
b. Breastfeeding at 6 months of age	12.1%	13.1%	4.7%	-61.4%	19	409	4.6%

(1) Item appears also on other WCH Section Agreement Addenda.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI >=30.0): Adapted from the National Academy of Sciences. Weight Gain During Pregnancy: Reexamining the Guidelines. Washington, DC : National Academy Press, 2009.

SFY 2013-2014 AGREEMENT ADDENDA DATA

ATTACHMENT A: WIC QUALITY ASSURANCE DELIVERABLES

Henderson County

Process Outcome Objectives	2003 to 2005	2006 to 2008	2009 to 2011	% Change from 2003 to 2011	2011 Data		
					Num	Den	Percent
1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2)	53.5%	45.4%	48.3%	-9.6%	243	512	47.5%
1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services.	63.0%	62.7%	63.4%	0.7%	500	729	68.6%
1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services.	57.5%	50.3%	54.7%	-4.8%	1,893	3,319	57.0%
2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy.	36.2%	34.9%	38.5%	6.4%	399	977	40.8%
2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum.	64.3%	66.0%	71.9%	11.9%	405	560	72.3%
3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1)	88.0%	84.1%					
3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1)	80.8%	75.1%					
4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were:							
a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1)	66.5%	63.7%	63.8%	-4.0%	628	968	64.9%
b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile)	17.0%	18.3%	18.8%	10.8%	184	968	19.0%
c. Obese (BMI at or greater than the 95th percentile).	13.9%	15.5%	14.1%	2.0%	123	968	12.7%
d. Underweight (BMI less than the 5th percentile).	2.7%	2.5%	3.2%	20.0%	33	968	3.4%
4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3)							
a. Recommended prenatal weight gain	36.5%	33.0%	31.6%	-13.2%	132	452	29.2%
b. Excessive prenatal weight gain	43.9%	47.6%	47.3%	7.6%	212	452	46.9%
c. Inadequate prenatal weight gain	19.6%	19.5%	21.1%	7.6%	108	452	23.9%
5A. Percentage of women who participated in WIC who initiated breastfeeding.	68.9%	76.9%	82.3%	19.4%	615	754	81.6%
5B. Percentage of infants who participated in WIC who were:							
a. Breastfeeding at 6 weeks of age (1)	46.5%	48.7%	52.7%	13.1%	397	776	51.2%
b. Breastfeeding at 6 months of age	26.9%	28.5%	32.0%	19.0%	251	776	32.3%

(1) Item appears also on other WCH Section Agreement Addenda.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI >=30.0): Adapted from the National Academy of Sciences. Weight Gain During Pregnancy: Reexamining the Guidelines. Washington, DC : National Academy Press, 2009.

SFY 2013-2014 AGREEMENT ADDENDA DATA

ATTACHMENT A: WIC QUALITY ASSURANCE DELIVERABLES

Hertford County

Process Outcome Objectives	2003 to 2005	2006 to 2008	2009 to 2011	% Change from 2003 to 2011	2011 Data		
					Num	Den	Percent
1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2)	79.4%	78.4%	76.3%	-3.9%	124	178	69.7%
1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services.	77.2%	72.2%	68.2%	-11.6%	127	203	62.6%
1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services.	58.3%	53.3%	55.8%	-4.4%	520	905	57.5%
2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy.	27.8%	28.2%	25.1%	-9.7%	107	412	26.0%
2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum.	74.8%	84.8%	85.9%	14.9%	156	183	85.2%
3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1)	100.0%	100.0%	100.0%	0.0%			
3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1)	61.4%	86.0%	85.4%	39.2%			
4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were:							
a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1)	65.1%	66.5%	66.2%	1.7%	254	387	65.6%
b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile)	16.6%	15.1%	15.4%	-7.4%	59	387	15.2%
c. Obese (BMI at or greater than the 95th percentile).	15.2%	12.9%	12.7%	-16.3%	55	387	14.2%
d. Underweight (BMI less than the 5th percentile).	3.0%	5.5%	5.7%	85.6%	19	387	4.9%
4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3)							
a. Recommended prenatal weight gain	29.7%	30.6%	28.2%	-5.2%	48	164	29.3%
b. Excessive prenatal weight gain	50.4%	47.1%	44.2%	-12.2%	60	164	36.6%
c. Inadequate prenatal weight gain	19.9%	22.3%	27.6%	38.5%	56	164	34.1%
5A. Percentage of women who participated in WIC who initiated breastfeeding.	33.8%	32.5%	30.6%	-9.4%	65	234	27.8%
5B. Percentage of infants who participated in WIC who were:							
a. Breastfeeding at 6 weeks of age (1)	13.9%	18.9%	17.7%	27.5%	41	236	17.4%
b. Breastfeeding at 6 months of age	4.2%	6.2%	7.4%	76.3%	17	236	7.2%

(1) Item appears also on other WCH Section Agreement Addenda.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI >=30.0): Adapted from the National Academy of Sciences. Weight Gain During Pregnancy: Reexamining the Guidelines. Washington, DC : National Academy Press, 2009.

SFY 2013-2014 AGREEMENT ADDENDA DATA

ATTACHMENT A: WIC QUALITY ASSURANCE DELIVERABLES

Hoke County

Process Outcome Objectives	2003 to 2005	2006 to 2008	2009 to 2011	% Change from 2003 to 2011	2011 Data		
					Num	Den	Percent
1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2)	78.4%	76.4%	71.4%	-9.0%	236	339	69.6%
1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services.	77.1%	74.2%	71.8%	-6.9%	336	481	69.9%
1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services.	58.9%	54.6%	54.9%	-6.8%	1,072	1,865	57.5%
2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy.	33.1%	26.3%	26.1%	-21.1%	245	940	26.1%
2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum.	71.0%	73.1%	75.3%	5.9%	321	447	71.8%
3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1)	97.3%	95.7%	97.6%	0.3%			
3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1)	95.2%	92.8%	94.6%	-0.6%			
4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were:							
a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1)	64.6%	60.0%	61.7%	-4.5%	446	692	64.5%
b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile)	14.9%	17.3%	17.9%	20.0%	109	692	15.8%
c. Obese (BMI at or greater than the 95th percentile).	16.4%	19.8%	17.8%	8.8%	114	692	16.5%
d. Underweight (BMI less than the 5th percentile).	4.1%	2.9%	2.6%	-37.2%	23	692	3.3%
4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3)							
a. Recommended prenatal weight gain	31.9%	33.7%	28.1%	-11.6%	130	464	28.0%
b. Excessive prenatal weight gain	47.7%	48.2%	47.5%	-0.4%	215	464	46.3%
c. Inadequate prenatal weight gain	20.4%	18.1%	24.3%	19.2%	119	464	25.6%
5A. Percentage of women who participated in WIC who initiated breastfeeding.	45.8%	46.1%	51.3%	12.1%	331	621	53.3%
5B. Percentage of infants who participated in WIC who were:							
a. Breastfeeding at 6 weeks of age (1)	24.9%	27.3%	27.9%	12.1%	175	580	30.2%
b. Breastfeeding at 6 months of age	12.6%	14.3%	16.1%	27.6%	107	580	18.4%

(1) Item appears also on other WCH Section Agreement Addenda.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI >=30.0): Adapted from the National Academy of Sciences. Weight Gain During Pregnancy: Reexamining the Guidelines. Washington, DC : National Academy Press, 2009.

SFY 2013-2014 AGREEMENT ADDENDA DATA

ATTACHMENT A: WIC QUALITY ASSURANCE DELIVERABLES

Hyde County

Process Outcome Objectives	2003 to 2005	2006 to 2008	2009 to 2011	% Change from 2003 to 2011	2011 Data		
					Num	Den	Percent
1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2)	76.8%	70.2%	77.4%	0.7%	26	32	81.3%
1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services.	61.3%	70.4%	79.3%	29.4%	30	39	76.9%
1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services.	67.0%	68.2%	71.2%	6.3%	109	151	72.2%
2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy.	40.0%	33.8%	28.2%	-29.6%	17	80	21.3%
2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum.	64.5%	78.3%	62.5%	-3.1%	21	32	65.6%
3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1)	50.0%	100.0%	80.0%	60.0%			
3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1)	88.6%	93.9%	100.0%	12.9%			
4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were:							
a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1)	62.5%	68.7%	62.6%	0.2%	56	81	69.1%
b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile)	11.6%	12.0%	12.6%	8.3%	10	81	12.3%
c. Obese (BMI at or greater than the 95th percentile).	19.8%	11.2%	18.9%	-4.6%	14	81	17.3%
d. Underweight (BMI less than the 5th percentile).	6.0%	8.0%	5.9%	-2.5%	1	81	1.2%
4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3)							
a. Recommended prenatal weight gain	25.8%	22.7%	39.7%	53.8%	8	26	30.8%
b. Excessive prenatal weight gain	51.7%	52.3%	48.7%	-5.7%	14	26	53.8%
c. Inadequate prenatal weight gain	22.5%	25.0%	11.5%	-48.7%	4	26	15.4%
5A. Percentage of women who participated in WIC who initiated breastfeeding.	51.6%	47.2%	39.7%	-23.1%	17	43	39.5%
5B. Percentage of infants who participated in WIC who were:							
a. Breastfeeding at 6 weeks of age (1)	30.3%	34.3%	27.8%	-8.4%	9	39	23.1%
b. Breastfeeding at 6 months of age	16.8%	18.0%	18.3%	8.8%	7	39	17.9%

(1) Item appears also on other WCH Section Agreement Addenda.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI >=30.0): Adapted from the National Academy of Sciences. Weight Gain During Pregnancy: Reexamining the Guidelines. Washington, DC : National Academy Press, 2009.

SFY 2013-2014 AGREEMENT ADDENDA DATA

ATTACHMENT A: WIC QUALITY ASSURANCE DELIVERABLES

Iredell County

Process Outcome Objectives	2003 to 2005	2006 to 2008	2009 to 2011	% Change from 2003 to 2011	2011 Data		
					Num	Den	Percent
1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2)	72.5%	75.7%	76.7%	5.8%	684	889	76.9%
1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services.	70.4%	72.9%	74.6%	5.9%	834	1,055	79.1%
1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services.	55.6%	56.7%	60.5%	9.0%	2,885	4,492	64.2%
2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy.	24.6%	30.1%	30.3%	23.4%	573	1,710	33.5%
2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum.	52.8%	58.1%	78.2%	48.1%	747	906	82.5%
3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1)	85.2%	88.4%	95.5%	12.0%			
3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1)	69.2%	75.3%	77.5%	11.9%			
4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were:							
a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1)	69.4%	68.5%	64.0%	-7.8%	920	1,460	63.0%
b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile)	15.1%	13.9%	17.2%	13.9%	268	1,460	18.4%
c. Obese (BMI at or greater than the 95th percentile).	12.6%	14.3%	15.9%	26.4%	231	1,460	15.8%
d. Underweight (BMI less than the 5th percentile).	2.9%	3.4%	2.9%	-0.7%	41	1,460	2.8%
4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3)							
a. Recommended prenatal weight gain	32.1%	30.2%	31.9%	-0.6%	255	871	29.3%
b. Excessive prenatal weight gain	49.0%	49.5%	46.8%	-4.4%	409	871	47.0%
c. Inadequate prenatal weight gain	18.9%	20.4%	21.3%	12.4%	207	871	23.8%
5A. Percentage of women who participated in WIC who initiated breastfeeding.	48.1%	53.2%	60.5%	25.8%	668	1,123	59.5%
5B. Percentage of infants who participated in WIC who were:							
a. Breastfeeding at 6 weeks of age (1)	23.5%	32.6%	31.1%	32.4%	393	1,178	33.4%
b. Breastfeeding at 6 months of age	13.0%	17.8%	17.1%	32.2%	212	1,178	18.0%

(1) Item appears also on other WCH Section Agreement Addenda.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI >=30.0): Adapted from the National Academy of Sciences. Weight Gain During Pregnancy: Reexamining the Guidelines. Washington, DC : National Academy Press, 2009.

SFY 2013-2014 AGREEMENT ADDENDA DATA

ATTACHMENT A: WIC QUALITY ASSURANCE DELIVERABLES

Jackson County

Process Outcome Objectives	2003 to 2005	2006 to 2008	2009 to 2011	% Change from 2003 to 2011	2011 Data		
					Num	Den	Percent
1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2)	82.7%	79.5%	78.1%	-5.6%	179	247	72.5%
1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services.	68.8%	69.4%	71.2%	3.4%	210	293	71.7%
1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services.	68.1%	62.6%	65.5%	-3.8%	804	1,190	67.6%
2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy.	47.3%	49.6%	46.7%	-1.2%	211	490	43.1%
2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum.	61.2%	64.4%	64.4%	5.2%	135	231	58.4%
3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1)	33.3%	50.0%					
3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1)	70.9%	62.7%	61.5%	-13.2%			
4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were:							
a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1)	60.5%	59.6%	65.0%	7.5%	241	350	68.9%
b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile)	18.9%	18.8%	15.4%	-18.6%	53	350	15.1%
c. Obese (BMI at or greater than the 95th percentile).	17.5%	18.5%	16.4%	-6.2%	46	350	13.1%
d. Underweight (BMI less than the 5th percentile).	3.1%	3.1%	3.2%	1.7%	10	350	2.9%
4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3)							
a. Recommended prenatal weight gain	33.0%	29.1%	30.8%	-6.6%	78	264	29.5%
b. Excessive prenatal weight gain	49.7%	54.2%	50.5%	1.4%	129	264	48.9%
c. Inadequate prenatal weight gain	17.3%	16.8%	18.7%	8.5%	57	264	21.6%
5A. Percentage of women who participated in WIC who initiated breastfeeding.	51.4%	46.6%	52.1%	1.5%	160	286	55.9%
5B. Percentage of infants who participated in WIC who were:							
a. Breastfeeding at 6 weeks of age (1)	33.6%	34.0%	38.5%	14.3%	109	275	39.6%
b. Breastfeeding at 6 months of age	16.7%	18.6%	24.1%	43.9%	79	275	28.7%

(1) Item appears also on other WCH Section Agreement Addenda.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI >=30.0): Adapted from the National Academy of Sciences. Weight Gain During Pregnancy: Reexamining the Guidelines. Washington, DC : National Academy Press, 2009.

SFY 2013-2014 AGREEMENT ADDENDA DATA

ATTACHMENT A: WIC QUALITY ASSURANCE DELIVERABLES

Johnston County

Process Outcome Objectives	2003 to 2005	2006 to 2008	2009 to 2011	% Change from 2003 to 2011	2011 Data		
					Num	Den	Percent
1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2)	64.4%	56.3%	59.0%	-8.3%	610	999	61.1%
1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services.	70.5%	68.9%	70.6%	0.2%	999	1,430	69.9%
1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services.	51.9%	46.8%	51.4%	-0.9%	3,174	5,934	53.5%
2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy.	24.8%	14.1%	18.6%	-24.9%	415	1,486	27.9%
2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum.	65.9%	63.0%	76.1%	15.5%	734	971	75.6%
3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1)	72.9%	66.8%	66.3%	-9.1%			
3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1)	77.9%	60.8%	59.7%	-23.4%			
4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were:							
a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1)	62.4%	61.6%	64.3%	3.0%	866	1,397	62.0%
b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile)	17.5%	16.8%	15.1%	-13.7%	229	1,397	16.4%
c. Obese (BMI at or greater than the 95th percentile).	16.7%	18.0%	15.8%	-5.7%	254	1,397	18.2%
d. Underweight (BMI less than the 5th percentile).	3.3%	3.6%	4.8%	44.7%	48	1,397	3.4%
4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3)							
a. Recommended prenatal weight gain	32.3%	37.4%	31.0%	-4.0%	300	1,067	28.1%
b. Excessive prenatal weight gain	43.6%	43.6%	50.0%	14.7%	530	1,067	49.7%
c. Inadequate prenatal weight gain	24.1%	19.0%	19.0%	-21.2%	237	1,067	22.2%
5A. Percentage of women who participated in WIC who initiated breastfeeding.	34.2%	47.0%	50.6%	47.7%	668	1,308	51.1%
5B. Percentage of infants who participated in WIC who were:							
a. Breastfeeding at 6 weeks of age (1)	20.9%	28.2%	29.7%	42.0%	366	1,293	28.3%
b. Breastfeeding at 6 months of age	10.5%	15.1%	16.5%	57.2%	210	1,293	16.2%

(1) Item appears also on other WCH Section Agreement Addenda.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI >=30.0): Adapted from the National Academy of Sciences. Weight Gain During Pregnancy: Reexamining the Guidelines. Washington, DC : National Academy Press, 2009.

SFY 2013-2014 AGREEMENT ADDENDA DATA

ATTACHMENT A: WIC QUALITY ASSURANCE DELIVERABLES

Jones County

Process Outcome Objectives	2003 to 2005	2006 to 2008	2009 to 2011	% Change from 2003 to 2011	2011 Data		
					Num	Den	Percent
1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2)	87.8%	81.8%	85.6%	-2.4%	61	69	88.4%
1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services.	79.1%	81.1%	80.1%	1.3%	58	81	71.6%
1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services.	71.7%	68.5%	72.6%	1.3%	241	320	75.3%
2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy.	47.1%	46.2%	46.0%	-2.3%	78	193	40.4%
2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum.	76.7%	76.9%	72.0%	-6.2%	57	82	69.5%
3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1)	90.8%	96.2%	100.0%	10.1%			
3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1)	85.8%	86.5%	89.1%	3.8%			
4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were:							
a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1)	68.3%	72.2%	67.8%	-0.8%	31	45	68.9%
b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile)	17.7%	14.2%	16.1%	-9.2%	5	45	11.1%
c. Obese (BMI at or greater than the 95th percentile).	11.1%	10.5%	10.4%	-6.0%	5	45	11.1%
d. Underweight (BMI less than the 5th percentile).	2.9%	3.0%	5.7%	97.3%	4	45	8.9%
4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3)							
a. Recommended prenatal weight gain	34.0%	35.7%	25.2%	-25.8%	21	90	23.3%
b. Excessive prenatal weight gain	45.5%	48.8%	46.6%	2.6%	43	90	47.8%
c. Inadequate prenatal weight gain	20.6%	15.5%	28.2%	36.8%	26	90	28.9%
5A. Percentage of women who participated in WIC who initiated breastfeeding.	44.8%	46.4%	47.6%	6.3%	42	102	41.2%
5B. Percentage of infants who participated in WIC who were:							
a. Breastfeeding at 6 weeks of age (1)	28.4%	30.2%	35.9%	26.4%	37	98	37.8%
b. Breastfeeding at 6 months of age	13.7%	17.1%	18.0%	31.0%	25	98	25.5%

(1) Item appears also on other WCH Section Agreement Addenda.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI >=30.0): Adapted from the National Academy of Sciences. Weight Gain During Pregnancy: Reexamining the Guidelines. Washington, DC : National Academy Press, 2009.

SFY 2013-2014 AGREEMENT ADDENDA DATA

ATTACHMENT A: WIC QUALITY ASSURANCE DELIVERABLES

Lee County

Process Outcome Objectives	2003 to 2005	2006 to 2008	2009 to 2011	% Change from 2003 to 2011	2011 Data		
					Num	Den	Percent
1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2)	69.4%	71.5%	56.1%	-19.3%	235	415	56.6%
1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services.	77.0%	78.1%	77.5%	0.6%	499	621	80.4%
1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services.	59.8%	59.5%	62.9%	5.1%	1,725	2,681	64.3%
2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy.	30.1%	23.3%	21.0%	-30.3%	251	1,003	25.0%
2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum.	81.7%	73.2%	68.2%	-16.5%	314	467	67.2%
3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1)	95.8%	100.0%					
3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1)	75.6%	79.4%	76.5%	1.2%			
4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were:							
a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1)	63.6%	67.4%	63.8%	0.2%	636	985	64.6%
b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile)	16.6%	14.8%	16.3%	-1.4%	158	985	16.0%
c. Obese (BMI at or greater than the 95th percentile).	17.6%	15.4%	17.1%	-2.6%	161	985	16.3%
d. Underweight (BMI less than the 5th percentile).	2.2%	2.4%	2.8%	26.3%	30	985	3.0%
4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3)							
a. Recommended prenatal weight gain	29.7%	30.8%	32.1%	8.3%	156	499	31.3%
b. Excessive prenatal weight gain	43.3%	47.8%	44.6%	3.1%	227	499	45.5%
c. Inadequate prenatal weight gain	27.1%	21.4%	23.3%	-14.0%	116	499	23.2%
5A. Percentage of women who participated in WIC who initiated breastfeeding.	50.6%	55.2%	50.3%	-0.5%	300	559	53.7%
5B. Percentage of infants who participated in WIC who were:							
a. Breastfeeding at 6 weeks of age (1)	31.3%	31.7%	30.3%	-3.1%	188	625	30.1%
b. Breastfeeding at 6 months of age	17.0%	15.9%	17.1%	0.6%	94	625	15.0%

(1) Item appears also on other WCH Section Agreement Addenda.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI >=30.0): Adapted from the National Academy of Sciences. Weight Gain During Pregnancy: Reexamining the Guidelines. Washington, DC : National Academy Press, 2009.

SFY 2013-2014 AGREEMENT ADDENDA DATA

ATTACHMENT A: WIC QUALITY ASSURANCE DELIVERABLES

Lenoir County

Process Outcome Objectives	2003 to 2005	2006 to 2008	2009 to 2011	% Change from 2003 to 2011	2011 Data		
					Num	Den	Percent
1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2)	74.3%	71.9%	71.8%	-3.4%	343	462	74.2%
1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services.	76.5%	75.2%	73.7%	-3.6%	410	544	75.4%
1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services.	55.4%	51.9%	51.2%	-7.6%	1,249	2,385	52.4%
2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy.	35.5%	27.3%	26.2%	-26.2%	267	1,027	26.0%
2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum.	44.8%	57.9%	74.5%	66.4%	376	471	79.8%
3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1)	86.2%	58.7%	50.0%	-42.0%			
3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1)	95.8%	79.5%	73.3%	-23.6%			
4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were:							
a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1)	69.8%	68.9%	66.5%	-4.7%	458	723	63.3%
b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile)	11.0%	10.3%	14.2%	28.9%	121	723	16.7%
c. Obese (BMI at or greater than the 95th percentile).	10.4%	11.1%	12.7%	22.3%	113	723	15.6%
d. Underweight (BMI less than the 5th percentile).	8.8%	9.7%	6.6%	-24.8%	31	723	4.3%
4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3)							
a. Recommended prenatal weight gain	30.5%	27.2%	27.8%	-8.8%	133	470	28.3%
b. Excessive prenatal weight gain	46.4%	48.9%	47.0%	1.3%	240	470	51.1%
c. Inadequate prenatal weight gain	23.1%	23.9%	25.2%	9.1%	97	470	20.6%
5A. Percentage of women who participated in WIC who initiated breastfeeding.	32.4%	39.9%	48.4%	49.3%	325	605	53.7%
5B. Percentage of infants who participated in WIC who were:							
a. Breastfeeding at 6 weeks of age (1)	18.7%	20.0%	22.8%	21.9%	120	565	21.2%
b. Breastfeeding at 6 months of age	8.2%	9.5%	10.5%	27.1%	51	565	9.0%

(1) Item appears also on other WCH Section Agreement Addenda.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI >=30.0): Adapted from the National Academy of Sciences. Weight Gain During Pregnancy: Reexamining the Guidelines. Washington, DC : National Academy Press, 2009.

SFY 2013-2014 AGREEMENT ADDENDA DATA

ATTACHMENT A: WIC QUALITY ASSURANCE DELIVERABLES

Lincoln County

Process Outcome Objectives	2003 to 2005	2006 to 2008	2009 to 2011	% Change from 2003 to 2011	2011 Data		
					Num	Den	Percent
1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2)	75.2%	74.2%	70.6%	-6.1%	263	382	68.8%
1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services.	74.3%	76.5%	74.3%	-0.1%	376	498	75.5%
1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services.	55.2%	59.3%	61.7%	11.7%	1,418	2,220	63.9%
2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy.	37.4%	38.2%	36.1%	-3.4%	339	907	37.4%
2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum.	73.4%	72.9%	77.8%	6.0%	362	465	77.8%
3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1)	65.2%	58.8%	80.0%	22.7%			
3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1)	61.8%	59.4%	57.1%	-7.5%			
4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were:							
a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1)	67.2%	72.8%	70.6%	5.0%	561	795	70.6%
b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile)	10.1%	9.4%	10.6%	4.4%	79	795	9.9%
c. Obese (BMI at or greater than the 95th percentile).	8.6%	9.8%	10.0%	15.9%	75	795	9.4%
d. Underweight (BMI less than the 5th percentile).	14.0%	7.9%	8.8%	-37.0%	80	795	10.1%
4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3)							
a. Recommended prenatal weight gain	29.6%	30.3%	30.8%	3.8%	116	423	27.4%
b. Excessive prenatal weight gain	49.8%	48.8%	46.2%	-7.1%	205	423	48.5%
c. Inadequate prenatal weight gain	20.6%	20.8%	23.0%	11.7%	102	423	24.1%
5A. Percentage of women who participated in WIC who initiated breastfeeding.	49.2%	49.1%	62.2%	26.6%	346	578	59.9%
5B. Percentage of infants who participated in WIC who were:							
a. Breastfeeding at 6 weeks of age (1)	26.4%	28.3%	31.5%	19.4%	179	572	31.3%
b. Breastfeeding at 6 months of age	10.2%	11.5%	14.2%	39.3%	80	572	14.0%

(1) Item appears also on other WCH Section Agreement Addenda.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI >=30.0): Adapted from the National Academy of Sciences. Weight Gain During Pregnancy: Reexamining the Guidelines. Washington, DC : National Academy Press, 2009.

SFY 2013-2014 AGREEMENT ADDENDA DATA

ATTACHMENT A: WIC QUALITY ASSURANCE DELIVERABLES

Macon County

Process Outcome Objectives	2003 to 2005	2006 to 2008	2009 to 2011	% Change from 2003 to 2011	2011 Data		
					Num	Den	Percent
1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2)	78.6%	72.3%	55.5%	-29.3%	124	168	73.8%
1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services.	73.1%	71.4%	73.1%	0.0%	177	254	69.7%
1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services.	61.1%	57.7%	62.9%	2.9%	752	1,163	64.7%
2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy.	49.9%	40.2%	38.5%	-23.0%	186	496	37.5%
2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum.	50.1%	71.3%	74.9%	49.4%	184	244	75.4%
3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1)	76.5%	89.5%	100.0%	30.8%			
3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1)	79.1%	77.7%	95.6%	20.8%			
4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were:							
a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1)	64.7%	67.9%	60.8%	-6.0%	73	137	53.3%
b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile)	16.5%	16.8%	18.9%	14.1%	20	137	14.6%
c. Obese (BMI at or greater than the 95th percentile).	15.9%	13.9%	16.7%	4.9%	35	137	25.5%
d. Underweight (BMI less than the 5th percentile).	2.8%	1.3%	3.6%	28.4%	9	137	6.6%
4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3)							
a. Recommended prenatal weight gain	32.2%	38.5%	31.7%	-1.3%	55	201	27.4%
b. Excessive prenatal weight gain	47.0%	44.1%	47.9%	2.1%	98	201	48.8%
c. Inadequate prenatal weight gain	20.9%	17.4%	20.3%	-2.7%	48	201	23.9%
5A. Percentage of women who participated in WIC who initiated breastfeeding.	65.0%	68.7%	64.8%	-0.4%	192	292	65.8%
5B. Percentage of infants who participated in WIC who were:							
a. Breastfeeding at 6 weeks of age (1)	42.1%	39.0%	40.5%	-3.7%	113	268	42.2%
b. Breastfeeding at 6 months of age	21.6%	22.5%	26.8%	24.2%	81	268	30.2%

(1) Item appears also on other WCH Section Agreement Addenda.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI >=30.0): Adapted from the National Academy of Sciences. Weight Gain During Pregnancy: Reexamining the Guidelines. Washington, DC : National Academy Press, 2009.

SFY 2013-2014 AGREEMENT ADDENDA DATA

ATTACHMENT A: WIC QUALITY ASSURANCE DELIVERABLES

Madison County

Process Outcome Objectives	2003 to 2005	2006 to 2008	2009 to 2011	% Change from 2003 to 2011	2011 Data		
					Num	Den	Percent
1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2)	80.7%	73.3%	69.8%	-13.5%	82	118	69.5%
1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services.	70.6%	66.8%	64.1%	-9.2%	95	153	62.1%
1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services.	64.6%	53.7%	55.9%	-13.4%	342	579	59.1%
2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy.	46.3%	47.8%	41.4%	-10.5%	135	318	42.5%
2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum.	72.6%	69.1%	73.5%	1.2%	103	146	70.5%
3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1)		93.8%	91.7%				
3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1)		84.6%	71.4%				
4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were:							
a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1)	62.3%	66.1%	67.2%	7.8%	43	69	62.3%
b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile)	16.3%	15.7%	14.7%	-9.9%	10	69	14.5%
c. Obese (BMI at or greater than the 95th percentile).	19.2%	14.4%	10.8%	-43.6%	10	69	14.5%
d. Underweight (BMI less than the 5th percentile).	2.2%	3.8%	7.3%	232.4%	6	69	8.7%
4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3)							
a. Recommended prenatal weight gain	28.3%	33.1%	30.7%	8.4%	33	148	22.3%
b. Excessive prenatal weight gain	46.9%	48.9%	48.9%	4.3%	79	148	53.4%
c. Inadequate prenatal weight gain	24.8%	18.0%	20.4%	-17.8%	36	148	24.3%
5A. Percentage of women who participated in WIC who initiated breastfeeding.	57.4%	49.2%	63.4%	10.6%	128	191	67.0%
5B. Percentage of infants who participated in WIC who were:							
a. Breastfeeding at 6 weeks of age (1)	30.7%	39.3%	50.8%	65.8%	65	158	41.1%
b. Breastfeeding at 6 months of age	16.9%	24.0%	32.4%	91.4%	32	158	20.3%

(1) Item appears also on other WCH Section Agreement Addenda.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI >=30.0): Adapted from the National Academy of Sciences. Weight Gain During Pregnancy: Reexamining the Guidelines. Washington, DC : National Academy Press, 2009.

SFY 2013-2014 AGREEMENT ADDENDA DATA

ATTACHMENT A: WIC QUALITY ASSURANCE DELIVERABLES

Martin-Tyrrell-Washington District

Process Outcome Objectives	2003 to 2005	2006 to 2008	2009 to 2011	% Change from 2003 to 2011	2011 Data		
					Num	Den	Percent
1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2)	80.0%	83.6%	85.1%	6.3%	246	305	80.7%
1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services.	74.6%	70.4%	78.6%	5.3%	301	357	84.3%
1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services.	65.0%	61.9%	61.9%	-4.7%	1,029	1,664	61.8%
2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy.	37.4%	32.3%	30.6%	-18.1%	242	911	26.6%
2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum.	79.1%	82.0%	79.7%	0.8%	247	313	78.9%
3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1)	97.1%	94.8%	98.5%	1.4%			
3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1)	94.7%	92.5%	93.3%	-1.5%			
4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were:							
a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1)	65.7%	65.5%	66.5%	1.2%	491	739	66.4%
b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile)	13.8%	13.4%	12.3%	-11.1%	100	739	13.5%
c. Obese (BMI at or greater than the 95th percentile).	12.7%	14.2%	14.6%	14.7%	111	739	15.0%
d. Underweight (BMI less than the 5th percentile).	7.8%	7.0%	6.7%	-14.6%	37	739	5.0%
4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3)							
a. Recommended prenatal weight gain	29.4%	30.0%	28.9%	-1.6%	79	295	26.8%
b. Excessive prenatal weight gain	42.9%	43.3%	43.3%	1.1%	128	295	43.4%
c. Inadequate prenatal weight gain	27.7%	26.6%	27.7%	0.1%	88	295	29.8%
5A. Percentage of women who participated in WIC who initiated breastfeeding.	41.3%	38.1%	26.6%	-35.5%	82	372	22.0%
5B. Percentage of infants who participated in WIC who were:							
a. Breastfeeding at 6 weeks of age (1)	24.6%	23.3%	14.8%	-39.9%	40	354	11.3%
b. Breastfeeding at 6 months of age	11.9%	11.4%	4.7%	-60.8%	15	354	4.2%

(1) Item appears also on other WCH Section Agreement Addenda.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI >=30.0): Adapted from the National Academy of Sciences. *Weight Gain During Pregnancy: Reexamining the Guidelines*. Washington, DC : National Academy Press, 2009.

SFY 2013-2014 AGREEMENT ADDENDA DATA

ATTACHMENT A: WIC QUALITY ASSURANCE DELIVERABLES

Mecklenburg County

Process Outcome Objectives	2003 to 2005	2006 to 2008	2009 to 2011	% Change from 2003 to 2011	2011 Data		
					Num	Den	Percent
1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2)	57.6%	62.4%	63.5%	10.1%	3,093	4,948	62.5%
1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services.	59.0%	61.9%	64.6%	9.5%	4,937	7,573	65.2%
1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services.	44.4%	42.7%	47.6%	7.2%	15,793	32,510	48.6%
2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy.	16.7%	16.4%	21.6%	29.2%	2,002	8,134	24.6%
2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum.	49.6%	47.8%	44.7%	-10.0%	2,715	5,260	51.6%
3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1)	18.8%	36.4%	66.7%	255.6%			
3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1)	20.8%	18.9%	40.9%	96.4%			
4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were:							
a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1)	63.3%	62.8%	61.1%	-3.4%	4,826	7,934	60.8%
b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile)	17.0%	17.5%	17.7%	3.7%	1,415	7,934	17.8%
c. Obese (BMI at or greater than the 95th percentile).	16.0%	16.6%	18.0%	12.5%	1,445	7,934	18.2%
d. Underweight (BMI less than the 5th percentile).	3.8%	3.0%	3.2%	-13.6%	248	7,934	3.1%
4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3)							
a. Recommended prenatal weight gain	31.2%	33.1%	31.2%	0.0%	1,349	4,647	29.0%
b. Excessive prenatal weight gain	49.5%	47.4%	46.9%	-5.3%	2,072	4,647	44.6%
c. Inadequate prenatal weight gain	19.2%	19.5%	21.9%	13.7%	1,226	4,647	26.4%
5A. Percentage of women who participated in WIC who initiated breastfeeding.	49.9%	55.5%	53.7%	7.7%	3,325	6,295	52.8%
5B. Percentage of infants who participated in WIC who were:							
a. Breastfeeding at 6 weeks of age (1)	29.7%	31.8%	33.8%	13.9%	2,363	7,199	32.8%
b. Breastfeeding at 6 months of age	15.8%	16.0%	18.8%	19.1%	1,320	7,199	18.3%

(1) Item appears also on other WCH Section Agreement Addenda.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI >=30.0): Adapted from the National Academy of Sciences. Weight Gain During Pregnancy: Reexamining the Guidelines. Washington, DC : National Academy Press, 2009.

SFY 2013-2014 AGREEMENT ADDENDA DATA

ATTACHMENT A: WIC QUALITY ASSURANCE DELIVERABLES

Montgomery County

Process Outcome Objectives	2003 to 2005	2006 to 2008	2009 to 2011	% Change from 2003 to 2011	2011 Data		
					Num	Den	Percent
1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2)	71.2%	68.6%	77.3%	8.6%	153	212	72.2%
1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services.	73.9%	79.0%	80.4%	8.8%	229	282	81.2%
1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services.	57.6%	63.0%	65.9%	14.4%	739	1,094	67.6%
2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy.	38.3%	38.4%	36.0%	-6.0%	281	755	37.2%
2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum.	81.6%	81.3%	82.0%	0.5%	243	302	80.5%
3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1)	97.9%	100.0%	97.4%	-0.6%			
3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1)	94.0%	95.2%	96.6%	2.7%			
4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were:							
a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1)	60.4%	61.7%	63.3%	4.7%	243	386	63.0%
b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile)	18.2%	17.0%	17.3%	-5.0%	73	386	18.9%
c. Obese (BMI at or greater than the 95th percentile).	18.2%	19.1%	16.8%	-7.5%	57	386	14.8%
d. Underweight (BMI less than the 5th percentile).	3.2%	2.3%	2.7%	-16.7%	13	386	3.4%
4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3)							
a. Recommended prenatal weight gain	30.2%	35.0%	35.4%	17.3%	83	258	32.2%
b. Excessive prenatal weight gain	46.5%	44.9%	41.0%	-11.8%	106	258	41.1%
c. Inadequate prenatal weight gain	23.3%	20.1%	23.5%	1.2%	69	258	26.7%
5A. Percentage of women who participated in WIC who initiated breastfeeding.	53.3%	57.9%	60.3%	13.2%	230	352	65.3%
5B. Percentage of infants who participated in WIC who were:							
a. Breastfeeding at 6 weeks of age (1)	33.3%	32.8%	33.6%	0.8%	74	302	24.5%
b. Breastfeeding at 6 months of age	15.0%	17.5%	15.6%	4.0%	32	302	10.6%

(1) Item appears also on other WCH Section Agreement Addenda.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI < 18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI ≥ 30.0): Adapted from the National Academy of Sciences. Weight Gain During Pregnancy: Reexamining the Guidelines. Washington, DC : National Academy Press, 2009.

SFY 2013-2014 AGREEMENT ADDENDA DATA

ATTACHMENT A: WIC QUALITY ASSURANCE DELIVERABLES

Moore County

Process Outcome Objectives	2003 to 2005	2006 to 2008	2009 to 2011	% Change from 2003 to 2011	2011 Data		
					Num	Den	Percent
1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2)	70.9%	49.5%	61.8%	-12.8%	292	449	65.0%
1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services.	78.2%	80.4%	80.2%	2.6%	435	555	78.4%
1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services.	64.1%	61.2%	60.4%	-5.9%	1,458	2,309	63.1%
2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy.	31.5%	28.9%	28.4%	-9.7%	334	1,097	30.4%
2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum.	70.2%	70.9%	81.2%	15.6%	397	476	83.4%
3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1)	93.5%	95.5%					
3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1)	89.4%	91.1%					
4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were:							
a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1)	65.3%	66.4%	63.7%	-2.5%	496	752	66.0%
b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile)	16.9%	15.9%	16.7%	-1.3%	111	752	14.8%
c. Obese (BMI at or greater than the 95th percentile).	14.2%	14.6%	16.1%	12.8%	123	752	16.4%
d. Underweight (BMI less than the 5th percentile).	3.5%	3.1%	3.5%	-0.1%	22	752	2.9%
4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3)							
a. Recommended prenatal weight gain	28.3%	27.4%	26.9%	-4.8%	122	446	27.4%
b. Excessive prenatal weight gain	51.2%	53.9%	50.0%	-2.4%	205	446	46.0%
c. Inadequate prenatal weight gain	20.5%	18.7%	23.1%	12.5%	119	446	26.7%
5A. Percentage of women who participated in WIC who initiated breastfeeding.	42.9%	52.9%	55.6%	29.8%	333	586	56.8%
5B. Percentage of infants who participated in WIC who were:							
a. Breastfeeding at 6 weeks of age (1)	24.9%	26.7%	26.5%	6.4%	163	587	27.8%
b. Breastfeeding at 6 months of age	11.2%	11.7%	13.5%	20.5%	83	587	14.1%

(1) Item appears also on other WCH Section Agreement Addenda.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI >=30.0): Adapted from the National Academy of Sciences. *Weight Gain During Pregnancy: Reexamining the Guidelines*. Washington, DC : National Academy Press, 2009.

SFY 2013-2014 AGREEMENT ADDENDA DATA

ATTACHMENT A: WIC QUALITY ASSURANCE DELIVERABLES

Nash County

Process Outcome Objectives	2003 to 2005	2006 to 2008	2009 to 2011	% Change from 2003 to 2011	2011 Data		
					Num	Den	Percent
1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2)	75.6%	76.7%	77.8%	2.8%	537	696	77.2%
1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services.	74.8%	73.6%	73.1%	-2.2%	634	829	76.5%
1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services.	52.3%	48.1%	50.4%	-3.6%	1,867	3,531	52.9%
2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy.	33.2%	31.0%	29.6%	-10.9%	486	1,825	26.6%
2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum.	78.9%	80.3%	80.6%	2.2%	691	846	81.7%
3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1)	91.0%	87.9%	92.9%	2.1%			
3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1)	88.4%	81.9%	83.6%	-5.5%			
4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were:							
a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1)	65.2%	66.0%	62.7%	-3.8%	685	1,076	63.7%
b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile)	15.0%	14.0%	16.0%	6.5%	168	1,076	15.6%
c. Obese (BMI at or greater than the 95th percentile).	13.8%	14.2%	16.5%	19.7%	173	1,076	16.1%
d. Underweight (BMI less than the 5th percentile).	6.1%	5.8%	4.9%	-19.7%	50	1,076	4.6%
4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3)							
a. Recommended prenatal weight gain	30.6%	30.7%	30.1%	-1.6%	189	688	27.5%
b. Excessive prenatal weight gain	45.9%	40.3%	45.3%	-1.3%	328	688	47.7%
c. Inadequate prenatal weight gain	23.5%	29.0%	24.7%	4.7%	171	688	24.9%
5A. Percentage of women who participated in WIC who initiated breastfeeding.	46.7%	43.8%	46.7%	0.0%	441	1,016	43.4%
5B. Percentage of infants who participated in WIC who were:							
a. Breastfeeding at 6 weeks of age (1)	18.6%	22.2%	22.6%	21.7%	227	960	23.6%
b. Breastfeeding at 6 months of age	8.6%	10.1%	11.6%	34.3%	117	960	12.2%

(1) Item appears also on other WCH Section Agreement Addenda.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI >=30.0): Adapted from the National Academy of Sciences. *Weight Gain During Pregnancy: Reexamining the Guidelines*. Washington, DC : National Academy Press, 2009.

SFY 2013-2014 AGREEMENT ADDENDA DATA

ATTACHMENT A: WIC QUALITY ASSURANCE DELIVERABLES

New Hanover County

Process Outcome Objectives	2003 to 2005	2006 to 2008	2009 to 2011	% Change from 2003 to 2011	2011 Data		
					Num	Den	Percent
1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2)	72.1%	72.0%	73.5%	1.9%	687	978	70.2%
1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services.	65.9%	67.2%	66.3%	0.6%	876	1,328	66.0%
1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services.	46.6%	45.9%	51.1%	9.8%	2,867	5,322	53.9%
2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy.	30.0%	34.0%	28.9%	-3.6%	684	2,583	26.5%
2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum.	70.8%	67.7%	69.3%	-2.1%	773	1,085	71.2%
3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1)	66.7%	71.4%	57.1%	-14.3%			
3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1)	49.2%	68.4%	63.5%	29.1%			
4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were:							
a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1)	65.8%	66.1%	65.7%	-0.2%	878	1,330	66.0%
b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile)	16.4%	16.2%	15.7%	-3.9%	202	1,330	15.2%
c. Obese (BMI at or greater than the 95th percentile).	14.9%	15.2%	13.7%	-8.1%	178	1,330	13.4%
d. Underweight (BMI less than the 5th percentile).	2.9%	2.5%	4.9%	68.4%	72	1,330	5.4%
4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3)							
a. Recommended prenatal weight gain	30.0%	31.2%	30.1%	0.4%	300	1,037	28.9%
b. Excessive prenatal weight gain	49.5%	49.0%	49.6%	0.2%	503	1,037	48.5%
c. Inadequate prenatal weight gain	20.6%	19.7%	20.3%	-1.1%	234	1,037	22.6%
5A. Percentage of women who participated in WIC who initiated breastfeeding.	42.3%	50.9%	58.1%	37.4%	832	1,397	59.6%
5B. Percentage of infants who participated in WIC who were:							
a. Breastfeeding at 6 weeks of age (1)	27.3%	32.7%	31.8%	16.5%	458	1,345	34.1%
b. Breastfeeding at 6 months of age	13.9%	16.8%	17.2%	24.1%	255	1,345	19.0%

(1) Item appears also on other WCH Section Agreement Addenda.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI >=30.0): Adapted from the National Academy of Sciences. Weight Gain During Pregnancy: Reexamining the Guidelines. Washington, DC : National Academy Press, 2009.

SFY 2013-2014 AGREEMENT ADDENDA DATA

ATTACHMENT A: WIC QUALITY ASSURANCE DELIVERABLES

Northampton County

Process Outcome Objectives	2003 to 2005	2006 to 2008	2009 to 2011	% Change from 2003 to 2011	2011 Data		
					Num	Den	Percent
1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2)	80.2%	81.4%	80.1%	-0.1%	111	139	79.9%
1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services.	73.9%	71.2%	72.1%	-2.5%	115	170	67.6%
1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services.	63.3%	62.7%	60.6%	-4.2%	551	856	64.4%
2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy.	39.1%	32.7%	32.7%	-16.2%	127	363	35.0%
2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum.	70.9%	75.8%	73.5%	3.7%	125	160	78.1%
3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1)	95.0%	95.2%	100.0%	5.2%			
3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1)	87.1%	89.8%	92.5%	6.1%			
4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were:							
a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1)	61.8%	61.4%	61.8%	0.1%	225	365	61.6%
b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile)	16.4%	16.1%	15.8%	-3.8%	67	365	18.4%
c. Obese (BMI at or greater than the 95th percentile).	16.2%	18.0%	17.3%	6.7%	56	365	15.3%
d. Underweight (BMI less than the 5th percentile).	5.6%	4.4%	5.1%	-9.8%	17	365	4.7%
4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3)							
a. Recommended prenatal weight gain	25.3%	28.1%	28.6%	13.0%	46	149	30.9%
b. Excessive prenatal weight gain	52.2%	49.9%	45.3%	-13.2%	55	149	36.9%
c. Inadequate prenatal weight gain	22.5%	22.0%	26.1%	16.2%	48	149	32.2%
5A. Percentage of women who participated in WIC who initiated breastfeeding.	15.5%	26.8%	33.8%	117.5%	84	201	41.8%
5B. Percentage of infants who participated in WIC who were:							
a. Breastfeeding at 6 weeks of age (1)	11.2%	14.4%	18.4%	64.3%	43	192	22.4%
b. Breastfeeding at 6 months of age	5.9%	5.3%	7.3%	23.3%	17	192	8.9%

(1) Item appears also on other WCH Section Agreement Addenda.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI >=30.0): Adapted from the National Academy of Sciences. Weight Gain During Pregnancy: Reexamining the Guidelines. Washington, DC : National Academy Press, 2009.

SFY 2013-2014 AGREEMENT ADDENDA DATA

ATTACHMENT A: WIC QUALITY ASSURANCE DELIVERABLES

Onslow County

Process Outcome Objectives	2003 to 2005	2006 to 2008	2009 to 2011	% Change from 2003 to 2011	2011 Data		
					Num	Den	Percent
1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2)	74.0%	74.0%	76.3%	3.1%	658	874	75.3%
1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services.	76.9%	75.6%	75.5%	-1.9%	857	1,130	75.8%
1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services.	60.6%	55.1%	53.1%	-12.4%	2,404	4,333	55.5%
2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy.	32.8%	27.1%	26.6%	-18.7%	1,471	5,929	24.8%
2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum.	38.5%	41.7%	35.0%	-9.1%	824	2,271	36.3%
3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1)	85.3%	82.8%	90.9%	6.6%			
3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1)	80.1%	77.1%	82.1%	2.6%			
4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were:							
a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1)	72.2%	70.1%	68.9%	-4.6%	1,793	2,576	69.6%
b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile)	14.3%	15.5%	16.1%	12.5%	392	2,576	15.2%
c. Obese (BMI at or greater than the 95th percentile).	9.6%	11.1%	11.6%	20.8%	290	2,576	11.3%
d. Underweight (BMI less than the 5th percentile).	3.9%	3.3%	3.5%	-11.2%	101	2,576	3.9%
4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3)							
a. Recommended prenatal weight gain	34.2%	32.4%	30.8%	-9.9%	674	2,323	29.0%
b. Excessive prenatal weight gain	48.2%	51.2%	51.0%	5.8%	1,208	2,323	52.0%
c. Inadequate prenatal weight gain	17.6%	16.5%	18.2%	3.3%	441	2,323	19.0%
5A. Percentage of women who participated in WIC who initiated breastfeeding.	51.3%	53.6%	48.7%	-5.1%	1,549	3,075	50.4%
5B. Percentage of infants who participated in WIC who were:							
a. Breastfeeding at 6 weeks of age (1)	31.1%	32.0%	30.6%	-1.5%	1,051	3,487	30.1%
b. Breastfeeding at 6 months of age	12.6%	12.9%	12.0%	-4.9%	445	3,487	12.8%

(1) Item appears also on other WCH Section Agreement Addenda.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI >=30.0): Adapted from the National Academy of Sciences. Weight Gain During Pregnancy: Reexamining the Guidelines. Washington, DC : National Academy Press, 2009.

SFY 2013-2014 AGREEMENT ADDENDA DATA

ATTACHMENT A: WIC QUALITY ASSURANCE DELIVERABLES

Orange County

Process Outcome Objectives	2003 to 2005	2006 to 2008	2009 to 2011	% Change from 2003 to 2011	2011 Data		
					Num	Den	Percent
1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2)	55.6%	67.1%	68.6%	23.3%	244	365	66.8%
1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services.	62.6%	59.3%	58.5%	-6.5%	374	650	57.5%
1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services.	53.3%	52.7%	54.9%	2.9%	1,504	2,606	57.7%
2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy.	28.3%	28.2%	27.3%	-3.6%	639	2,405	26.6%
2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum.	60.5%	61.0%	67.7%	11.9%	787	1,116	70.5%
3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1)	83.7%	83.9%	91.9%	9.8%			
3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1)	32.3%	50.3%	64.1%	98.5%			
4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were:							
a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1)	59.3%	64.0%	65.1%	9.8%	845	1,315	64.3%
b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile)	18.0%	16.9%	15.1%	-16.4%	207	1,315	15.7%
c. Obese (BMI at or greater than the 95th percentile).	19.1%	16.1%	14.8%	-22.6%	221	1,315	16.8%
d. Underweight (BMI less than the 5th percentile).	3.6%	3.1%	5.0%	41.0%	42	1,315	3.2%
4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3)							
a. Recommended prenatal weight gain	31.8%	27.5%	34.0%	7.0%	153	471	32.5%
b. Excessive prenatal weight gain	51.9%	51.3%	44.6%	-14.0%	209	471	44.4%
c. Inadequate prenatal weight gain	16.4%	21.2%	21.4%	30.6%	109	471	23.1%
5A. Percentage of women who participated in WIC who initiated breastfeeding.	61.3%	68.4%	68.8%	12.3%	1,012	1,407	71.9%
5B. Percentage of infants who participated in WIC who were:							
a. Breastfeeding at 6 weeks of age (1)	49.8%	52.4%	52.8%	6.0%	782	1,435	54.5%
b. Breastfeeding at 6 months of age	29.7%	32.6%	32.8%	10.5%	484	1,435	33.7%

(1) Item appears also on other WCH Section Agreement Addenda.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI >=30.0): Adapted from the National Academy of Sciences. Weight Gain During Pregnancy: Reexamining the Guidelines. Washington, DC : National Academy Press, 2009.

SFY 2013-2014 AGREEMENT ADDENDA DATA

ATTACHMENT A: WIC QUALITY ASSURANCE DELIVERABLES

Pamlico County

Process Outcome Objectives	2003 to 2005	2006 to 2008	2009 to 2011	% Change from 2003 to 2011	2011 Data		
					Num	Den	Percent
1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2)	81.5%	85.6%	83.2%	2.1%	44	55	80.0%
1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services.	78.7%	76.5%	75.5%	-4.1%	61	80	76.3%
1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services.	59.5%	59.6%	68.2%	14.5%	285	384	74.2%
2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy.	54.2%	50.1%	47.6%	-12.2%	90	188	47.9%
2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum.	71.3%	73.9%	75.4%	5.8%	61	79	77.2%
3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1)	76.9%	50.0%	100.0%	30.0%			
3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1)	87.0%	80.7%	64.3%	-26.1%			
4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were:							
a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1)	59.7%	65.3%	59.7%	0.0%	99	155	63.9%
b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile)	20.2%	17.0%	20.3%	0.8%	26	155	16.8%
c. Obese (BMI at or greater than the 95th percentile).	17.8%	14.9%	17.3%	-3.1%	26	155	16.8%
d. Underweight (BMI less than the 5th percentile).	2.3%	2.8%	2.7%	16.1%	4	155	2.6%
4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3)							
a. Recommended prenatal weight gain	29.6%	28.4%	31.9%	7.6%	23	76	30.3%
b. Excessive prenatal weight gain	52.8%	50.0%	39.6%	-25.0%	23	76	30.3%
c. Inadequate prenatal weight gain	17.6%	21.6%	28.5%	62.1%	30	76	39.5%
5A. Percentage of women who participated in WIC who initiated breastfeeding.	58.6%	56.0%	49.1%	-16.2%	48	102	47.1%
5B. Percentage of infants who participated in WIC who were:							
a. Breastfeeding at 6 weeks of age (1)	32.8%	45.3%	28.3%	-13.5%	29	98	29.6%
b. Breastfeeding at 6 months of age	16.7%	30.2%	15.0%	-10.6%	17	98	17.3%

(1) Item appears also on other WCH Section Agreement Addenda.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI >=30.0): Adapted from the National Academy of Sciences. Weight Gain During Pregnancy: Reexamining the Guidelines. Washington, DC : National Academy Press, 2009.

SFY 2013-2014 AGREEMENT ADDENDA DATA

ATTACHMENT A: WIC QUALITY ASSURANCE DELIVERABLES

Pender County

Process Outcome Objectives	2003 to 2005	2006 to 2008	2009 to 2011	% Change from 2003 to 2011	2011 Data		
					Num	Den	Percent
1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2)	69.9%	70.8%	65.2%	-6.8%	185	287	64.5%
1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services.	71.5%	76.2%	72.0%	0.6%	278	387	71.8%
1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services.	57.6%	57.8%	57.8%	0.5%	1,029	1,750	58.8%
2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy.	34.7%	32.0%	29.3%	-15.5%	219	698	31.4%
2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum.	73.8%	78.3%	76.4%	3.4%	230	300	76.7%
3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1)	80.5%	82.5%	83.4%	3.7%			
3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1)	78.7%	78.4%	80.6%	2.5%			
4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were:							
a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1)	66.1%	60.1%	55.3%	-16.4%	275	488	56.4%
b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile)	13.0%	17.5%	19.6%	50.5%	90	488	18.4%
c. Obese (BMI at or greater than the 95th percentile).	13.8%	19.7%	22.5%	63.2%	104	488	21.3%
d. Underweight (BMI less than the 5th percentile).	7.1%	2.7%	2.6%	-63.0%	19	488	3.9%
4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3)							
a. Recommended prenatal weight gain	31.4%	31.1%	29.7%	-5.4%	97	320	30.3%
b. Excessive prenatal weight gain	45.4%	48.3%	44.9%	-1.1%	137	320	42.8%
c. Inadequate prenatal weight gain	23.2%	20.6%	25.3%	9.3%	86	320	26.9%
5A. Percentage of women who participated in WIC who initiated breastfeeding.	34.9%	46.3%	50.2%	43.8%	243	420	57.9%
5B. Percentage of infants who participated in WIC who were:							
a. Breastfeeding at 6 weeks of age (1)	16.5%	27.2%	26.5%	60.1%	108	435	24.8%
b. Breastfeeding at 6 months of age	5.8%	10.6%	13.3%	127.9%	66	435	15.2%

(1) Item appears also on other WCH Section Agreement Addenda.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI >=30.0): Adapted from the National Academy of Sciences. Weight Gain During Pregnancy: Reexamining the Guidelines. Washington, DC : National Academy Press, 2009.

SFY 2013-2014 AGREEMENT ADDENDA DATA

ATTACHMENT A: WIC QUALITY ASSURANCE DELIVERABLES

Person County

Process Outcome Objectives	2003 to 2005	2006 to 2008	2009 to 2011	% Change from 2003 to 2011	2011 Data		
					Num	Den	Percent
1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2)	79.9%	83.2%	84.2%	5.5%	213	260	81.9%
1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services.	74.4%	82.1%	81.3%	9.2%	255	306	83.3%
1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services.	64.8%	62.4%	68.0%	4.9%	900	1,304	69.0%
2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy.	38.9%	37.3%	34.8%	-10.6%	173	543	31.9%
2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum.	77.1%	71.5%	78.1%	1.3%	216	256	84.4%
3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1)	91.4%	89.2%	50.0%	-45.3%			
3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1)	91.2%	90.2%	97.2%	6.6%			
4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were:							
a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1)	68.3%	63.8%	63.4%	-7.1%	312	473	66.0%
b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile)	14.6%	16.9%	15.9%	9.3%	62	473	13.1%
c. Obese (BMI at or greater than the 95th percentile).	14.5%	15.8%	17.1%	18.1%	86	473	18.2%
d. Underweight (BMI less than the 5th percentile).	2.7%	3.6%	3.5%	33.6%	13	473	2.7%
4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3)							
a. Recommended prenatal weight gain	32.9%	33.6%	28.7%	-12.9%	64	257	24.9%
b. Excessive prenatal weight gain	44.7%	47.7%	50.3%	12.5%	129	257	50.2%
c. Inadequate prenatal weight gain	22.4%	18.8%	21.0%	-6.1%	64	257	24.9%
5A. Percentage of women who participated in WIC who initiated breastfeeding.	35.6%	37.8%	43.9%	23.3%	124	305	40.7%
5B. Percentage of infants who participated in WIC who were:							
a. Breastfeeding at 6 weeks of age (1)	20.4%	22.4%	25.0%	22.4%	92	312	29.5%
b. Breastfeeding at 6 months of age	9.5%	9.8%	14.3%	49.9%	60	312	19.2%

(1) Item appears also on other WCH Section Agreement Addenda.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI >=30.0): Adapted from the National Academy of Sciences. Weight Gain During Pregnancy: Reexamining the Guidelines. Washington, DC : National Academy Press, 2009.

SFY 2013-2014 AGREEMENT ADDENDA DATA

ATTACHMENT A: WIC QUALITY ASSURANCE DELIVERABLES

Pitt County

Process Outcome Objectives	2003 to 2005	2006 to 2008	2009 to 2011	% Change from 2003 to 2011	2011 Data		
					Num	Den	Percent
1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2)	74.9%	76.5%	75.2%	0.5%	827	1,089	75.9%
1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services.	75.1%	76.3%	78.6%	4.7%	1,085	1,380	78.6%
1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services.	60.2%	57.2%	60.2%	0.1%	3,561	5,678	62.7%
2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy.	32.0%	30.6%	29.6%	-7.5%	892	3,040	29.3%
2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum.	81.5%	82.2%	80.0%	-1.9%	1,011	1,262	80.1%
3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1)	89.5%	100.0%					
3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1)	87.6%	83.5%					
4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were:							
a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1)	59.7%	64.7%	61.9%	3.8%	1,212	1,988	61.0%
b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile)	18.6%	17.0%	17.4%	-6.4%	365	1,988	18.4%
c. Obese (BMI at or greater than the 95th percentile).	19.3%	15.6%	17.7%	-8.1%	346	1,988	17.4%
d. Underweight (BMI less than the 5th percentile).	2.5%	2.7%	3.0%	20.4%	65	1,988	3.3%
4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3)							
a. Recommended prenatal weight gain	31.4%	32.3%	31.8%	1.3%	328	1,127	29.1%
b. Excessive prenatal weight gain	45.5%	46.5%	47.4%	4.2%	533	1,127	47.3%
c. Inadequate prenatal weight gain	23.2%	21.3%	20.8%	-10.0%	266	1,127	23.6%
5A. Percentage of women who participated in WIC who initiated breastfeeding.	44.9%	48.6%	47.0%	4.8%	687	1,549	44.4%
5B. Percentage of infants who participated in WIC who were:							
a. Breastfeeding at 6 weeks of age (1)	23.7%	27.7%	30.6%	29.1%	462	1,429	32.3%
b. Breastfeeding at 6 months of age	10.9%	13.1%	15.7%	44.5%	244	1,429	17.1%

(1) Item appears also on other WCH Section Agreement Addenda.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI >=30.0): Adapted from the National Academy of Sciences. Weight Gain During Pregnancy: Reexamining the Guidelines. Washington, DC : National Academy Press, 2009.

SFY 2013-2014 AGREEMENT ADDENDA DATA

ATTACHMENT A: WIC QUALITY ASSURANCE DELIVERABLES

Randolph County

Process Outcome Objectives	2003 to 2005	2006 to 2008	2009 to 2011	% Change from 2003 to 2011	2011 Data		
					Num	Den	Percent
1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2)	67.5%	66.9%	72.5%	7.5%	622	873	71.2%
1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services.	73.9%	75.6%	76.0%	2.8%	930	1,209	76.9%
1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services.	53.6%	52.6%	56.5%	5.3%	3,213	5,433	59.1%
2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy.	30.8%	28.6%	30.2%	-2.0%	566	1,784	31.7%
2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum.	61.2%	54.1%	75.7%	23.8%	785	937	83.8%
3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1)	89.2%	85.7%	100.0%	12.1%			
3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1)	81.7%	74.5%	80.4%	-1.6%			
4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were:							
a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1)	56.7%	62.9%	64.3%	13.3%	1,134	1,794	63.2%
b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile)	18.5%	17.1%	15.9%	-13.9%	295	1,794	16.4%
c. Obese (BMI at or greater than the 95th percentile).	22.4%	17.3%	16.1%	-27.9%	299	1,794	16.7%
d. Underweight (BMI less than the 5th percentile).	2.4%	2.8%	3.7%	53.0%	66	1,794	3.7%
4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3)							
a. Recommended prenatal weight gain	29.9%	32.5%	32.2%	7.7%	282	947	29.8%
b. Excessive prenatal weight gain	53.2%	49.9%	48.8%	-8.1%	472	947	49.8%
c. Inadequate prenatal weight gain	17.0%	17.5%	19.0%	11.9%	193	947	20.4%
5A. Percentage of women who participated in WIC who initiated breastfeeding.	61.5%	58.9%	60.0%	-2.5%	746	1,172	63.7%
5B. Percentage of infants who participated in WIC who were:							
a. Breastfeeding at 6 weeks of age (1)	32.8%	37.2%	34.9%	6.4%	445	1,188	37.5%
b. Breastfeeding at 6 months of age	16.2%	19.0%	16.7%	3.4%	217	1,188	18.3%

(1) Item appears also on other WCH Section Agreement Addenda.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI >=30.0): Adapted from the National Academy of Sciences. Weight Gain During Pregnancy: Reexamining the Guidelines. Washington, DC : National Academy Press, 2009.

SFY 2013-2014 AGREEMENT ADDENDA DATA

ATTACHMENT A: WIC QUALITY ASSURANCE DELIVERABLES

Richmond County

Process Outcome Objectives	2003 to 2005	2006 to 2008	2009 to 2011	% Change from 2003 to 2011	2011 Data		
					Num	Den	Percent
1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2)	79.7%	78.2%	77.3%	-3.0%	338	465	72.7%
1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services.	83.4%	82.1%	81.0%	-2.9%	439	541	81.1%
1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services.	72.1%	67.2%	61.3%	-15.0%	1,364	2,245	60.8%
2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy.	34.3%	31.6%	32.5%	-5.4%	356	1,134	31.4%
2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum.	75.9%	71.5%	66.0%	-13.1%	285	440	64.8%
3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1)							
3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1)							
4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were:							
a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1)	68.9%	67.6%	68.1%	-1.2%	527	808	65.2%
b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile)	8.4%	10.9%	10.6%	26.8%	110	808	13.6%
c. Obese (BMI at or greater than the 95th percentile).	8.1%	10.9%	11.3%	39.5%	111	808	13.7%
d. Underweight (BMI less than the 5th percentile).	14.6%	10.7%	10.0%	-31.6%	60	808	7.4%
4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3)							
a. Recommended prenatal weight gain	30.5%	27.7%	27.6%	-9.6%	114	449	25.4%
b. Excessive prenatal weight gain	49.0%	49.1%	49.0%	0.1%	221	449	49.2%
c. Inadequate prenatal weight gain	20.5%	23.2%	23.4%	14.1%	114	449	25.4%
5A. Percentage of women who participated in WIC who initiated breastfeeding.	41.7%	29.2%	28.7%	-31.0%	169	552	30.6%
5B. Percentage of infants who participated in WIC who were:							
a. Breastfeeding at 6 weeks of age (1)	21.4%	19.0%	14.9%	-30.1%	97	588	16.5%
b. Breastfeeding at 6 months of age	10.1%	8.9%	6.6%	-35.0%	46	588	7.8%

(1) Item appears also on other WCH Section Agreement Addenda.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI >=30.0): Adapted from the National Academy of Sciences. Weight Gain During Pregnancy: Reexamining the Guidelines. Washington, DC : National Academy Press, 2009.

SFY 2013-2014 AGREEMENT ADDENDA DATA

ATTACHMENT A: WIC QUALITY ASSURANCE DELIVERABLES

Robeson County

Process Outcome Objectives	2003 to 2005	2006 to 2008	2009 to 2011	% Change from 2003 to 2011	2011 Data		
					Num	Den	Percent
1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2)	72.8%	74.6%	72.7%	-0.2%	984	1,420	69.3%
1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services.	76.3%	75.7%	77.1%	1.1%	1,364	1,737	78.5%
1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services.	52.7%	47.9%	50.1%	-4.8%	4,002	7,649	52.3%
2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy.	31.3%	20.8%	22.1%	-29.5%	556	2,334	23.8%
2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum.	81.3%	83.0%	81.1%	-0.2%	966	1,192	81.0%
3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1)	89.8%	88.5%	85.6%	-4.7%			
3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1)	89.1%	84.4%	88.1%	-1.1%			
4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were:							
a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1)	66.4%	65.1%	58.2%	-12.4%	1,058	1,791	59.1%
b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile)	13.8%	15.1%	18.6%	34.7%	302	1,791	16.9%
c. Obese (BMI at or greater than the 95th percentile).	13.0%	15.4%	20.3%	56.0%	376	1,791	21.0%
d. Underweight (BMI less than the 5th percentile).	6.8%	4.5%	3.0%	-56.5%	55	1,791	3.1%
4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3)							
a. Recommended prenatal weight gain	31.7%	30.1%	28.7%	-9.5%	372	1,352	27.5%
b. Excessive prenatal weight gain	41.5%	42.5%	45.0%	8.6%	592	1,352	43.8%
c. Inadequate prenatal weight gain	26.8%	27.4%	26.2%	-2.0%	388	1,352	28.7%
5A. Percentage of women who participated in WIC who initiated breastfeeding.	35.6%	36.5%	34.5%	-2.9%	627	1,586	39.5%
5B. Percentage of infants who participated in WIC who were:							
a. Breastfeeding at 6 weeks of age (1)	19.2%	20.1%	19.8%	3.3%	364	1,698	21.4%
b. Breastfeeding at 6 months of age	9.6%	10.5%	9.6%	0.9%	165	1,698	9.7%

(1) Item appears also on other WCH Section Agreement Addenda.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI >=30.0): Adapted from the National Academy of Sciences. Weight Gain During Pregnancy: Reexamining the Guidelines. Washington, DC : National Academy Press, 2009.

SFY 2013-2014 AGREEMENT ADDENDA DATA

ATTACHMENT A: WIC QUALITY ASSURANCE DELIVERABLES

Rockingham County

Process Outcome Objectives	2003 to 2005	2006 to 2008	2009 to 2011	% Change from 2003 to 2011	2011 Data		
					Num	Den	Percent
1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2)	77.6%	77.7%	78.4%	1.0%	400	520	76.9%
1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services.	76.6%	75.9%	77.0%	0.5%	514	677	75.9%
1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services.	55.7%	54.1%	58.2%	4.5%	1,837	3,032	60.6%
2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy.	33.3%	36.2%	33.3%	-0.1%	527	1,577	33.4%
2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum.	76.3%	73.9%	76.4%	0.1%	484	624	77.6%
3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1)	79.8%	89.7%	92.3%	15.7%			
3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1)	52.8%	59.3%	69.2%	31.0%			
4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were:							
a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1)	70.6%	65.3%	64.7%	-8.4%	714	1,075	66.4%
b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile)	11.0%	16.4%	15.6%	41.9%	164	1,075	15.3%
c. Obese (BMI at or greater than the 95th percentile).	11.5%	16.0%	16.3%	42.0%	162	1,075	15.1%
d. Underweight (BMI less than the 5th percentile).	7.0%	2.2%	3.5%	-50.5%	35	1,075	3.3%
4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3)							
a. Recommended prenatal weight gain	30.6%	31.9%	29.4%	-4.1%	154	549	28.1%
b. Excessive prenatal weight gain	46.6%	50.3%	48.9%	5.0%	272	549	49.5%
c. Inadequate prenatal weight gain	22.8%	17.8%	21.7%	-4.7%	123	549	22.4%
5A. Percentage of women who participated in WIC who initiated breastfeeding.	40.0%	43.1%	45.7%	14.2%	389	760	51.2%
5B. Percentage of infants who participated in WIC who were:							
a. Breastfeeding at 6 weeks of age (1)	24.7%	26.8%	25.2%	2.2%	176	717	24.5%
b. Breastfeeding at 6 months of age	12.8%	13.2%	11.3%	-11.6%	84	717	11.7%

(1) Item appears also on other WCH Section Agreement Addenda.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI >=30.0): Adapted from the National Academy of Sciences. Weight Gain During Pregnancy: Reexamining the Guidelines. Washington, DC : National Academy Press, 2009.

SFY 2013-2014 AGREEMENT ADDENDA DATA

ATTACHMENT A: WIC QUALITY ASSURANCE DELIVERABLES

Rowan County

Process Outcome Objectives	2003 to 2005	2006 to 2008	2009 to 2011	% Change from 2003 to 2011	2011 Data		
					Num	Den	Percent
1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2)	74.6%	70.8%	67.9%	-8.9%	578	862	67.1%
1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services.	72.8%	72.1%	74.4%	2.2%	850	1,142	74.4%
1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services.	56.8%	54.3%	55.2%	-2.8%	3,013	5,322	56.6%
2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy.	34.5%	20.6%	26.7%	-22.5%	422	1,386	30.4%
2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum.	75.8%	80.4%	78.4%	3.4%	625	796	78.5%
3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1)	70.8%	63.2%	75.0%	5.9%			
3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1)	75.0%	68.9%	78.6%	4.8%			
4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were:							
a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1)	64.1%	65.4%	67.1%	4.8%	1,053	1,591	66.2%
b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile)	17.6%	16.1%	14.7%	-17.0%	250	1,591	15.7%
c. Obese (BMI at or greater than the 95th percentile).	16.0%	14.7%	13.6%	-14.8%	217	1,591	13.6%
d. Underweight (BMI less than the 5th percentile).	2.3%	3.8%	4.6%	99.1%	71	1,591	4.5%
4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3)							
a. Recommended prenatal weight gain	31.4%	31.7%	30.6%	-2.8%	248	880	28.2%
b. Excessive prenatal weight gain	46.2%	45.6%	47.8%	3.5%	447	880	50.8%
c. Inadequate prenatal weight gain	22.4%	22.7%	21.7%	-3.2%	185	880	21.0%
5A. Percentage of women who participated in WIC who initiated breastfeeding.	47.8%	50.7%	52.2%	9.1%	561	1,033	54.3%
5B. Percentage of infants who participated in WIC who were:							
a. Breastfeeding at 6 weeks of age (1)	29.9%	29.9%	27.9%	-6.5%	338	1,121	30.2%
b. Breastfeeding at 6 months of age	13.1%	12.2%	11.2%	-14.6%	126	1,121	11.2%

(1) Item appears also on other WCH Section Agreement Addenda.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI >=30.0): Adapted from the National Academy of Sciences. Weight Gain During Pregnancy: Reexamining the Guidelines. Washington, DC : National Academy Press, 2009.

SFY 2013-2014 AGREEMENT ADDENDA DATA

ATTACHMENT A: WIC QUALITY ASSURANCE DELIVERABLES

Rutherford-Polk-McDowell District

Process Outcome Objectives	2003 to 2005	2006 to 2008	2009 to 2011	% Change from 2003 to 2011	2011 Data		
					Num	Den	Percent
1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2)	66.8%	66.7%	76.6%	14.6%	614	848	72.4%
1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services.	74.9%	76.7%	74.9%	0.0%	807	1,041	77.5%
1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services.	61.9%	60.3%	60.9%	-1.5%	2,765	4,396	62.9%
2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy.	32.5%	35.2%	33.4%	2.8%	885	2,568	34.5%
2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum.	69.9%	72.4%	76.2%	9.0%	780	981	79.5%
3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1)	91.4%	95.0%	96.8%	5.9%			
3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1)	91.4%	90.7%	91.3%	-0.1%			
4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were:							
a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1)	67.9%	67.3%	66.3%	-2.4%	1,069	1,632	65.5%
b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile)	14.9%	15.6%	15.1%	1.4%	250	1,632	15.3%
c. Obese (BMI at or greater than the 95th percentile).	12.9%	13.4%	13.3%	2.9%	236	1,632	14.5%
d. Underweight (BMI less than the 5th percentile).	4.2%	3.7%	5.3%	24.7%	77	1,632	4.7%
4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3)							
a. Recommended prenatal weight gain	33.4%	29.5%	28.9%	-13.4%	224	846	26.5%
b. Excessive prenatal weight gain	37.4%	40.5%	48.3%	29.0%	433	846	51.2%
c. Inadequate prenatal weight gain	29.2%	30.0%	22.8%	-21.9%	189	846	22.3%
5A. Percentage of women who participated in WIC who initiated breastfeeding.	46.4%	54.4%	60.3%	30.0%	701	1,153	60.8%
5B. Percentage of infants who participated in WIC who were:							
a. Breastfeeding at 6 weeks of age (1)	28.1%	31.1%	30.0%	6.6%	338	1,102	30.7%
b. Breastfeeding at 6 months of age	14.7%	14.5%	16.2%	10.0%	190	1,102	17.2%

(1) Item appears also on other WCH Section Agreement Addenda.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI >=30.0): Adapted from the National Academy of Sciences. Weight Gain During Pregnancy: Reexamining the Guidelines. Washington, DC : National Academy Press, 2009.

SFY 2013-2014 AGREEMENT ADDENDA DATA

ATTACHMENT A: WIC QUALITY ASSURANCE DELIVERABLES

Sampson County

Process Outcome Objectives	2003 to 2005	2006 to 2008	2009 to 2011	% Change from 2003 to 2011	2011 Data		
					Num	Den	Percent
1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2)	76.3%	77.7%	75.6%	-0.9%	356	496	71.8%
1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services.	74.9%	77.7%	80.0%	6.8%	594	753	78.9%
1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services.	56.0%	55.8%	59.2%	5.6%	1,982	3,226	61.4%
2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy.	41.5%	34.8%	27.1%	-34.8%	334	1,261	26.5%
2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum.	71.9%	77.8%	79.9%	11.2%	420	514	81.7%
3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1)	92.3%	72.2%	100.0%	8.3%			
3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1)	80.0%	84.0%	91.3%	14.1%			
4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were:							
a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1)	61.1%	54.3%	56.6%	-7.3%	712	1,340	53.1%
b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile)	15.2%	19.4%	19.6%	29.2%	275	1,340	20.5%
c. Obese (BMI at or greater than the 95th percentile).	17.6%	23.8%	20.9%	18.6%	311	1,340	23.2%
d. Underweight (BMI less than the 5th percentile).	6.2%	2.5%	2.9%	-52.9%	42	1,340	3.1%
4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3)							
a. Recommended prenatal weight gain	31.9%	31.0%	29.6%	-7.3%	168	575	29.2%
b. Excessive prenatal weight gain	48.3%	49.5%	45.2%	-6.4%	238	575	41.4%
c. Inadequate prenatal weight gain	19.8%	19.4%	25.2%	27.2%	169	575	29.4%
5A. Percentage of women who participated in WIC who initiated breastfeeding.	26.3%	30.4%	24.7%	-6.0%	213	682	31.2%
5B. Percentage of infants who participated in WIC who were:							
a. Breastfeeding at 6 weeks of age (1)	16.1%	19.5%	16.9%	4.9%	116	637	18.2%
b. Breastfeeding at 6 months of age	6.8%	12.2%	10.2%	50.4%	82	637	12.9%

(1) Item appears also on other WCH Section Agreement Addenda.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI >=30.0): Adapted from the National Academy of Sciences. Weight Gain During Pregnancy: Reexamining the Guidelines. Washington, DC : National Academy Press, 2009.

SFY 2013-2014 AGREEMENT ADDENDA DATA

ATTACHMENT A: WIC QUALITY ASSURANCE DELIVERABLES

Scotland County

Process Outcome Objectives	2003 to 2005	2006 to 2008	2009 to 2011	% Change from 2003 to 2011	2011 Data		
					Num	Den	Percent
1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2)	86.0%	84.5%	83.8%	-2.6%	311	385	80.8%
1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services.	80.9%	81.4%	80.9%	0.1%	349	436	80.0%
1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services.	65.7%	61.1%	63.4%	-3.5%	1,205	1,882	64.0%
2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy.	55.0%	50.9%	43.8%	-20.2%	377	940	40.1%
2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum.	80.2%	79.6%	80.3%	0.1%	344	433	79.4%
3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1)	66.7%						
3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1)	76.9%						
4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were:							
a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1)	67.0%	68.1%	66.2%	-1.1%	517	773	66.9%
b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile)	14.3%	14.8%	14.6%	1.8%	110	773	14.2%
c. Obese (BMI at or greater than the 95th percentile).	13.5%	12.6%	13.8%	1.9%	111	773	14.4%
d. Underweight (BMI less than the 5th percentile).	5.2%	4.5%	5.4%	4.4%	35	773	4.5%
4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3)							
a. Recommended prenatal weight gain	29.1%	27.4%	24.8%	-14.8%	90	393	22.9%
b. Excessive prenatal weight gain	46.9%	46.6%	46.0%	-2.0%	170	393	43.3%
c. Inadequate prenatal weight gain	24.0%	26.1%	29.2%	21.9%	133	393	33.8%
5A. Percentage of women who participated in WIC who initiated breastfeeding.	35.0%	25.8%	28.5%	-18.4%	172	519	33.1%
5B. Percentage of infants who participated in WIC who were:							
a. Breastfeeding at 6 weeks of age (1)	11.2%	9.4%	12.2%	8.7%	75	512	14.6%
b. Breastfeeding at 6 months of age	3.8%	2.4%	5.6%	47.5%	40	512	7.8%

(1) Item appears also on other WCH Section Agreement Addenda.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI >=30.0): Adapted from the National Academy of Sciences. Weight Gain During Pregnancy: Reexamining the Guidelines. Washington, DC : National Academy Press, 2009.

SFY 2013-2014 AGREEMENT ADDENDA DATA

ATTACHMENT A: WIC QUALITY ASSURANCE DELIVERABLES

Stanly County

Process Outcome Objectives	2003 to 2005	2006 to 2008	2009 to 2011	% Change from 2003 to 2011	2011 Data		
					Num	Den	Percent
1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2)	74.8%	79.4%	78.9%	5.4%	282	375	75.2%
1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services.	79.2%	73.8%	76.8%	-3.0%	339	447	75.8%
1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services.	59.7%	54.2%	55.3%	-7.5%	1,120	1,937	57.8%
2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy.	38.2%	40.2%	34.1%	-10.8%	295	844	35.0%
2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum.	69.4%	45.5%	64.4%	-7.2%	315	388	81.2%
3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1)	100.0%	88.9%	100.0%	0.0%			
3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1)	82.9%	85.6%	83.8%	1.1%			
4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were:							
a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1)	69.7%	70.2%	63.5%	-8.9%	396	604	65.6%
b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile)	12.7%	12.4%	15.8%	24.2%	83	604	13.7%
c. Obese (BMI at or greater than the 95th percentile).	12.8%	11.2%	16.3%	27.0%	93	604	15.4%
d. Underweight (BMI less than the 5th percentile).	4.7%	6.2%	4.4%	-7.6%	32	604	5.3%
4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3)							
a. Recommended prenatal weight gain	33.0%	28.6%	29.0%	-12.3%	106	378	28.0%
b. Excessive prenatal weight gain	40.1%	44.1%	47.7%	18.8%	184	378	48.7%
c. Inadequate prenatal weight gain	26.9%	27.3%	23.4%	-13.0%	88	378	23.3%
5A. Percentage of women who participated in WIC who initiated breastfeeding.	65.7%	64.2%	66.9%	1.8%	328	482	68.0%
5B. Percentage of infants who participated in WIC who were:							
a. Breastfeeding at 6 weeks of age (1)	15.6%	18.2%	24.1%	54.7%	137	497	27.6%
b. Breastfeeding at 6 months of age	5.8%	9.5%	11.7%	100.8%	68	497	13.7%

(1) Item appears also on other WCH Section Agreement Addenda.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI >=30.0): Adapted from the National Academy of Sciences. Weight Gain During Pregnancy: Reexamining the Guidelines. Washington, DC : National Academy Press, 2009.

SFY 2013-2014 AGREEMENT ADDENDA DATA

ATTACHMENT A: WIC QUALITY ASSURANCE DELIVERABLES

Stokes County

Process Outcome Objectives	2003 to 2005	2006 to 2008	2009 to 2011	% Change from 2003 to 2011	2011 Data		
					Num	Den	Percent
1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2)	75.2%	75.9%	72.3%	-3.9%	157	227	69.2%
1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services.	75.9%	76.1%	73.7%	-2.9%	211	274	77.0%
1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services.	56.5%	55.9%	57.7%	2.2%	717	1,226	58.5%
2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy.	34.1%	39.8%	33.6%	-1.4%	178	520	34.2%
2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum.	78.2%	79.3%	75.9%	-2.9%	170	226	75.2%
3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1)	93.9%	88.8%	82.4%	-12.3%			
3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1)	87.7%	88.5%	91.4%	4.2%			
4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were:							
a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1)	66.6%	62.3%	66.7%	0.1%	277	396	69.9%
b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile)	15.8%	18.3%	16.3%	3.4%	60	396	15.2%
c. Obese (BMI at or greater than the 95th percentile).	13.4%	16.3%	14.1%	4.8%	48	396	12.1%
d. Underweight (BMI less than the 5th percentile).	4.2%	3.1%	2.9%	-29.7%	11	396	2.8%
4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3)							
a. Recommended prenatal weight gain	32.2%	28.6%	28.0%	-13.1%	67	240	27.9%
b. Excessive prenatal weight gain	52.1%	55.2%	47.4%	-9.0%	109	240	45.4%
c. Inadequate prenatal weight gain	15.7%	16.1%	24.6%	56.5%	64	240	26.7%
5A. Percentage of women who participated in WIC who initiated breastfeeding.	50.8%	50.1%	50.2%	-1.2%	167	294	56.8%
5B. Percentage of infants who participated in WIC who were:							
a. Breastfeeding at 6 weeks of age (1)	21.1%	24.2%	23.1%	9.7%	84	335	25.1%
b. Breastfeeding at 6 months of age	9.4%	10.8%	10.3%	8.7%	37	335	11.0%

(1) Item appears also on other WCH Section Agreement Addenda.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI >=30.0): Adapted from the National Academy of Sciences. Weight Gain During Pregnancy: Reexamining the Guidelines. Washington, DC : National Academy Press, 2009.

SFY 2013-2014 AGREEMENT ADDENDA DATA

ATTACHMENT A: WIC QUALITY ASSURANCE DELIVERABLES

Surry County

Process Outcome Objectives	2003 to 2005	2006 to 2008	2009 to 2011	% Change from 2003 to 2011	2011 Data		
					Num	Den	Percent
1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2)	72.5%	68.7%	68.1%	-6.0%	319	461	69.2%
1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services.	76.7%	75.9%	74.7%	-2.6%	460	614	74.9%
1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services.	55.4%	51.5%	56.6%	2.2%	1,567	2,682	58.4%
2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy.	46.6%	45.3%	41.1%	-11.8%	387	915	42.3%
2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum.	79.5%	80.5%	80.9%	1.8%	400	502	79.7%
3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1)	83.5%	82.4%	78.8%	-5.5%			
3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1)	89.0%	79.7%	75.2%	-15.5%			
4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were:							
a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1)	68.2%	65.3%	61.1%	-10.5%	298	481	62.0%
b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile)	13.2%	16.4%	16.9%	27.7%	82	481	17.0%
c. Obese (BMI at or greater than the 95th percentile).	12.2%	15.0%	19.1%	56.6%	87	481	18.1%
d. Underweight (BMI less than the 5th percentile).	6.4%	3.4%	3.0%	-53.3%	14	481	2.9%
4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3)							
a. Recommended prenatal weight gain	28.1%	29.5%	29.8%	6.0%	134	466	28.8%
b. Excessive prenatal weight gain	52.5%	53.9%	48.9%	-6.8%	236	466	50.6%
c. Inadequate prenatal weight gain	19.4%	16.6%	21.3%	9.8%	96	466	20.6%
5A. Percentage of women who participated in WIC who initiated breastfeeding.	53.0%	42.3%	53.0%	-0.1%	345	655	52.7%
5B. Percentage of infants who participated in WIC who were:							
a. Breastfeeding at 6 weeks of age (1)	33.3%	28.9%	31.3%	-6.0%	191	614	31.1%
b. Breastfeeding at 6 months of age	17.1%	13.0%	16.2%	-5.3%	96	614	15.6%

(1) Item appears also on other WCH Section Agreement Addenda.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI >=30.0): Adapted from the National Academy of Sciences. Weight Gain During Pregnancy: Reexamining the Guidelines. Washington, DC : National Academy Press, 2009.

SFY 2013-2014 AGREEMENT ADDENDA DATA

ATTACHMENT A: WIC QUALITY ASSURANCE DELIVERABLES

Swain County

Process Outcome Objectives	2003 to 2005	2006 to 2008	2009 to 2011	% Change from 2003 to 2011	2011 Data		
					Num	Den	Percent
1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2)	84.8%	85.8%	79.3%	-6.4%	117	154	76.0%
1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services.	76.2%	73.3%	67.1%	-12.0%	120	173	69.4%
1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services.	78.6%	74.2%	74.0%	-5.9%	539	716	75.3%
2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy.	50.3%	49.2%	36.5%	-27.5%	66	184	35.9%
2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum.	79.2%	75.1%	59.9%	-24.3%	37	91	40.7%
3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1)	100.0%						
3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1)	88.9%						
4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were:							
a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1)	72.7%	67.7%	66.4%	-8.7%	116	178	65.2%
b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile)	14.3%	16.9%	16.3%	13.5%	25	178	14.0%
c. Obese (BMI at or greater than the 95th percentile).	9.4%	13.7%	13.2%	39.5%	28	178	15.7%
d. Underweight (BMI less than the 5th percentile).	3.5%	1.7%	4.2%	18.1%	9	178	5.1%
4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3)							
a. Recommended prenatal weight gain	31.0%	27.3%	29.1%	-6.1%	32	141	22.7%
b. Excessive prenatal weight gain	48.5%	56.4%	51.5%	6.2%	69	141	48.9%
c. Inadequate prenatal weight gain	20.5%	16.3%	19.3%	-5.5%	40	141	28.4%
5A. Percentage of women who participated in WIC who initiated breastfeeding.	82.6%	66.2%	76.3%	-7.6%	62	111	55.9%
5B. Percentage of infants who participated in WIC who were:							
a. Breastfeeding at 6 weeks of age (1)	37.1%	29.6%	46.2%	24.4%	51	95	53.7%
b. Breastfeeding at 6 months of age	20.8%	17.0%	25.8%	24.5%	24	95	25.3%

(1) Item appears also on other WCH Section Agreement Addenda.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI >=30.0): Adapted from the National Academy of Sciences. Weight Gain During Pregnancy: Reexamining the Guidelines. Washington, DC : National Academy Press, 2009.

SFY 2013-2014 AGREEMENT ADDENDA DATA

ATTACHMENT A: WIC QUALITY ASSURANCE DELIVERABLES

Toe River District

Process Outcome Objectives	2003 to 2005	2006 to 2008	2009 to 2011	% Change from 2003 to 2011	2011 Data		
					Num	Den	Percent
1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2)	72.8%	72.6%	71.8%	-1.5%	178	275	64.7%
1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services.	63.4%	63.6%	67.8%	7.0%	239	352	67.9%
1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services.	66.8%	59.9%	64.6%	-3.4%	985	1,463	67.3%
2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy.	41.5%	43.6%	39.6%	-4.5%	316	841	37.6%
2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum.	70.7%	68.2%	67.2%	-4.9%	249	372	66.9%
3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1)	89.2%	94.4%	100.0%	12.0%			
3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1)	92.1%	87.9%	87.5%	-4.9%			
4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were:							
a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1)	65.6%	68.1%	64.3%	-2.0%	364	561	64.9%
b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile)	16.4%	15.5%	17.1%	4.7%	101	561	18.0%
c. Obese (BMI at or greater than the 95th percentile).	15.7%	12.7%	14.8%	-5.5%	78	561	13.9%
d. Underweight (BMI less than the 5th percentile).	2.4%	3.6%	3.8%	59.7%	18	561	3.2%
4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3)							
a. Recommended prenatal weight gain	33.5%	32.7%	31.0%	-7.4%	90	290	31.0%
b. Excessive prenatal weight gain	43.0%	46.1%	44.1%	2.5%	120	290	41.4%
c. Inadequate prenatal weight gain	23.5%	21.1%	24.9%	5.9%	80	290	27.6%
5A. Percentage of women who participated in WIC who initiated breastfeeding.	63.9%	62.6%	64.2%	0.4%	272	444	61.3%
5B. Percentage of infants who participated in WIC who were:							
a. Breastfeeding at 6 weeks of age (1)	40.6%	36.3%	41.6%	2.5%	164	423	38.8%
b. Breastfeeding at 6 months of age	26.2%	22.8%	27.6%	5.5%	115	423	27.2%

(1) Item appears also on other WCH Section Agreement Addenda.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI >=30.0): Adapted from the National Academy of Sciences. Weight Gain During Pregnancy: Reexamining the Guidelines. Washington, DC : National Academy Press, 2009.

SFY 2013-2014 AGREEMENT ADDENDA DATA

ATTACHMENT A: WIC QUALITY ASSURANCE DELIVERABLES

Transylvania County

Process Outcome Objectives	2003 to 2005	2006 to 2008	2009 to 2011	% Change from 2003 to 2011	2011 Data		
					Num	Den	Percent
1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2)	77.1%	71.8%	68.8%	-10.8%	122	177	68.9%
1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services.	71.4%	65.8%	65.5%	-8.3%	133	205	64.9%
1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services.	58.4%	53.0%	57.4%	-1.8%	541	912	59.3%
2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy.	50.0%	50.9%	47.1%	-5.8%	177	375	47.2%
2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum.	78.2%	75.4%	72.6%	-7.2%	127	186	68.3%
3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1)	33.3%	50.0%	100.0%	200.0%			
3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1)	85.7%	83.9%	58.8%	-31.4%			
4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were:							
a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1)	67.0%	67.5%	64.8%	-3.3%	165	265	62.3%
b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile)	15.9%	15.8%	16.7%	5.1%	48	265	18.1%
c. Obese (BMI at or greater than the 95th percentile).	14.0%	14.0%	14.8%	5.1%	43	265	16.2%
d. Underweight (BMI less than the 5th percentile).	3.0%	2.7%	3.7%	22.7%	9	265	3.4%
4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3)							
a. Recommended prenatal weight gain	29.0%	33.0%	30.1%	3.7%	47	163	28.8%
b. Excessive prenatal weight gain	49.6%	47.8%	49.1%	-1.1%	80	163	49.1%
c. Inadequate prenatal weight gain	21.3%	19.2%	20.8%	-2.4%	36	163	22.1%
5A. Percentage of women who participated in WIC who initiated breastfeeding.	80.9%	75.6%	81.1%	0.3%	181	222	81.5%
5B. Percentage of infants who participated in WIC who were:							
a. Breastfeeding at 6 weeks of age (1)	49.5%	53.1%	49.9%	0.9%	99	226	43.8%
b. Breastfeeding at 6 months of age	29.4%	30.8%	29.8%	1.3%	61	226	27.0%

(1) Item appears also on other WCH Section Agreement Addenda.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI >=30.0): Adapted from the National Academy of Sciences. Weight Gain During Pregnancy: Reexamining the Guidelines. Washington, DC : National Academy Press, 2009.

SFY 2013-2014 AGREEMENT ADDENDA DATA

ATTACHMENT A: WIC QUALITY ASSURANCE DELIVERABLES

Union County

Process Outcome Objectives	2003 to 2005	2006 to 2008	2009 to 2011	% Change from 2003 to 2011	2011 Data		
					Num	Den	Percent
1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2)	62.1%	57.1%	57.3%	-7.8%	500	866	57.7%
1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services.	61.8%	64.5%	67.2%	8.8%	880	1,292	68.1%
1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services.	52.0%	49.8%	54.5%	4.7%	3,281	5,637	58.2%
2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy.	27.4%	18.1%	15.0%	-45.1%	314	1,258	25.0%
2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum.	76.9%	78.0%	80.7%	5.0%	712	875	81.4%
3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1)	89.7%	87.1%	89.2%	-0.6%			
3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1)	89.2%	85.8%	87.2%	-2.2%			
4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were:							
a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1)	66.1%	66.5%	59.2%	-10.4%	1,223	1,906	64.2%
b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile)	16.6%	15.7%	18.8%	13.0%	344	1,906	18.0%
c. Obese (BMI at or greater than the 95th percentile).	14.2%	14.5%	19.6%	38.2%	296	1,906	15.5%
d. Underweight (BMI less than the 5th percentile).	3.1%	3.3%	2.4%	-22.3%	43	1,906	2.3%
4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3)							
a. Recommended prenatal weight gain	34.9%	38.1%	31.9%	-8.7%	290	983	29.5%
b. Excessive prenatal weight gain	41.6%	40.0%	44.1%	6.0%	428	983	43.5%
c. Inadequate prenatal weight gain	23.5%	21.8%	24.0%	2.2%	265	983	27.0%
5A. Percentage of women who participated in WIC who initiated breastfeeding.	68.9%	68.6%	71.0%	3.0%	906	1,232	73.5%
5B. Percentage of infants who participated in WIC who were:							
a. Breastfeeding at 6 weeks of age (1)	35.6%	36.2%	38.0%	7.0%	496	1,237	40.1%
b. Breastfeeding at 6 months of age	17.5%	20.0%	22.6%	29.0%	289	1,237	23.4%

(1) Item appears also on other WCH Section Agreement Addenda.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI >=30.0): Adapted from the National Academy of Sciences. Weight Gain During Pregnancy: Reexamining the Guidelines. Washington, DC : National Academy Press, 2009.

SFY 2013-2014 AGREEMENT ADDENDA DATA

ATTACHMENT A: WIC QUALITY ASSURANCE DELIVERABLES

Wake County

Process Outcome Objectives	2003 to 2005	2006 to 2008	2009 to 2011	% Change from 2003 to 2011	2011 Data		
					Num	Den	Percent
1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2)	70.2%	68.0%	67.7%	-3.6%	2,226	3,355	66.3%
1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services.	67.4%	67.7%	67.2%	-0.3%	3,522	5,359	65.7%
1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services.	51.6%	50.8%	55.7%	8.1%	13,500	23,005	58.7%
2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy.	19.1%	19.7%	17.8%	-7.1%	1,997	10,613	18.8%
2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum.	68.3%	71.8%	71.8%	5.1%	3,135	4,220	74.3%
3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1)	86.5%	84.4%	89.8%	3.8%			
3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1)	63.5%	63.4%	69.4%	9.3%			
4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were:							
a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1)	64.5%	63.8%	63.0%	-2.2%	5,476	8,575	63.9%
b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile)	16.4%	17.3%	17.0%	3.5%	1,446	8,575	16.9%
c. Obese (BMI at or greater than the 95th percentile).	16.2%	16.4%	16.4%	1.4%	1,347	8,575	15.7%
d. Underweight (BMI less than the 5th percentile).	2.9%	2.5%	3.5%	22.5%	306	8,575	3.6%
4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3)							
a. Recommended prenatal weight gain	33.3%	32.8%	31.1%	-6.5%	1,183	4,155	28.5%
b. Excessive prenatal weight gain	42.8%	39.8%	46.8%	9.3%	2,018	4,155	48.6%
c. Inadequate prenatal weight gain	23.9%	27.4%	22.1%	-7.7%	954	4,155	23.0%
5A. Percentage of women who participated in WIC who initiated breastfeeding.	57.2%	68.3%	69.0%	20.6%	3,827	5,410	70.7%
5B. Percentage of infants who participated in WIC who were:							
a. Breastfeeding at 6 weeks of age (1)	39.4%	56.3%	55.0%	39.6%	2,689	4,850	55.4%
b. Breastfeeding at 6 months of age	29.5%	32.5%	30.5%	3.4%	1,398	4,850	28.8%

(1) Item appears also on other WCH Section Agreement Addenda.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI >=30.0): Adapted from the National Academy of Sciences. Weight Gain During Pregnancy: Reexamining the Guidelines. Washington, DC : National Academy Press, 2009.

SFY 2013-2014 AGREEMENT ADDENDA DATA

ATTACHMENT A: WIC QUALITY ASSURANCE DELIVERABLES

Warren County

Process Outcome Objectives	2003 to 2005	2006 to 2008	2009 to 2011	% Change from 2003 to 2011	2011 Data		
					Num	Den	Percent
1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2)	86.1%	85.5%	82.5%	-4.1%	95	118	80.5%
1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services.	76.1%	78.4%	77.1%	1.3%	118	145	81.4%
1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services.	68.9%	66.7%	66.6%	-3.4%	541	817	66.2%
2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy.	49.7%	46.4%	42.0%	-15.4%	108	279	38.7%
2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum.	80.0%	78.2%	71.2%	-11.0%	96	136	70.6%
3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1)	97.7%	98.8%	98.3%	0.6%			
3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1)	95.9%	92.5%	98.6%	2.7%			
4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were:							
a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1)	70.4%	70.9%	67.7%	-3.8%	220	330	66.7%
b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile)	13.3%	10.0%	12.8%	-4.4%	55	330	16.7%
c. Obese (BMI at or greater than the 95th percentile).	10.2%	10.6%	10.9%	7.0%	39	330	11.8%
d. Underweight (BMI less than the 5th percentile).	6.1%	8.5%	8.6%	41.9%	16	330	4.8%
4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3)							
a. Recommended prenatal weight gain	31.8%	26.8%	29.2%	-8.1%	35	117	29.9%
b. Excessive prenatal weight gain	42.1%	44.1%	46.7%	10.9%	55	117	47.0%
c. Inadequate prenatal weight gain	26.1%	29.1%	24.1%	-7.9%	27	117	23.1%
5A. Percentage of women who participated in WIC who initiated breastfeeding.	34.1%	37.7%	47.2%	38.5%	95	171	55.6%
5B. Percentage of infants who participated in WIC who were:							
a. Breastfeeding at 6 weeks of age (1)	22.3%	27.9%	26.5%	19.0%	56	195	28.7%
b. Breastfeeding at 6 months of age	11.3%	14.2%	15.0%	32.7%	39	195	20.0%

(1) Item appears also on other WCH Section Agreement Addenda.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI >=30.0): Adapted from the National Academy of Sciences. Weight Gain During Pregnancy: Reexamining the Guidelines. Washington, DC : National Academy Press, 2009.

SFY 2013-2014 AGREEMENT ADDENDA DATA

ATTACHMENT A: WIC QUALITY ASSURANCE DELIVERABLES

Wayne County

Process Outcome Objectives	2003 to 2005	2006 to 2008	2009 to 2011	% Change from 2003 to 2011	2011 Data		
					Num	Den	Percent
1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2)	71.6%	78.6%	70.0%	-2.3%	617	921	67.0%
1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services.	72.8%	77.1%	72.3%	-0.6%	871	1,239	70.3%
1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services.	55.1%	53.2%	53.6%	-2.8%	2,711	4,924	55.1%
2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy.	30.9%	37.4%	31.4%	1.5%	754	2,634	28.6%
2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum.	67.8%	71.6%	70.3%	3.8%	880	1,320	66.7%
3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1)	93.1%	85.0%					
3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1)	85.6%	83.3%					
4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were:							
a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1)	64.1%	63.8%	65.4%	2.0%	939	1,488	63.1%
b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile)	16.9%	17.4%	15.1%	-10.6%	247	1,488	16.6%
c. Obese (BMI at or greater than the 95th percentile).	16.2%	16.3%	15.6%	-3.8%	241	1,488	16.2%
d. Underweight (BMI less than the 5th percentile).	2.8%	2.5%	3.9%	41.1%	61	1,488	4.1%
4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3)							
a. Recommended prenatal weight gain	31.3%	31.5%	31.2%	-0.5%	317	1,036	30.6%
b. Excessive prenatal weight gain	47.2%	45.0%	44.0%	-6.7%	461	1,036	44.5%
c. Inadequate prenatal weight gain	21.5%	23.5%	24.8%	15.4%	258	1,036	24.9%
5A. Percentage of women who participated in WIC who initiated breastfeeding.	51.9%	49.1%	51.5%	-0.8%	857	1,506	56.9%
5B. Percentage of infants who participated in WIC who were:							
a. Breastfeeding at 6 weeks of age (1)	26.8%	28.1%	29.4%	9.8%	417	1,331	31.3%
b. Breastfeeding at 6 months of age	11.5%	11.4%	13.9%	20.7%	211	1,331	15.9%

(1) Item appears also on other WCH Section Agreement Addenda.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI < 18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI ≥ 30.0): Adapted from the National Academy of Sciences. *Weight Gain During Pregnancy: Reexamining the Guidelines*. Washington, DC : National Academy Press, 2009.

SFY 2013-2014 AGREEMENT ADDENDA DATA

ATTACHMENT A: WIC QUALITY ASSURANCE DELIVERABLES

Wilkes County

Process Outcome Objectives	2003 to 2005	2006 to 2008	2009 to 2011	% Change from 2003 to 2011	2011 Data		
					Num	Den	Percent
1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2)	79.0%	79.7%	79.3%	0.4%	335	436	76.8%
1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services.	78.6%	80.9%	77.2%	-1.8%	368	513	71.7%
1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services.	62.3%	60.4%	63.3%	1.6%	1,470	2,336	62.9%
2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy.	42.5%	39.9%	39.5%	-7.1%	311	775	40.1%
2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum.	81.5%	79.9%	78.4%	-3.9%	343	450	76.2%
3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1)	84.6%	91.3%	97.2%	14.9%			
3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1)	84.6%	83.7%	82.8%	-2.1%			
4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were:							
a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1)	61.5%	63.7%	60.7%	-1.2%	519	838	61.9%
b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile)	17.8%	17.6%	17.8%	0.1%	138	838	16.5%
c. Obese (BMI at or greater than the 95th percentile).	18.7%	16.7%	17.5%	-6.2%	139	838	16.6%
d. Underweight (BMI less than the 5th percentile).	2.1%	2.0%	3.9%	89.8%	42	838	5.0%
4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3)							
a. Recommended prenatal weight gain	30.0%	31.6%	30.5%	1.7%	116	422	27.5%
b. Excessive prenatal weight gain	50.1%	38.0%	45.2%	-9.8%	206	422	48.8%
c. Inadequate prenatal weight gain	19.9%	30.3%	24.3%	22.1%	100	422	23.7%
5A. Percentage of women who participated in WIC who initiated breastfeeding.	47.0%	54.2%	59.5%	26.7%	307	523	58.7%
5B. Percentage of infants who participated in WIC who were:							
a. Breastfeeding at 6 weeks of age (1)	26.4%	25.4%	25.8%	-2.6%	133	543	24.5%
b. Breastfeeding at 6 months of age	12.8%	11.5%	12.3%	-3.7%	73	543	13.4%

(1) Item appears also on other WCH Section Agreement Addenda.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI >=30.0): Adapted from the National Academy of Sciences. Weight Gain During Pregnancy: Reexamining the Guidelines. Washington, DC : National Academy Press, 2009.

SFY 2013-2014 AGREEMENT ADDENDA DATA

ATTACHMENT A: WIC QUALITY ASSURANCE DELIVERABLES

Wilson County

Process Outcome Objectives	2003 to 2005	2006 to 2008	2009 to 2011	% Change from 2003 to 2011	2011 Data		
					Num	Den	Percent
1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2)	83.4%	84.1%	84.5%	1.3%	512	615	83.3%
1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services.	78.2%	75.9%	75.8%	-3.1%	618	823	75.1%
1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services.	57.2%	57.6%	57.2%	-0.1%	2,122	3,592	59.1%
2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy.	30.2%	24.6%	20.8%	-31.1%	371	1,700	21.8%
2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum.	67.7%	68.5%	69.8%	3.1%	516	725	71.2%
3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1)	99.3%	99.2%	100.0%	0.8%			
3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1)	97.3%	97.2%	97.0%	-0.3%			
4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were:							
a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1)	62.4%	62.8%	62.7%	0.5%	889	1,382	64.3%
b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile)	16.8%	18.3%	16.2%	-3.9%	202	1,382	14.6%
c. Obese (BMI at or greater than the 95th percentile).	18.2%	16.7%	17.1%	-6.1%	242	1,382	17.5%
d. Underweight (BMI less than the 5th percentile).	2.7%	2.1%	4.1%	54.0%	49	1,382	3.5%
4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3)							
a. Recommended prenatal weight gain	30.5%	32.4%	32.6%	6.7%	238	687	34.6%
b. Excessive prenatal weight gain	41.3%	39.1%	43.1%	4.4%	301	687	43.8%
c. Inadequate prenatal weight gain	28.2%	28.5%	24.3%	-13.8%	148	687	21.5%
5A. Percentage of women who participated in WIC who initiated breastfeeding.	38.5%	41.9%	38.1%	-0.9%	370	839	44.1%
5B. Percentage of infants who participated in WIC who were:							
a. Breastfeeding at 6 weeks of age (1)	24.0%	25.7%	23.2%	-3.0%	209	851	24.6%
b. Breastfeeding at 6 months of age	12.1%	13.4%	15.7%	29.2%	148	851	17.4%

(1) Item appears also on other WCH Section Agreement Addenda.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI >=30.0): Adapted from the National Academy of Sciences. Weight Gain During Pregnancy: Reexamining the Guidelines. Washington, DC : National Academy Press, 2009.

SFY 2013-2014 AGREEMENT ADDENDA DATA

ATTACHMENT A: WIC QUALITY ASSURANCE DELIVERABLES

Yadkin County

Process Outcome Objectives	2003 to 2005	2006 to 2008	2009 to 2011	% Change from 2003 to 2011	2011 Data		
					Num	Den	Percent
1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2)	72.8%	73.4%	75.0%	3.1%	153	204	75.0%
1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services.	67.8%	73.0%	76.7%	13.2%	236	296	79.7%
1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services.	55.6%	58.7%	63.5%	14.3%	884	1,333	66.3%
2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy.	48.0%	44.8%	43.7%	-8.9%	233	538	43.3%
2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum.	71.5%	77.9%	78.2%	9.4%	208	265	78.5%
3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1)	98.5%	98.5%	96.9%	-1.6%			
3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1)	97.7%	95.8%	96.5%	-1.2%			
4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were:							
a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1)	62.0%	64.3%	63.0%	1.5%	336	535	62.8%
b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile)	16.5%	16.0%	16.4%	-1.1%	91	535	17.0%
c. Obese (BMI at or greater than the 95th percentile).	18.6%	16.9%	18.3%	-2.0%	96	535	17.9%
d. Underweight (BMI less than the 5th percentile).	2.8%	2.7%	2.4%	-14.1%	12	535	2.2%
4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3)							
a. Recommended prenatal weight gain	36.4%	33.3%	31.3%	-14.1%	76	242	31.4%
b. Excessive prenatal weight gain	44.1%	49.3%	45.5%	3.2%	117	242	48.3%
c. Inadequate prenatal weight gain	19.5%	17.4%	23.2%	19.0%	49	242	20.2%
5A. Percentage of women who participated in WIC who initiated breastfeeding.	65.5%	69.2%	70.5%	7.6%	231	342	67.5%
5B. Percentage of infants who participated in WIC who were:							
a. Breastfeeding at 6 weeks of age (1)	37.5%	43.3%	42.1%	12.1%	147	322	45.7%
b. Breastfeeding at 6 months of age	16.2%	21.8%	22.4%	37.9%	82	322	25.5%

(1) Item appears also on other WCH Section Agreement Addenda.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI >=30.0): Adapted from the National Academy of Sciences. Weight Gain During Pregnancy: Reexamining the Guidelines. Washington, DC : National Academy Press, 2009.

SFY 2013-2014 AGREEMENT ADDENDA DATA

ATTACHMENT A: WIC QUALITY ASSURANCE DELIVERABLES

Allegheny

Process Outcome Objectives	2003 to 2005	2006 to 2008	2009 to 2011	% Change from 2003 to 2011	2011 Data		
					Num	Den	Percent
1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2)	85.2%	74.1%	73.6%	-13.7%	38	59	64.4%
1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services.	76.3%	77.4%	84.1%	10.3%	64	70	91.4%
1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services.	72.4%	68.5%	69.4%	-4.2%	247	345	71.6%
2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy.	49.8%	46.4%	41.4%	-16.8%	46	122	37.7%
2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum.	74.7%	75.6%	76.5%	2.5%	48	63	76.2%
3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1)	90.2%	93.7%					
3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1)	91.5%	89.9%					
4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were:							
a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1)	59.8%	55.3%	55.1%	-7.8%	84	169	49.7%
b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile)	18.4%	18.8%	20.8%	13.1%	34	169	20.1%
c. Obese (BMI at or greater than the 95th percentile).	20.3%	24.5%	22.2%	9.1%	48	169	28.4%
d. Underweight (BMI less than the 5th percentile).	1.5%	1.4%	1.9%	28.2%	3	169	1.8%
4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3)							
a. Recommended prenatal weight gain	34.6%	33.8%	31.2%	-9.8%	19	56	33.9%
b. Excessive prenatal weight gain	47.1%	49.3%	48.2%	2.5%	22	56	39.3%
c. Inadequate prenatal weight gain	18.4%	16.9%	20.6%	12.0%	15	56	26.8%
5A. Percentage of women who participated in WIC who initiated breastfeeding.	56.5%	65.2%	36.5%	-35.3%	47	82	57.3%
5B. Percentage of infants who participated in WIC who were:							
a. Breastfeeding at 6 weeks of age (1)	15.4%	27.8%	14.1%	-8.3%	14	94	14.9%
b. Breastfeeding at 6 months of age	4.3%	15.8%	6.0%	38.2%	8	94	8.5%

(1) Item appears also on other WCH Section Agreement Addenda.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI < 18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI ≥ 30.0): Adapted from the National Academy of Sciences. *Weight Gain During Pregnancy: Reexamining the Guidelines*. Washington, DC : National Academy Press, 2009.

SFY 2013-2014 AGREEMENT ADDENDA DATA

ATTACHMENT A: WIC QUALITY ASSURANCE DELIVERABLES

Ashe

Process Outcome Objectives	2003 to 2005	2006 to 2008	2009 to 2011	% Change from 2003 to 2011	2011 Data		
					Num	Den	Percent
1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2)	77.6%	75.3%	71.5%	-7.9%	97	150	64.7%
1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services.	73.2%	72.8%	68.6%	-6.4%	113	176	64.2%
1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services.	70.3%	62.1%	61.9%	-11.9%	468	770	60.8%
2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy.	52.6%	52.3%	42.7%	-18.8%	121	295	41.0%
2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum.	73.8%	81.0%	77.8%	5.4%	123	159	77.4%
3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1)	89.9%	93.2%					
3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1)	81.9%	84.4%					
4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were:							
a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1)	70.9%	64.5%	66.4%	-6.4%	120	178	67.4%
b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile)	11.3%	14.9%	15.3%	34.5%	24	178	13.5%
c. Obese (BMI at or greater than the 95th percentile).	11.7%	16.0%	12.9%	10.9%	26	178	14.6%
d. Underweight (BMI less than the 5th percentile).	6.1%	4.6%	5.5%	-11.1%	8	178	4.5%
4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3)							
a. Recommended prenatal weight gain	30.3%	31.4%	29.4%	-3.0%	42	122	34.4%
b. Excessive prenatal weight gain	46.5%	45.5%	46.4%	-0.2%	55	122	45.1%
c. Inadequate prenatal weight gain	23.1%	23.1%	24.2%	4.4%	25	122	20.5%
5A. Percentage of women who participated in WIC who initiated breastfeeding.	48.4%	50.2%	57.9%	19.6%	97	183	53.0%
5B. Percentage of infants who participated in WIC who were:							
a. Breastfeeding at 6 weeks of age (1)	30.8%	28.8%	30.5%	-1.2%	60	188	31.9%
b. Breastfeeding at 6 months of age	13.4%	13.1%	19.3%	43.9%	41	188	21.8%

(1) Item appears also on other WCH Section Agreement Addenda.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI < 18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI ≥ 30.0): Adapted from the National Academy of Sciences. *Weight Gain During Pregnancy: Reexamining the Guidelines*. Washington, DC : National Academy Press, 2009.

SFY 2013-2014 AGREEMENT ADDENDA DATA

ATTACHMENT A: WIC QUALITY ASSURANCE DELIVERABLES

Avery

Process Outcome Objectives	2003 to 2005	2006 to 2008	2009 to 2011	% Change from 2003 to 2011	2011 Data		
					Num	Den	Percent
1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2)	73.0%	73.6%	70.8%	-3.1%	57	90	63.3%
1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services.	60.9%	65.5%	68.9%	13.2%	81	116	69.8%
1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services.	65.8%	60.3%	64.8%	-1.5%	293	452	64.8%
2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy.	46.8%	46.7%	44.0%	-5.9%	104	241	43.2%
2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum.	72.2%	68.6%	68.5%	-5.2%	73	108	67.6%
3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1)	100.0%	93.2%	100.0%	0.0%			
3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1)	83.3%	86.5%	87.5%	5.0%			
4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were:							
a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1)	65.8%	62.4%	66.5%	1.1%	124	179	69.3%
b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile)	15.6%	18.8%	16.0%	2.0%	28	179	15.6%
c. Obese (BMI at or greater than the 95th percentile).	17.2%	15.9%	12.8%	-25.5%	18	179	10.1%
d. Underweight (BMI less than the 5th percentile).	1.4%	3.0%	4.7%	243.3%	9	179	5.0%
4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3)							
a. Recommended prenatal weight gain	31.4%	30.1%	32.6%	3.8%	33	98	33.7%
b. Excessive prenatal weight gain	45.0%	46.5%	46.9%	4.1%	45	98	45.9%
c. Inadequate prenatal weight gain	23.6%	23.4%	20.5%	-12.9%	20	98	20.4%
5A. Percentage of women who participated in WIC who initiated breastfeeding.	56.9%	55.2%	58.7%	3.2%	82	131	62.6%
5B. Percentage of infants who participated in WIC who were:							
a. Breastfeeding at 6 weeks of age (1)	38.3%	32.5%	38.3%	0.0%	54	143	37.8%
b. Breastfeeding at 6 months of age	28.2%	22.8%	24.8%	-12.2%	37	143	25.9%

(1) Item appears also on other WCH Section Agreement Addenda.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI < 18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI ≥ 30.0): Adapted from the National Academy of Sciences. *Weight Gain During Pregnancy: Reexamining the Guidelines*. Washington, DC : National Academy Press, 2009.

SFY 2013-2014 AGREEMENT ADDENDA DATA

ATTACHMENT A: WIC QUALITY ASSURANCE DELIVERABLES

Bertie

Process Outcome Objectives	2003 to 2005	2006 to 2008	2009 to 2011	% Change from 2003 to 2011	2011 Data		
					Num	Den	Percent
1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2)	82.0%	82.6%	78.8%	-3.9%	120	159	75.5%
1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services.	73.0%	72.3%	71.3%	-2.3%	119	165	72.1%
1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services.	66.0%	61.3%	58.8%	-10.9%	422	691	61.1%
2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy.	39.2%	28.3%	24.2%	-38.2%	80	326	24.5%
2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum.	47.4%	27.0%	30.0%	-36.7%	44	155	28.4%
3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1)	63.5%	39.2%					
3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1)	71.2%	45.6%					
4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were:							
a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1)	64.3%	63.1%	64.9%	1.0%	213	322	66.1%
b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile)	13.0%	14.0%	12.6%	-3.0%	39	322	12.1%
c. Obese (BMI at or greater than the 95th percentile).	15.3%	17.1%	14.2%	-7.0%	41	322	12.7%
d. Underweight (BMI less than the 5th percentile).	7.5%	5.9%	8.3%	10.9%	29	322	9.0%
4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3)							
a. Recommended prenatal weight gain	26.0%	29.0%	27.1%	4.6%	24	127	18.9%
b. Excessive prenatal weight gain	51.1%	45.6%	35.9%	-29.6%	34	127	26.8%
c. Inadequate prenatal weight gain	23.0%	25.4%	36.9%	60.7%	69	127	54.3%
5A. Percentage of women who participated in WIC who initiated breastfeeding.	26.5%	13.2%	28.4%	7.0%	76	189	40.2%
5B. Percentage of infants who participated in WIC who were:							
a. Breastfeeding at 6 weeks of age (1)	8.4%	6.6%	8.2%	-2.3%	18	177	10.2%
b. Breastfeeding at 6 months of age	2.9%	2.4%	2.0%	-30.7%	4	177	2.3%

(1) Item appears also on other WCH Section Agreement Addenda.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI < 18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI ≥ 30.0): Adapted from the National Academy of Sciences. *Weight Gain During Pregnancy: Reexamining the Guidelines*. Washington, DC : National Academy Press, 2009.

SFY 2013-2014 AGREEMENT ADDENDA DATA

ATTACHMENT A: WIC QUALITY ASSURANCE DELIVERABLES

Camden

Process Outcome Objectives	2003 to 2005	2006 to 2008	2009 to 2011	% Change from 2003 to 2011	2011 Data		
					Num	Den	Percent
1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2)	80.0%	78.3%	79.6%	-0.5%	21	31	67.7%
1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services.	69.0%	75.8%	72.2%	4.7%	24	34	70.6%
1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services.	60.2%	56.9%	54.6%	-9.3%	91	162	56.2%
2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy.	39.2%	35.4%	28.8%	-26.6%	408	1,586	25.7%
2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum.	34.6%	38.2%	48.2%	39.5%	305	614	49.7%
3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1)	63.6%	81.8%					
3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1)	64.4%	53.8%					
4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were:							
a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1)	68.6%	69.7%	66.1%	-3.7%	83	117	70.9%
b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile)	6.5%	6.3%	12.6%	95.7%	8	117	6.8%
c. Obese (BMI at or greater than the 95th percentile).	7.9%	12.9%	15.3%	93.0%	16	117	13.7%
d. Underweight (BMI less than the 5th percentile).	17.0%	11.1%	6.0%	-64.8%	10	117	8.5%
4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3)							
a. Recommended prenatal weight gain	36.8%	30.4%	26.0%	-29.4%	8	30	26.7%
b. Excessive prenatal weight gain	39.1%	42.4%	49.0%	25.5%	18	30	60.0%
c. Inadequate prenatal weight gain	24.1%	27.2%	25.0%	3.6%	4	30	13.3%
5A. Percentage of women who participated in WIC who initiated breastfeeding.	24.7%	21.2%	29.4%	19.0%	284	799	35.5%
5B. Percentage of infants who participated in WIC who were:							
a. Breastfeeding at 6 weeks of age (1)	9.7%	9.9%	13.5%	38.8%	132	758	17.4%
b. Breastfeeding at 6 months of age	3.9%	3.5%	6.2%	60.0%	84	758	11.1%

(1) Item appears also on other WCH Section Agreement Addenda.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI < 18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI ≥ 30.0): Adapted from the National Academy of Sciences. *Weight Gain During Pregnancy: Reexamining the Guidelines*. Washington, DC : National Academy Press, 2009.

SFY 2013-2014 AGREEMENT ADDENDA DATA

ATTACHMENT A: WIC QUALITY ASSURANCE DELIVERABLES

Chowan

Process Outcome Objectives	2003 to 2005	2006 to 2008	2009 to 2011	% Change from 2003 to 2011	2011 Data		
					Num	Den	Percent
1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2)	82.0%	81.0%	79.5%	-3.1%	90	109	82.6%
1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services.	83.2%	84.0%	85.2%	2.4%	100	121	82.6%
1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services.	67.5%	65.2%	68.6%	1.5%	336	473	71.0%
2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy.	39.2%	35.4%	28.8%	-26.6%	408	1,586	25.7%
2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum.	34.6%	38.2%	48.2%	39.5%	305	614	49.7%
3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1)	88.9%	75.4%					
3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1)	84.7%	85.4%					
4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were:							
a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1)	63.7%	65.6%	61.2%	-3.9%	132	223	59.2%
b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile)	17.9%	15.2%	13.7%	-23.5%	29	223	13.0%
c. Obese (BMI at or greater than the 95th percentile).	16.7%	15.6%	18.2%	8.7%	45	223	20.2%
d. Underweight (BMI less than the 5th percentile).	1.6%	3.5%	6.9%	323.7%	17	223	7.6%
4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3)							
a. Recommended prenatal weight gain	33.3%	34.7%	30.2%	-9.3%	28	103	27.2%
b. Excessive prenatal weight gain	41.4%	43.6%	45.7%	10.4%	51	103	49.5%
c. Inadequate prenatal weight gain	25.3%	21.6%	24.1%	-4.8%	24	103	23.3%
5A. Percentage of women who participated in WIC who initiated breastfeeding.	24.7%	21.2%	29.4%	19.0%	284	799	35.5%
5B. Percentage of infants who participated in WIC who were:							
a. Breastfeeding at 6 weeks of age (1)	9.7%	9.9%	13.5%	38.8%	132	758	17.4%
b. Breastfeeding at 6 months of age	3.9%	3.5%	6.2%	60.0%	84	758	11.1%

(1) Item appears also on other WCH Section Agreement Addenda.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI < 18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI ≥ 30.0): Adapted from the National Academy of Sciences. *Weight Gain During Pregnancy: Reexamining the Guidelines*. Washington, DC : National Academy Press, 2009.

SFY 2013-2014 AGREEMENT ADDENDA DATA

ATTACHMENT A: WIC QUALITY ASSURANCE DELIVERABLES

Currituck

Process Outcome Objectives	2003 to 2005	2006 to 2008	2009 to 2011	% Change from 2003 to 2011	2011 Data		
					Num	Den	Percent
1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2)	78.1%	71.9%	66.0%	-15.5%	53	99	53.5%
1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services.	75.5%	67.0%	67.9%	-10.0%	84	119	70.6%
1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services.	56.5%	51.3%	49.9%	-11.6%	267	512	52.1%
2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy.	39.2%	35.4%	28.8%	-26.6%	408	1,586	25.7%
2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum.	34.6%	38.2%	48.2%	39.5%	305	614	49.7%
3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1)	89.3%	82.7%					
3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1)	83.6%	81.9%					
4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were:							
a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1)	67.0%	65.5%	68.3%	2.0%	109	178	61.2%
b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile)	16.9%	15.4%	15.9%	-5.7%	34	178	19.1%
c. Obese (BMI at or greater than the 95th percentile).	13.4%	16.0%	11.9%	-11.1%	25	178	14.0%
d. Underweight (BMI less than the 5th percentile).	2.7%	3.1%	3.8%	40.4%	10	178	5.6%
4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3)							
a. Recommended prenatal weight gain	36.8%	27.7%	30.7%	-16.4%	10	55	18.2%
b. Excessive prenatal weight gain	45.8%	50.3%	44.7%	-2.4%	28	55	50.9%
c. Inadequate prenatal weight gain	17.4%	22.0%	24.6%	41.1%	17	55	30.9%
5A. Percentage of women who participated in WIC who initiated breastfeeding.	24.7%	21.2%	29.4%	19.0%	284	799	35.5%
5B. Percentage of infants who participated in WIC who were:							
a. Breastfeeding at 6 weeks of age (1)	9.7%	9.9%	13.5%	38.8%	132	758	17.4%
b. Breastfeeding at 6 months of age	3.9%	3.5%	6.2%	60.0%	84	758	11.1%

(1) Item appears also on other WCH Section Agreement Addenda.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI < 18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI ≥ 30.0): Adapted from the National Academy of Sciences. *Weight Gain During Pregnancy: Reexamining the Guidelines*. Washington, DC : National Academy Press, 2009.

SFY 2013-2014 AGREEMENT ADDENDA DATA

ATTACHMENT A: WIC QUALITY ASSURANCE DELIVERABLES

Gates

Process Outcome Objectives	2003 to 2005	2006 to 2008	2009 to 2011	% Change from 2003 to 2011	2011 Data		
					Num	Den	Percent
1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2)	81.1%	80.0%	54.0%	-33.4%	19	57	33.3%
1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services.	73.4%	73.7%	77.8%	6.1%	54	68	79.4%
1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services.	59.8%	60.6%	62.4%	4.3%	188	288	65.3%
2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy.	28.4%	28.0%	27.1%	-4.7%	36	127	28.3%
2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum.	45.4%	57.9%	71.8%	58.3%	48	64	75.0%
3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1)	86.7%	92.3%					
3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1)	85.1%	90.6%					
4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were:							
a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1)	63.7%	65.1%	64.3%	0.9%	27	43	62.8%
b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile)	16.6%	14.1%	15.1%	-9.3%	9	43	20.9%
c. Obese (BMI at or greater than the 95th percentile).	15.7%	17.0%	13.1%	-16.8%	3	43	7.0%
d. Underweight (BMI less than the 5th percentile).	3.9%	3.7%	7.5%	91.9%	4	43	9.3%
4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3)							
a. Recommended prenatal weight gain	28.8%	37.7%	32.4%	12.8%	12	28	42.9%
b. Excessive prenatal weight gain	41.3%	46.8%	33.8%	-18.1%	7	28	25.0%
c. Inadequate prenatal weight gain	30.0%	15.6%	33.8%	12.6%	9	28	32.1%
5A. Percentage of women who participated in WIC who initiated breastfeeding.	29.8%	30.7%	13.2%	-55.8%	12	85	14.1%
5B. Percentage of infants who participated in WIC who were:							
a. Breastfeeding at 6 weeks of age (1)	12.2%	10.6%	11.6%	-4.3%	10	86	11.6%
b. Breastfeeding at 6 months of age	5.7%	3.2%	2.0%	-64.5%	2	86	2.3%

(1) Item appears also on other WCH Section Agreement Addenda.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI < 18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI ≥ 30.0): Adapted from the National Academy of Sciences. *Weight Gain During Pregnancy: Reexamining the Guidelines*. Washington, DC : National Academy Press, 2009.

SFY 2013-2014 AGREEMENT ADDENDA DATA

ATTACHMENT A: WIC QUALITY ASSURANCE DELIVERABLES

Granville

Process Outcome Objectives	2003 to 2005	2006 to 2008	2009 to 2011	% Change from 2003 to 2011	2011 Data		
					Num	Den	Percent
1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2)	78.1%	73.4%	76.3%	-2.3%	210	273	76.9%
1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services.	78.7%	74.8%	76.2%	-3.1%	285	362	78.7%
1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services.	61.0%	58.1%	60.3%	-1.2%	922	1,534	60.1%
2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy.	32.6%	32.7%	29.0%	-11.1%	408	1,500	27.2%
2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum.	81.7%	77.8%	73.9%	-9.5%	558	735	75.9%
3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1)	92.6%	93.0%	94.7%	2.3%			
3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1)	93.1%	90.1%	83.6%	-10.1%			
4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were:							
a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1)	61.2%	63.0%	65.6%	7.2%	398	617	64.5%
b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile)	18.2%	18.2%	14.0%	-23.0%	83	617	13.5%
c. Obese (BMI at or greater than the 95th percentile).	16.8%	16.3%	16.4%	-1.9%	104	617	16.9%
d. Underweight (BMI less than the 5th percentile).	3.9%	2.6%	3.9%	2.0%	32	617	5.2%
4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3)							
a. Recommended prenatal weight gain	27.3%	31.0%	31.0%	13.3%	83	309	26.9%
b. Excessive prenatal weight gain	37.4%	44.4%	47.6%	27.3%	156	309	50.5%
c. Inadequate prenatal weight gain	35.3%	24.6%	21.4%	-39.3%	70	309	22.7%
5A. Percentage of women who participated in WIC who initiated breastfeeding.	43.3%	52.9%	41.4%	-4.3%	408	922	44.3%
5B. Percentage of infants who participated in WIC who were:							
a. Breastfeeding at 6 weeks of age (1)	28.7%	34.5%	36.3%	26.6%	320	959	33.4%
b. Breastfeeding at 6 months of age	20.9%	22.5%	26.8%	28.5%	238	959	24.8%

(1) Item appears also on other WCH Section Agreement Addenda.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI < 18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI ≥ 30.0): Adapted from the National Academy of Sciences. *Weight Gain During Pregnancy: Reexamining the Guidelines*. Washington, DC : National Academy Press, 2009.

SFY 2013-2014 AGREEMENT ADDENDA DATA

ATTACHMENT A: WIC QUALITY ASSURANCE DELIVERABLES

Martin

Process Outcome Objectives	2003 to 2005	2006 to 2008	2009 to 2011	% Change from 2003 to 2011	2011 Data		
					Num	Den	Percent
1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2)	75.9%	81.8%	84.2%	11.0%	131	169	77.5%
1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services.	74.7%	68.0%	77.5%	3.8%	159	187	85.0%
1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services.	61.7%	56.6%	60.9%	-1.3%	544	916	59.4%
2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy.	37.4%	32.3%	30.6%	-18.1%	242	911	26.6%
2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum.	79.1%	82.0%	79.7%	0.8%	247	313	78.9%
3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1)	91.1%	92.7%	95.5%	4.8%			
3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1)	92.0%	90.1%	90.2%	-2.0%			
4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were:							
a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1)	64.4%	64.4%	65.3%	1.4%	263	398	66.1%
b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile)	14.8%	16.2%	13.7%	-7.3%	57	398	14.3%
c. Obese (BMI at or greater than the 95th percentile).	14.3%	15.5%	15.7%	9.8%	64	398	16.1%
d. Underweight (BMI less than the 5th percentile).	6.5%	3.9%	5.3%	-19.0%	14	398	3.5%
4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3)							
a. Recommended prenatal weight gain	31.5%	31.5%	27.8%	-11.6%	38	157	24.2%
b. Excessive prenatal weight gain	41.6%	41.4%	43.6%	4.8%	70	157	44.6%
c. Inadequate prenatal weight gain	26.9%	27.2%	28.6%	6.2%	49	157	31.2%
5A. Percentage of women who participated in WIC who initiated breastfeeding.	41.3%	38.1%	26.6%	-35.5%	82	372	22.0%
5B. Percentage of infants who participated in WIC who were:							
a. Breastfeeding at 6 weeks of age (1)	24.6%	23.3%	14.8%	-39.9%	40	354	11.3%
b. Breastfeeding at 6 months of age	11.9%	11.4%	4.7%	-60.8%	15	354	4.2%

(1) Item appears also on other WCH Section Agreement Addenda.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI < 18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI ≥ 30.0): Adapted from the National Academy of Sciences. *Weight Gain During Pregnancy: Reexamining the Guidelines*. Washington, DC : National Academy Press, 2009.

SFY 2013-2014 AGREEMENT ADDENDA DATA

ATTACHMENT A: WIC QUALITY ASSURANCE DELIVERABLES

McDowell

Process Outcome Objectives	2003 to 2005	2006 to 2008	2009 to 2011	% Change from 2003 to 2011	2011 Data		
					Num	Den	Percent
1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2)	66.3%	73.5%	72.1%	8.6%	185	295	62.7%
1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services.	70.9%	73.7%	74.9%	5.6%	287	370	77.6%
1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services.	64.8%	62.7%	62.7%	-3.1%	1,026	1,593	64.4%
2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy.	32.5%	35.2%	33.4%	2.8%	885	2,568	34.5%
2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum.	69.9%	72.4%	76.2%	9.0%	780	981	79.5%
3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1)	85.7%	100.0%	80.0%	-6.7%			
3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1)	87.6%	86.2%	80.0%	-8.7%			
4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were:							
a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1)	68.6%	68.0%	68.3%	-0.4%	423	629	67.2%
b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile)	14.0%	15.6%	13.9%	-0.8%	95	629	15.1%
c. Obese (BMI at or greater than the 95th percentile).	12.4%	13.1%	11.4%	-8.3%	82	629	13.0%
d. Underweight (BMI less than the 5th percentile).	5.0%	3.3%	6.4%	27.9%	29	629	4.6%
4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3)							
a. Recommended prenatal weight gain	35.4%	29.5%	29.2%	-17.6%	86	291	29.6%
b. Excessive prenatal weight gain	34.1%	41.1%	48.4%	42.0%	146	291	50.2%
c. Inadequate prenatal weight gain	30.5%	29.4%	22.5%	-26.4%	59	291	20.3%
5A. Percentage of women who participated in WIC who initiated breastfeeding.	46.4%	54.4%	60.3%	30.0%	701	1,153	60.8%
5B. Percentage of infants who participated in WIC who were:							
a. Breastfeeding at 6 weeks of age (1)	28.1%	31.1%	30.0%	6.6%	338	1,102	30.7%
b. Breastfeeding at 6 months of age	14.7%	14.5%	16.2%	10.0%	190	1,102	17.2%

(1) Item appears also on other WCH Section Agreement Addenda.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI < 18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI ≥ 30.0): Adapted from the National Academy of Sciences. *Weight Gain During Pregnancy: Reexamining the Guidelines*. Washington, DC : National Academy Press, 2009.

SFY 2013-2014 AGREEMENT ADDENDA DATA

ATTACHMENT A: WIC QUALITY ASSURANCE DELIVERABLES

Mitchell

Process Outcome Objectives	2003 to 2005	2006 to 2008	2009 to 2011	% Change from 2003 to 2011	2011 Data		
					Num	Den	Percent
1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2)	77.5%	75.6%	77.6%	0.1%	57	78	73.1%
1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services.	67.0%	63.3%	71.3%	6.3%	75	109	68.8%
1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services.	72.8%	60.8%	65.1%	-10.5%	355	509	69.7%
2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy.	38.1%	41.9%	38.3%	0.6%	82	230	35.7%
2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum.	70.9%	72.4%	65.0%	-8.3%	68	113	60.2%
3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1)	96.8%	75.0%	100.0%	3.3%			
3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1)	92.8%	97.1%	100.0%	7.8%			
4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were:							
a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1)	69.1%	71.2%	64.4%	-6.9%	144	225	64.0%
b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile)	12.6%	12.6%	16.8%	33.7%	44	225	19.6%
c. Obese (BMI at or greater than the 95th percentile).	14.9%	11.7%	15.0%	0.6%	32	225	14.2%
d. Underweight (BMI less than the 5th percentile).	3.4%	4.5%	3.8%	12.4%	5	225	2.2%
4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3)							
a. Recommended prenatal weight gain	37.6%	35.6%	27.4%	-27.0%	28	92	30.4%
b. Excessive prenatal weight gain	41.3%	45.6%	45.9%	11.2%	36	92	39.1%
c. Inadequate prenatal weight gain	21.1%	18.8%	26.6%	26.1%	28	92	30.4%
5A. Percentage of women who participated in WIC who initiated breastfeeding.	53.0%	63.8%	63.5%	19.9%	91	139	65.5%
5B. Percentage of infants who participated in WIC who were:							
a. Breastfeeding at 6 weeks of age (1)	43.5%	42.5%	46.0%	5.7%	55	137	40.1%
b. Breastfeeding at 6 months of age	24.8%	25.8%	28.0%	13.1%	36	137	26.3%

(1) Item appears also on other WCH Section Agreement Addenda.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI < 18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI ≥ 30.0): Adapted from the National Academy of Sciences. *Weight Gain During Pregnancy: Reexamining the Guidelines*. Washington, DC : National Academy Press, 2009.

SFY 2013-2014 AGREEMENT ADDENDA DATA

ATTACHMENT A: WIC QUALITY ASSURANCE DELIVERABLES

Pasquotank

Process Outcome Objectives	2003 to 2005	2006 to 2008	2009 to 2011	% Change from 2003 to 2011	2011 Data		
					Num	Den	Percent
1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2)	81.0%	60.6%	60.1%	-25.8%	142	254	55.9%
1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services.	74.4%	71.2%	72.9%	-1.9%	242	327	74.0%
1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services.	57.7%	53.2%	54.0%	-6.4%	804	1,430	56.2%
2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy.	39.2%	35.4%	28.8%	-26.6%	408	1,586	25.7%
2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum.	34.6%	38.2%	48.2%	39.5%	305	614	49.7%
3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1)	63.8%	63.6%					
3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1)	66.3%	65.3%					
4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were:							
a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1)	66.2%	65.1%	64.0%	-3.4%	300	471	63.7%
b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile)	15.0%	15.3%	15.3%	2.2%	67	471	14.2%
c. Obese (BMI at or greater than the 95th percentile).	13.8%	16.5%	15.2%	10.2%	77	471	16.3%
d. Underweight (BMI less than the 5th percentile).	5.0%	3.1%	5.5%	10.2%	27	471	5.7%
4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3)							
a. Recommended prenatal weight gain	34.0%	32.7%	27.4%	-19.3%	53	226	23.5%
b. Excessive prenatal weight gain	42.2%	42.9%	40.4%	-4.3%	90	226	39.8%
c. Inadequate prenatal weight gain	23.8%	24.5%	32.1%	35.2%	83	226	36.7%
5A. Percentage of women who participated in WIC who initiated breastfeeding.	24.7%	21.2%	29.4%	19.0%	284	799	35.5%
5B. Percentage of infants who participated in WIC who were:							
a. Breastfeeding at 6 weeks of age (1)	9.7%	9.9%	13.5%	38.8%	132	758	17.4%
b. Breastfeeding at 6 months of age	3.9%	3.5%	6.2%	60.0%	84	758	11.1%

(1) Item appears also on other WCH Section Agreement Addenda.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI < 18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI ≥ 30.0): Adapted from the National Academy of Sciences. *Weight Gain During Pregnancy: Reexamining the Guidelines*. Washington, DC : National Academy Press, 2009.

SFY 2013-2014 AGREEMENT ADDENDA DATA

ATTACHMENT A: WIC QUALITY ASSURANCE DELIVERABLES

Perquimans

Process Outcome Objectives	2003 to 2005	2006 to 2008	2009 to 2011	% Change from 2003 to 2011	2011 Data		
					Num	Den	Percent
1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2)	78.9%	71.2%	74.0%	-6.2%	50	71	70.4%
1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services.	76.8%	71.3%	73.9%	-3.8%	68	88	77.3%
1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services.	65.0%	53.9%	55.5%	-14.6%	237	405	58.5%
2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy.	39.2%	35.4%	28.8%	-26.6%	408	1,586	25.7%
2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum.	34.6%	38.2%	48.2%	39.5%	305	614	49.7%
3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1)	85.1%	82.9%					
3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1)	72.8%	82.5%					
4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were:							
a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1)	62.9%	62.3%	58.1%	-7.6%	76	142	53.5%
b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile)	16.2%	18.0%	16.0%	-1.3%	25	142	17.6%
c. Obese (BMI at or greater than the 95th percentile).	17.4%	18.4%	21.4%	22.8%	32	142	22.5%
d. Underweight (BMI less than the 5th percentile).	3.4%	1.3%	4.4%	28.9%	9	142	6.3%
4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3)							
a. Recommended prenatal weight gain	35.3%	34.7%	27.6%	-21.7%	16	63	25.4%
b. Excessive prenatal weight gain	34.0%	44.6%	44.7%	31.5%	32	63	50.8%
c. Inadequate prenatal weight gain	30.7%	20.7%	27.6%	-10.0%	15	63	23.8%
5A. Percentage of women who participated in WIC who initiated breastfeeding.	24.7%	21.2%	29.4%	19.0%	284	799	35.5%
5B. Percentage of infants who participated in WIC who were:							
a. Breastfeeding at 6 weeks of age (1)	9.7%	9.9%	13.5%	38.8%	132	758	17.4%
b. Breastfeeding at 6 months of age	3.9%	3.5%	6.2%	60.0%	84	758	11.1%

(1) Item appears also on other WCH Section Agreement Addenda.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI < 18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI ≥ 30.0): Adapted from the National Academy of Sciences. *Weight Gain During Pregnancy: Reexamining the Guidelines*. Washington, DC : National Academy Press, 2009.

SFY 2013-2014 AGREEMENT ADDENDA DATA

ATTACHMENT A: WIC QUALITY ASSURANCE DELIVERABLES

Polk

Process Outcome Objectives	2003 to 2005	2006 to 2008	2009 to 2011	% Change from 2003 to 2011	2011 Data		
					Num	Den	Percent
1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2)	67.7%	55.8%	67.9%	0.4%	49	78	62.8%
1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services.	66.3%	65.5%	65.0%	-2.0%	73	103	70.9%
1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services.	54.9%	54.1%	56.3%	2.6%	265	473	56.0%
2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy.	32.5%	35.2%	33.4%	2.8%	885	2,568	34.5%
2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum.	69.9%	72.4%	76.2%	9.0%	780	981	79.5%
3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1)	86.4%	100.0%	100.0%	15.8%			
3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1)	85.6%	82.6%	81.8%	-4.4%			
4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were:							
a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1)	64.2%	69.6%	64.9%	1.1%	101	159	63.5%
b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile)	18.3%	15.9%	16.8%	-8.1%	29	159	18.2%
c. Obese (BMI at or greater than the 95th percentile).	15.8%	12.5%	16.1%	2.2%	27	159	17.0%
d. Underweight (BMI less than the 5th percentile).	1.8%	2.1%	2.2%	22.5%	2	159	1.3%
4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3)							
a. Recommended prenatal weight gain	33.3%	23.9%	30.9%	-7.3%	12	66	18.2%
b. Excessive prenatal weight gain	40.4%	42.8%	49.1%	21.7%	40	66	60.6%
c. Inadequate prenatal weight gain	26.3%	33.3%	20.0%	-24.0%	14	66	21.2%
5A. Percentage of women who participated in WIC who initiated breastfeeding.	46.4%	54.4%	60.3%	30.0%	701	1,153	60.8%
5B. Percentage of infants who participated in WIC who were:							
a. Breastfeeding at 6 weeks of age (1)	28.1%	31.1%	30.0%	6.6%	338	1,102	30.7%
b. Breastfeeding at 6 months of age	14.7%	14.5%	16.2%	10.0%	190	1,102	17.2%

(1) Item appears also on other WCH Section Agreement Addenda.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI < 18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI ≥ 30.0): Adapted from the National Academy of Sciences. *Weight Gain During Pregnancy: Reexamining the Guidelines*. Washington, DC : National Academy Press, 2009.

SFY 2013-2014 AGREEMENT ADDENDA DATA

ATTACHMENT A: WIC QUALITY ASSURANCE DELIVERABLES

Rutherford

Process Outcome Objectives	2003 to 2005	2006 to 2008	2009 to 2011	% Change from 2003 to 2011	2011 Data		
					Num	Den	Percent
1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2)	67.0%	64.0%	80.8%	20.6%	380	475	80.0%
1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services.	79.2%	80.9%	76.9%	-3.0%	447	568	78.7%
1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services.	61.5%	60.0%	60.6%	-1.4%	1,474	2,330	63.3%
2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy.	32.5%	35.2%	33.4%	2.8%	885	2,568	34.5%
2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum.	69.9%	72.4%	76.2%	9.0%	780	981	79.5%
3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1)	92.3%	93.9%	100.0%	8.3%			
3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1)	93.4%	93.7%	94.7%	1.4%			
4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were:							
a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1)	68.2%	66.3%	65.1%	-4.5%	545	844	64.6%
b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile)	14.9%	15.6%	15.7%	5.1%	126	844	14.9%
c. Obese (BMI at or greater than the 95th percentile).	12.8%	13.7%	14.2%	10.9%	127	844	15.0%
d. Underweight (BMI less than the 5th percentile).	4.1%	4.4%	5.0%	21.9%	46	844	5.5%
4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3)							
a. Recommended prenatal weight gain	31.7%	30.4%	28.4%	-10.5%	126	489	25.8%
b. Excessive prenatal weight gain	39.5%	39.5%	48.1%	21.7%	247	489	50.5%
c. Inadequate prenatal weight gain	28.8%	30.1%	23.5%	-18.2%	116	489	23.7%
5A. Percentage of women who participated in WIC who initiated breastfeeding.	46.4%	54.4%	60.3%	30.0%	701	1,153	60.8%
5B. Percentage of infants who participated in WIC who were:							
a. Breastfeeding at 6 weeks of age (1)	28.1%	31.1%	30.0%	6.6%	338	1,102	30.7%
b. Breastfeeding at 6 months of age	14.7%	14.5%	16.2%	10.0%	190	1,102	17.2%

(1) Item appears also on other WCH Section Agreement Addenda.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI < 18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI ≥ 30.0): Adapted from the National Academy of Sciences. *Weight Gain During Pregnancy: Reexamining the Guidelines*. Washington, DC : National Academy Press, 2009.

SFY 2013-2014 AGREEMENT ADDENDA DATA

ATTACHMENT A: WIC QUALITY ASSURANCE DELIVERABLES

Tyrrell

Process Outcome Objectives	2003 to 2005	2006 to 2008	2009 to 2011	% Change from 2003 to 2011	2011 Data		
					Num	Den	Percent
1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2)	80.3%	83.7%	88.9%	10.7%	21	25	84.0%
1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services.	72.4%	72.7%	79.1%	9.2%	30	36	83.3%
1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services.	66.0%	69.1%	68.6%	3.9%	114	156	73.1%
2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy.	37.4%	32.3%	30.6%	-18.1%	242	911	26.6%
2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum.	79.1%	82.0%	79.7%	0.8%	247	313	78.9%
3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1)	97.5%	91.4%	97.6%	0.1%			
3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1)	95.0%	93.4%	94.3%	-0.7%			
4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were:							
a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1)	63.3%	62.6%	69.4%	9.6%	89	118	75.4%
b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile)	20.0%	14.1%	11.4%	-43.1%	12	118	10.2%
c. Obese (BMI at or greater than the 95th percentile).	14.9%	20.3%	14.6%	-2.0%	11	118	9.3%
d. Underweight (BMI less than the 5th percentile).	1.8%	3.1%	4.6%	157.2%	6	118	5.1%
4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3)							
a. Recommended prenatal weight gain	28.2%	33.3%	28.9%	2.6%	7	27	25.9%
b. Excessive prenatal weight gain	45.6%	47.0%	43.3%	-5.0%	12	27	44.4%
c. Inadequate prenatal weight gain	26.2%	19.7%	27.8%	6.0%	8	27	29.6%
5A. Percentage of women who participated in WIC who initiated breastfeeding.	41.3%	38.1%	26.6%	-35.5%	82	372	22.0%
5B. Percentage of infants who participated in WIC who were:							
a. Breastfeeding at 6 weeks of age (1)	24.6%	23.3%	14.8%	-39.9%	40	354	11.3%
b. Breastfeeding at 6 months of age	11.9%	11.4%	4.7%	-60.8%	15	354	4.2%

(1) Item appears also on other WCH Section Agreement Addenda.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI < 18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI ≥ 30.0): Adapted from the National Academy of Sciences. *Weight Gain During Pregnancy: Reexamining the Guidelines*. Washington, DC : National Academy Press, 2009.

SFY 2013-2014 AGREEMENT ADDENDA DATA

ATTACHMENT A: WIC QUALITY ASSURANCE DELIVERABLES

Vance

Process Outcome Objectives	2003 to 2005	2006 to 2008	2009 to 2011	% Change from 2003 to 2011	2011 Data		
					Num	Den	Percent
1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2)	69.0%	76.6%	71.6%	3.8%	275	389	70.7%
1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services.	78.0%	74.7%	76.0%	-2.6%	389	519	75.0%
1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services.	57.4%	55.9%	56.0%	-2.5%	1,380	2,346	58.8%
2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy.	32.6%	32.7%	29.0%	-11.1%	408	1,500	27.2%
2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum.	81.7%	77.8%	73.9%	-9.5%	558	735	75.9%
3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1)	90.0%	92.9%	100.0%	11.1%			
3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1)	92.8%	91.6%	89.6%	-3.4%			
4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were:							
a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1)	66.6%	68.3%	66.1%	-0.8%	503	757	66.4%
b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile)	14.5%	13.2%	11.6%	-20.5%	89	757	11.8%
c. Obese (BMI at or greater than the 95th percentile).	15.0%	12.0%	11.6%	-22.7%	89	757	11.8%
d. Underweight (BMI less than the 5th percentile).	3.9%	6.4%	10.8%	178.4%	76	757	10.0%
4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3)							
a. Recommended prenatal weight gain	31.5%	29.8%	32.0%	1.6%	138	412	33.5%
b. Excessive prenatal weight gain	43.2%	42.0%	37.9%	-12.1%	153	412	37.1%
c. Inadequate prenatal weight gain	25.3%	28.1%	30.1%	18.7%	121	412	29.4%
5A. Percentage of women who participated in WIC who initiated breastfeeding.	43.3%	52.9%	41.4%	-4.3%	408	922	44.3%
5B. Percentage of infants who participated in WIC who were:							
a. Breastfeeding at 6 weeks of age (1)	28.7%	34.5%	36.3%	26.6%	320	959	33.4%
b. Breastfeeding at 6 months of age	20.9%	22.5%	26.8%	28.5%	238	959	24.8%

(1) Item appears also on other WCH Section Agreement Addenda.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI < 18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI ≥ 30.0): Adapted from the National Academy of Sciences. *Weight Gain During Pregnancy: Reexamining the Guidelines*. Washington, DC : National Academy Press, 2009.

SFY 2013-2014 AGREEMENT ADDENDA DATA

ATTACHMENT A: WIC QUALITY ASSURANCE DELIVERABLES

Washington

Process Outcome Objectives	2003 to 2005	2006 to 2008	2009 to 2011	% Change from 2003 to 2011	2011 Data		
					Num	Den	Percent
1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2)	86.6%	86.4%	85.5%	-1.2%	94	111	84.7%
1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services.	75.1%	73.6%	80.0%	6.7%	112	134	83.6%
1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services.	69.6%	67.9%	61.6%	-11.5%	371	592	62.7%
2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy.	37.4%	32.3%	30.6%	-18.1%	242	911	26.6%
2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum.	79.1%	82.0%	79.7%	0.8%	247	313	78.9%
3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1)	98.9%	98.0%	100.0%	1.1%			
3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1)	95.8%	93.6%	94.7%	-1.1%			
4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were:							
a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1)	68.0%	68.1%	66.9%	-1.7%	139	223	62.3%
b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile)	10.4%	9.4%	10.4%	0.3%	31	223	13.9%
c. Obese (BMI at or greater than the 95th percentile).	10.1%	9.8%	12.7%	26.1%	36	223	16.1%
d. Underweight (BMI less than the 5th percentile).	11.5%	12.7%	10.0%	-13.2%	17	223	7.6%
4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3)							
a. Recommended prenatal weight gain	26.8%	26.7%	30.6%	14.2%	34	111	30.6%
b. Excessive prenatal weight gain	43.8%	45.1%	42.9%	-2.1%	46	111	41.4%
c. Inadequate prenatal weight gain	29.3%	28.2%	26.4%	-9.9%	31	111	27.9%
5A. Percentage of women who participated in WIC who initiated breastfeeding.	41.3%	38.1%	26.6%	-35.5%	82	372	22.0%
5B. Percentage of infants who participated in WIC who were:							
a. Breastfeeding at 6 weeks of age (1)	24.6%	23.3%	14.8%	-39.9%	40	354	11.3%
b. Breastfeeding at 6 months of age	11.9%	11.4%	4.7%	-60.8%	15	354	4.2%

(1) Item appears also on other WCH Section Agreement Addenda.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI < 18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI ≥ 30.0): Adapted from the National Academy of Sciences. *Weight Gain During Pregnancy: Reexamining the Guidelines*. Washington, DC : National Academy Press, 2009.

SFY 2013-2014 AGREEMENT ADDENDA DATA

ATTACHMENT A: WIC QUALITY ASSURANCE DELIVERABLES

Watauga

Process Outcome Objectives	2003 to 2005	2006 to 2008	2009 to 2011	% Change from 2003 to 2011	2011 Data		
					Num	Den	Percent
1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2)	79.8%	50.9%	63.1%	-21.0%	108	161	67.1%
1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services.	60.6%	66.6%	59.9%	-1.2%	120	209	57.4%
1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services.	65.3%	60.3%	62.2%	-4.8%	526	832	63.2%
2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy.	52.1%	45.2%	39.3%	-24.7%	177	474	37.3%
2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum.	58.2%	66.4%	69.2%	18.9%	136	198	68.7%
3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1)	71.7%	82.4%					
3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1)	80.6%	83.1%					
4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were:							
a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1)	66.2%	73.3%	67.1%	1.3%	221	332	66.6%
b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile)	17.6%	13.0%	17.4%	-0.8%	62	332	18.7%
c. Obese (BMI at or greater than the 95th percentile).	12.2%	10.7%	11.8%	-2.9%	41	332	12.3%
d. Underweight (BMI less than the 5th percentile).	4.0%	3.0%	3.7%	-9.1%	8	332	2.4%
4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3)							
a. Recommended prenatal weight gain	31.6%	31.8%	33.0%	4.4%	56	179	31.3%
b. Excessive prenatal weight gain	49.1%	51.6%	50.2%	2.3%	97	179	54.2%
c. Inadequate prenatal weight gain	19.3%	16.5%	16.7%	-13.0%	26	179	14.5%
5A. Percentage of women who participated in WIC who initiated breastfeeding.	53.4%	63.1%	71.4%	33.8%	185	250	74.0%
5B. Percentage of infants who participated in WIC who were:							
a. Breastfeeding at 6 weeks of age (1)	42.7%	46.5%	49.8%	16.7%	130	242	53.7%
b. Breastfeeding at 6 months of age	27.7%	26.9%	29.2%	5.4%	74	242	30.6%

(1) Item appears also on other WCH Section Agreement Addenda.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI < 18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI ≥ 30.0): Adapted from the National Academy of Sciences. *Weight Gain During Pregnancy: Reexamining the Guidelines*. Washington, DC : National Academy Press, 2009.

SFY 2013-2014 AGREEMENT ADDENDA DATA

ATTACHMENT A: WIC QUALITY ASSURANCE DELIVERABLES

Yancey

Process Outcome Objectives	2003 to 2005	2006 to 2008	2009 to 2011	% Change from 2003 to 2011	2011 Data		
					Num	Den	Percent
1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2)	68.5%	68.7%	68.6%	0.1%	64	107	59.8%
1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services.	62.1%	62.0%	64.1%	3.3%	83	127	65.4%
1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services.	62.3%	58.8%	63.7%	2.4%	337	502	67.1%
2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy.	40.0%	42.5%	37.7%	-5.8%	130	370	35.1%
2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum.	69.1%	63.9%	67.8%	-1.9%	108	151	71.5%
3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1)	85.2%	100.0%	100.0%	17.3%			
3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1)	92.1%	86.1%	83.9%	-8.9%			
4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were:							
a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1)	62.5%	70.3%	62.3%	-0.3%	96	157	61.1%
b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile)	20.0%	15.3%	18.4%	-8.2%	29	157	18.5%
c. Obese (BMI at or greater than the 95th percentile).	15.2%	11.0%	16.2%	6.8%	28	157	17.8%
d. Underweight (BMI less than the 5th percentile).	2.3%	3.5%	3.1%	33.4%	4	157	2.5%
4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3)							
a. Recommended prenatal weight gain	32.1%	32.9%	32.9%	2.7%	29	100	29.0%
b. Excessive prenatal weight gain	42.5%	46.3%	39.2%	-7.8%	39	100	39.0%
c. Inadequate prenatal weight gain	25.4%	20.8%	27.8%	9.7%	32	100	32.0%
5A. Percentage of women who participated in WIC who initiated breastfeeding.	79.2%	68.1%	69.3%	-12.5%	99	174	56.9%
5B. Percentage of infants who participated in WIC who were:							
a. Breastfeeding at 6 weeks of age (1)	39.9%	34.8%	40.9%	2.4%	55	143	38.5%
b. Breastfeeding at 6 months of age	25.7%	20.2%	30.0%	16.9%	42	143	29.4%

(1) Item appears also on other WCH Section Agreement Addenda.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI < 18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI ≥ 30.0): Adapted from the National Academy of Sciences. *Weight Gain During Pregnancy: Reexamining the Guidelines*. Washington, DC : National Academy Press, 2009.