

Trend Data for the WIC Agreement Addenda SFY 2014-2015

The overall purpose of trend data is to track patterns or changes in data over time. For purposes of the WIC Agreement Addenda, trend data is being used to track the performance of the State and Local WIC Programs with respect to fifteen identified program indicators and sub-indicators. The goal for both the State and Local WIC Programs is to implement or enhance activities which improve performance over time for each indicator.

This compendium consists of two reports of WIC Program Quality Assurance Deliverables data. One report is WIC Local Agency specific and the other represents North Carolina state wide data. It may be useful to compare an agency's performance with respect to the state as a whole. To compare local agency data with other agencies' data, please refer to the Indicator-specific reports also available on the website <http://www.nutritionnc.com/wic/wicLAR.htm>.

Each report includes the following information.

❖ **Process Outcome Objectives**

This column includes a description of the Process Outcome Objectives (POO) data indicators of the WIC Program Quality Assurance Deliverable being assessed.

❖ **Trend Data**

The data displayed in these columns represent nine years of data divided into 3-year time periods (2004 to 2006, 2007 to 2009, and 2010 to 2012*) with the average of each three year time period shown.

- POO 1A, 1B, 1C, 4A, and 4B are calendar year (Jan – Dec) data.
- POO 2A, 2B, and 5A, are fiscal year (July – June) data, six months more recent than calendar year data.
- POO 5B is fiscal year (July – June) data which is twelve months prior to other fiscal year data.
- **POO 3A and 3B (Child Health data):** The trend and 2012 data used for Child Health for POO indicators (3A and 3B) are not presented because of inconsistencies in data collection systems over time.

❖ **% Change from 2004-2012**

This column displays the percent change from the first three year period to the most recent three-year time period along with the direction of the change (i.e., increase, decrease or no change).

❖ **2012 Data**

This column displays the most recent year's data including the numerator (Num) and denominator (Den) counts and the percent. Percentages calculated based on numbers less than 20 can show wide fluctuations from one year to the next. Therefore numbers below 20 in the denominator should be used with caution as the computed rates using small numbers are very unreliable.

Guidance for Completing the WIC Agreement Addendum & Local Agency Nutrition Education Plan

The consolidated planning process allows all budgets and agreement addenda for local public health programs receiving funding from the State to be submitted at one time in a consolidated manner. The WIC agreement addendum also serves as the federally-required Local Agency Nutrition Education Plan.

Completing the WIC Agreement Addendum

Section III: Scope of Work and Deliverables

- Deliverables #1 - 5 - These five deliverables do not require staff to complete or to write anything. Instead, local WIC agencies are agreeing to improve the percentage for each of the listed indicators. To assess performance levels and to determine actions for improving performance, staff should refer to the “Trend Data” under Agreement Addenda Data for SCY 2014-2015 accessible through the web link: <http://nutritionnc.com/wic/wicLAR.htm> after February 14th, 2014..

NOTE: The table below provides information related to Deliverable #4. Increase the percent of women with live term singleton births who receive WIC Program Services prenatally and who gain weight within the –2009 guidelines from the Institute of Medicine (IOM) (as adapted from National Academy of Sciences. Weight Gain During Pregnancy: Reexamining the Guidelines. Washington, DC : National Academy Press, 2009) This information is also on the “Trend Data Reports” associated with the WIC Agreement Addendum.

| <i>Woman's Prepregnancy BMI Category</i> | <i>Woman's Prepregnancy Weight</i> | <i>Recommended Range of Total Weight Gain in Pregnancy</i> |
|--|------------------------------------|--|
| Below 18.5 | Underweight | 28 to 40 Pounds |
| 18.5 - 24.9 | Normal Weight | 25 to 35 pounds |
| 25.0 - 29.9 | Overweight | 15 to 25 Pounds |
| 30.0 and Above | Obese | 11 to 20 Pounds |

- Deliverable #6 - Does not require staff to write anything, however, the agency agrees to comply with this requirement throughout the year.
- Deliverables #7 and 8 - These two deliverables relate to the Local Agency Nutrition Education Plan.
 - Deliverable #7 requires staff to attach to the WIC Agreement Addendum, a Nutrition Education Plan for SFY 2014-15 for at least one of the focus areas listed, as defined in the Agreement Addendum.
 - Deliverable #8 requires staff to attach to the WIC Agreement Addendum, a brief evaluation of the local agency’s 2013-14 plan as defined in the Agreement Addendum.

Section IV: Performance Measures/Reporting Requirements

Local agencies are agreeing to maintain an average monthly participation of at least 97% of their assigned monthly base caseload. The minimum average monthly participation for the SFY can be determined by multiplying the assigned base caseload by 97%.

Assuring the WIC Agreement Addendum and Budget are Complete

Prior to submission, local staff may wish to use the checklist below to ensure that their WIC agreement addendum and budget are complete.

Agreement Addendum

- The Health Director has signed and dated the first page.
- Contact information for the Local Health Department is placed on the first page.
- The evaluation of the 2013-14 Agreement Addendum's Nutrition Education Plan (Deliverable #8) has been included as an attachment to the Agreement Addendum.
- The Nutrition Education Plan for 2014-15 (Deliverable #7) has been included as an Attachment to the Agreement Addendum.

Budget

- Figures are added correctly and rounded to the nearest whole dollar (no cents!).
- At least 20% of the total WIC allocation is budgeted in Nutrition Education (5404).
Note: The minimum amount for Nutrition Education is listed on your budget page.
- No more than 10% of the total WIC allocation is budgeted in General Administration (5405).
- The amount budgeted for Breastfeeding (5409) equals or exceeds the minimum amount for Breastfeeding listed on your budget page.
- Total budget does not exceed assigned WIC allocation.
- Submitted budget is the original document (not a photocopy).

Resources for Nutrition Education Activities

The following list of websites offer ideas that staff may consider as they develop their nutrition education plans. This list of websites is by no means all inclusive.

<http://www.cdc.gov/physicalactivity/> (ideas for physical activity)

<http://www.surgeongeneral.gov/library/calls/breastfeeding/index.html> (ideas for breastfeeding education and support)

<http://www.fruitsandveggiesmorematters.org/> (ideas for eating more fruits & vegetables)

www.cspinet.org/nutrition/schoolkit.html or <http://www.nutritionnc.com/moveMilk.htm> (ideas for promoting consumption of 1% or less milk)

http://www.nal.usda.gov/wicworks/Learning_Center/loving_support.html (overall good website for ideas and resources for nutrition & breastfeeding education activities in the WIC Program)

<http://www.nchealthyweight.com/> (ideas for moving our children towards a healthy weight)

<http://www.eatsmartmovemorenc.com/> and <http://www.myeatsmartmovemore.com> (ideas for physical activity and nutrition activities)

<http://www.colormehealthy.com/> (ideas for getting preschoolers to move more and eat healthy)

<http://www.nutritionnc.com/> (ideas for a variety of nutrition activities & information on how to borrow materials from the Nutrition Education and Training (NET) Library)

SFY 2014-2015 AGREEMENT ADDENDA DATA
WIC QUALITY ASSURANCE DELIVERABLES
North Carolina

| Process Outcome Objectives | 2004 to 2006 | 2007 to 2009 | 2010 to 2012 | % Change from 2004 to 2012 | 2012 Data | | |
|---|--------------------|--------------------|--------------------|----------------------------------|-----------|---------|---------|
| | | | | | Num | Den | Percent |
| 1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2) | 71.0% | 71.4% | 71.5% | 0.8% | 41,970 | 56,771 | 73.9% |
| 1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services. | 70.8% | 71.7% | 71.1% | 0.4% | 52,409 | 73,811 | 71.0% |
| 1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services. | 53.0% | 54.5% | 55.9% | 5.4% | 216,826 | 381,405 | 56.8% |
| 2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy. | 30.1% | 29.6% | 29.0% | -3.5% | 39,045 | 138,792 | 28.1% |
| 2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum. | 67.4% | 68.9% | 72.8% | 8.1% | 48,247 | 64,257 | 75.1% |
| 3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1) | | | | | | | |
| 3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1) | | | | | | | |
| 4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were: | | | | | | | |
| a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1) | 65.4% | 65.1% | 63.8% | -2.4% | 71,801 | 112,204 | 64.0% |
| b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile) | 15.6% | 15.9% | 15.7% | 1.0% | 16,625 | 112,204 | 14.8% |
| c. Obese (BMI at or greater than the 95th percentile). | 14.8% | 15.3% | 15.3% | 3.0% | 16,442 | 112,204 | 14.7% |
| d. Underweight (BMI less than the 5th percentile). | 4.2% | 3.6% | 5.2% | 23.7% | 7,336 | 112,204 | 6.5% |
| 4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3) | | | | | | | |
| a. Recommended prenatal weight gain | 31.7% | 31.8% | 28.8% | -9.1% | 14,155 | 51,590 | 27.4% |
| b. Excessive prenatal weight gain | 46.3% | 45.7% | 47.7% | 3.1% | 25,057 | 51,590 | 48.6% |
| c. Inadequate prenatal weight gain | 22.0% | 22.5% | 23.5% | 6.7% | 12,378 | 51,590 | 24.0% |
| 5A. Percentage of women who participated in WIC who initiated breastfeeding. | 51.1% | 54.7% | 57.2% | 11.9% | 47,427 | 80,977 | 58.6% |
| 5B. Percentage of infants who participated in WIC who were: | | | | | | | |
| a. Breastfeeding at 6 weeks of age (1) | 29.5% | 32.4% | 34.0% | 15.0% | 29,464 | 81,644 | 36.1% |
| b. Breastfeeding at 6 months of age | 15.3% | 17.1% | 18.7% | 22.1% | 16,316 | 81,644 | 20.0% |

(1) Item appears also on other WCH Section Agreement Addenda. Data for 3A and 3B are not presented due to limitation in recent trend data.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI >=30.0): Adapted from the National Academy of Sciences. Weight Gain During Pregnancy: Reexamining the Guidelines. Washington, DC : National Academy Press, 2009.

SFY 2014-2015 AGREEMENT ADDENDA DATA

WIC QUALITY ASSURANCE DELIVERABLES

Alamance County

| Process Outcome Objectives | 2004 to 2006 | 2007 to 2009 | 2010 to 2012 | % Change from 2004 to 2012 | 2012 Data | | |
|---|--------------------|--------------------|--------------------|----------------------------------|-----------|-------|---------|
| | | | | | Num | Den | Percent |
| 1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2) | 75.5% | 77.6% | 75.8% | 0.4% | 724 | 906 | 79.9% |
| 1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services. | 74.7% | 75.8% | 72.0% | -3.7% | 840 | 1,201 | 69.9% |
| 1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services. | 51.8% | 56.5% | 58.3% | 12.5% | 3,641 | 6,207 | 58.7% |
| 2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy. | 40.9% | 42.8% | 35.0% | -14.4% | 716 | 2,207 | 32.4% |
| 2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum. | 46.0% | 72.3% | 77.6% | 68.6% | 849 | 1,103 | 77.0% |
| 3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1) | | | | | | | |
| 3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1) | | | | | | | |
| 4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were: | | | | | | | |
| a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1) | 62.5% | 65.4% | 61.9% | -1.0% | 1,163 | 1,873 | 62.1% |
| b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile) | 19.0% | 15.9% | 18.1% | -4.5% | 315 | 1,873 | 16.8% |
| c. Obese (BMI at or greater than the 95th percentile). | 16.7% | 16.5% | 16.8% | 0.2% | 328 | 1,873 | 17.5% |
| d. Underweight (BMI less than the 5th percentile). | 1.8% | 2.3% | 3.2% | 82.8% | 67 | 1,873 | 3.6% |
| 4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3) | | | | | | | |
| a. Recommended prenatal weight gain | 33.5% | 32.9% | 29.2% | -12.9% | 209 | 845 | 24.7% |
| b. Excessive prenatal weight gain | 47.3% | 44.9% | 46.9% | -0.8% | 431 | 845 | 51.0% |
| c. Inadequate prenatal weight gain | 19.2% | 22.2% | 23.9% | 24.6% | 205 | 845 | 24.3% |
| 5A. Percentage of women who participated in WIC who initiated breastfeeding. | 47.3% | 50.1% | 61.8% | 30.7% | 922 | 1,325 | 69.6% |
| 5B. Percentage of infants who participated in WIC who were: | | | | | | | |
| a. Breastfeeding at 6 weeks of age (1) | 32.1% | 32.1% | 36.1% | 12.4% | 458 | 1,239 | 37.0% |
| b. Breastfeeding at 6 months of age | 16.9% | 18.8% | 19.7% | 16.7% | 244 | 1,239 | 19.7% |

(1) Item appears also on other WCH Section Agreement Addenda. Data for 3A and 3B were not presented due to the limitation in recent trend datasets.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI >=30.0): Adapted from the National Academy of Sciences. Weight Gain During Pregnancy: Reexamining the Guidelines. Washington, DC : National Academy Press, 2009.

SFY 2014-2015 AGREEMENT ADDENDA DATA

WIC QUALITY ASSURANCE DELIVERABLES

Albemarle Regional Health Services District

| Process Outcome Objectives | 2004 to 2006 | 2007 to 2009 | 2010 to 2012 | % Change from 2004 to 2012 | 2012 Data | | |
|---|--------------------|--------------------|--------------------|----------------------------------|-----------|-------|---------|
| | | | | | Num | Den | Percent |
| 1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2) | 78.3% | 71.3% | 68.3% | -12.8% | 486 | 703 | 69.1% |
| 1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services. | 73.9% | 73.2% | 74.6% | 1.0% | 643 | 843 | 76.3% |
| 1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services. | 58.4% | 57.4% | 57.6% | -1.3% | 2,784 | 4,667 | 59.7% |
| 2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy. | 36.1% | 32.8% | 26.8% | -25.7% | 591 | 2,230 | 26.5% |
| 2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum. | 33.5% | 40.4% | 56.7% | 69.1% | 654 | 894 | 73.2% |
| 3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1) | | | | | | | |
| 3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1) | | | | | | | |
| 4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were: | | | | | | | |
| a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1) | 65.2% | 65.6% | 61.7% | -5.4% | 2,182 | 3,579 | 61.0% |
| b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile) | 14.8% | 14.6% | 13.6% | -8.1% | 453 | 3,579 | 12.7% |
| c. Obese (BMI at or greater than the 95th percentile). | 15.0% | 15.9% | 16.3% | 9.2% | 600 | 3,579 | 16.8% |
| d. Underweight (BMI less than the 5th percentile). | 5.0% | 3.8% | 8.4% | 66.0% | 344 | 3,579 | 9.6% |
| 4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3) | | | | | | | |
| a. Recommended prenatal weight gain | 32.0% | 32.0% | 26.7% | -16.7% | 150 | 581 | 25.8% |
| b. Excessive prenatal weight gain | 44.2% | 44.0% | 39.8% | -9.9% | 232 | 581 | 39.9% |
| c. Inadequate prenatal weight gain | 23.8% | 24.0% | 33.5% | 40.8% | 199 | 581 | 34.3% |
| 5A. Percentage of women who participated in WIC who initiated breastfeeding. | 20.5% | 22.0% | 32.9% | 60.8% | 388 | 1,111 | 34.9% |
| 5B. Percentage of infants who participated in WIC who were: | | | | | | | |
| a. Breastfeeding at 6 weeks of age (1) | 8.7% | 10.1% | 15.0% | 73.9% | 196 | 992 | 19.8% |
| b. Breastfeeding at 6 months of age | 3.3% | 3.1% | 7.2% | 119.5% | 90 | 992 | 9.1% |

(1) Item appears also on other WCH Section Agreement Addenda. Data for 3A and 3B were not presented due to the limitation in recent trend datasets.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI >=30.0): Adapted from the National Academy of Sciences. Weight Gain During Pregnancy: Reexamining the Guidelines. Washington, DC : National Academy Press, 2009.

SFY 2014-2015 AGREEMENT ADDENDA DATA

WIC QUALITY ASSURANCE DELIVERABLES

Alexander County

| Process Outcome Objectives | 2004 to 2006 | 2007 to 2009 | 2010 to 2012 | % Change from 2004 to 2012 | 2012 Data | | |
|---|--------------------|--------------------|--------------------|----------------------------------|-----------|-------|---------|
| | | | | | Num | Den | Percent |
| 1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2) | 73.1% | 73.3% | 73.2% | 0.1% | 164 | 237 | 69.2% |
| 1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services. | 78.5% | 77.9% | 73.8% | -6.1% | 184 | 249 | 73.9% |
| 1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services. | 62.9% | 57.2% | 58.3% | -7.4% | 728 | 1,202 | 60.6% |
| 2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy. | 41.3% | 36.6% | 37.1% | -10.3% | 127 | 372 | 34.1% |
| 2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum. | 51.0% | 76.6% | 79.9% | 56.7% | 164 | 202 | 81.2% |
| 3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1) | | | | | | | |
| 3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1) | | | | | | | |
| 4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were: | | | | | | | |
| a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1) | 63.6% | 67.3% | 67.0% | 5.4% | 154 | 235 | 65.5% |
| b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile) | 16.0% | 15.1% | 15.4% | -3.6% | 39 | 235 | 16.6% |
| c. Obese (BMI at or greater than the 95th percentile). | 16.0% | 13.9% | 12.4% | -22.4% | 30 | 235 | 12.8% |
| d. Underweight (BMI less than the 5th percentile). | 4.3% | 3.7% | 5.1% | 17.4% | 12 | 235 | 5.1% |
| 4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3) | | | | | | | |
| a. Recommended prenatal weight gain | 28.7% | 30.0% | 30.4% | 6.1% | 54 | 212 | 25.5% |
| b. Excessive prenatal weight gain | 46.5% | 49.6% | 47.5% | 2.2% | 103 | 212 | 48.6% |
| c. Inadequate prenatal weight gain | 24.9% | 20.4% | 22.1% | -11.1% | 55 | 212 | 25.9% |
| 5A. Percentage of women who participated in WIC who initiated breastfeeding. | 37.6% | 57.4% | 89.2% | 137.5% | 243 | 267 | 91.0% |
| 5B. Percentage of infants who participated in WIC who were: | | | | | | | |
| a. Breastfeeding at 6 weeks of age (1) | 25.7% | 27.2% | 32.4% | 26.1% | 96 | 287 | 33.4% |
| b. Breastfeeding at 6 months of age | 16.0% | 14.9% | 21.0% | 31.0% | 67 | 287 | 23.3% |

(1) Item appears also on other WCH Section Agreement Addenda. Data for 3A and 3B were not presented due to the limitation in recent trend datasets.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI >=30.0): Adapted from the National Academy of Sciences. Weight Gain During Pregnancy: Reexamining the Guidelines. Washington, DC : National Academy Press, 2009.

SFY 2014-2015 AGREEMENT ADDENDA DATA

WIC QUALITY ASSURANCE DELIVERABLES

Anson County

| Process Outcome Objectives | 2004 to 2006 | 2007 to 2009 | 2010 to 2012 | % Change from 2004 to 2012 | 2012 Data | | |
|---|--------------------|--------------------|--------------------|----------------------------------|-----------|-------|---------|
| | | | | | Num | Den | Percent |
| 1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2) | 83.0% | 83.1% | 84.3% | 1.5% | 166 | 195 | 85.1% |
| 1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services. | 74.0% | 79.8% | 75.8% | 2.4% | 164 | 213 | 77.0% |
| 1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services. | 60.9% | 61.2% | 61.1% | 0.3% | 675 | 1,067 | 63.3% |
| 2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy. | 50.7% | 50.3% | 43.0% | -15.2% | 181 | 432 | 41.9% |
| 2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum. | 80.4% | 81.5% | 82.6% | 2.7% | 181 | 222 | 81.5% |
| 3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1) | | | | | | | |
| 3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1) | | | | | | | |
| 4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were: | | | | | | | |
| a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1) | 66.0% | 67.6% | 57.7% | -12.6% | 245 | 416 | 58.9% |
| b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile) | 12.1% | 10.0% | 18.0% | 48.6% | 76 | 416 | 18.3% |
| c. Obese (BMI at or greater than the 95th percentile). | 14.2% | 12.0% | 20.0% | 41.2% | 80 | 416 | 19.2% |
| d. Underweight (BMI less than the 5th percentile). | 7.7% | 10.4% | 4.3% | -44.0% | 15 | 416 | 3.6% |
| 4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3) | | | | | | | |
| a. Recommended prenatal weight gain | 30.7% | 26.3% | 25.1% | -18.4% | 48 | 195 | 24.6% |
| b. Excessive prenatal weight gain | 47.4% | 47.7% | 50.5% | 6.4% | 97 | 195 | 49.7% |
| c. Inadequate prenatal weight gain | 21.8% | 26.0% | 24.4% | 11.9% | 50 | 195 | 25.6% |
| 5A. Percentage of women who participated in WIC who initiated breastfeeding. | 31.9% | 36.2% | 37.4% | 17.2% | 90 | 271 | 33.2% |
| 5B. Percentage of infants who participated in WIC who were: | | | | | | | |
| a. Breastfeeding at 6 weeks of age (1) | 18.0% | 17.0% | 15.0% | -16.9% | 37 | 246 | 15.0% |
| b. Breastfeeding at 6 months of age | 6.2% | 4.8% | 6.0% | -2.2% | 12 | 246 | 4.9% |

(1) Item appears also on other WCH Section Agreement Addenda. Data for 3A and 3B were not presented due to the limitation in recent trend datasets.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI >=30.0): Adapted from the National Academy of Sciences. *Weight Gain During Pregnancy: Reexamining the Guidelines*. Washington, DC : National Academy Press, 2009.

SFY 2014-2015 AGREEMENT ADDENDA DATA

WIC QUALITY ASSURANCE DELIVERABLES

Appalachian District

| Process Outcome Objectives | 2004 to 2006 | 2007 to 2009 | 2010 to 2012 | % Change from 2004 to 2012 | 2012 Data | | |
|---|--------------------|--------------------|--------------------|----------------------------------|-----------|-------|---------|
| | | | | | Num | Den | Percent |
| 1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2) | 74.0% | 64.4% | 73.6% | -0.5% | 304 | 382 | 79.6% |
| 1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services. | 69.5% | 70.8% | 66.3% | -4.5% | 317 | 473 | 67.0% |
| 1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services. | 64.7% | 63.9% | 62.5% | -3.4% | 1,488 | 2,352 | 63.3% |
| 2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy. | 49.9% | 46.2% | 38.9% | -22.1% | 344 | 930 | 37.0% |
| 2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum. | 68.8% | 74.3% | 73.3% | 6.5% | 309 | 409 | 75.6% |
| 3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1) | | | | | | | |
| 3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1) | | | | | | | |
| 4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were: | | | | | | | |
| a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1) | 66.9% | 66.2% | 63.3% | -5.4% | 544 | 859 | 63.3% |
| b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile) | 14.2% | 15.7% | 16.6% | 17.3% | 125 | 859 | 14.6% |
| c. Obese (BMI at or greater than the 95th percentile). | 14.6% | 14.4% | 15.5% | 5.6% | 137 | 859 | 15.9% |
| d. Underweight (BMI less than the 5th percentile). | 4.3% | 3.6% | 4.6% | 8.1% | 53 | 859 | 6.2% |
| 4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3) | | | | | | | |
| a. Recommended prenatal weight gain | 31.6% | 30.0% | 31.9% | 1.1% | 99 | 335 | 29.6% |
| b. Excessive prenatal weight gain | 48.0% | 49.0% | 49.4% | 3.1% | 176 | 335 | 52.5% |
| c. Inadequate prenatal weight gain | 20.5% | 21.1% | 18.6% | -8.9% | 60 | 335 | 17.9% |
| 5A. Percentage of women who participated in WIC who initiated breastfeeding. | 54.2% | 58.8% | 60.1% | 11.0% | 283 | 503 | 56.3% |
| 5B. Percentage of infants who participated in WIC who were: | | | | | | | |
| a. Breastfeeding at 6 weeks of age (1) | 34.4% | 36.4% | 38.2% | 11.0% | 203 | 493 | 41.2% |
| b. Breastfeeding at 6 months of age | 18.4% | 20.5% | 24.6% | 33.3% | 149 | 493 | 30.2% |

(1) Item appears also on other WCH Section Agreement Addenda. Data for 3A and 3B were not presented due to the limitation in recent trend datasets.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI >=30.0): Adapted from the National Academy of Sciences. Weight Gain During Pregnancy: Reexamining the Guidelines. Washington, DC : National Academy Press, 2009.

SFY 2014-2015 AGREEMENT ADDENDA DATA

WIC QUALITY ASSURANCE DELIVERABLES

Beaufort County

| Process Outcome Objectives | 2004 to 2006 | 2007 to 2009 | 2010 to 2012 | % Change from 2004 to 2012 | 2012 Data | | |
|---|--------------------|--------------------|--------------------|----------------------------------|-----------|-------|---------|
| | | | | | Num | Den | Percent |
| 1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2) | 83.6% | 78.8% | 82.6% | -1.2% | 267 | 311 | 85.9% |
| 1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services. | 76.5% | 76.7% | 70.8% | -7.4% | 265 | 371 | 71.4% |
| 1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services. | 58.6% | 58.7% | 62.6% | 6.7% | 1,165 | 1,851 | 62.9% |
| 2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy. | 36.4% | 38.5% | 34.5% | -5.5% | 355 | 1,030 | 34.5% |
| 2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum. | 73.6% | 73.6% | 79.5% | 8.0% | 340 | 417 | 81.5% |
| 3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1) | | | | | | | |
| 3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1) | | | | | | | |
| 4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were: | | | | | | | |
| a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1) | 63.8% | 66.2% | 64.6% | 1.2% | 523 | 787 | 66.5% |
| b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile) | 15.8% | 16.0% | 15.7% | -0.9% | 113 | 787 | 14.4% |
| c. Obese (BMI at or greater than the 95th percentile). | 17.8% | 14.9% | 15.8% | -11.1% | 117 | 787 | 14.9% |
| d. Underweight (BMI less than the 5th percentile). | 2.6% | 2.9% | 3.9% | 53.8% | 34 | 787 | 4.3% |
| 4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3) | | | | | | | |
| a. Recommended prenatal weight gain | 31.6% | 35.6% | 28.9% | -8.6% | 77 | 278 | 27.7% |
| b. Excessive prenatal weight gain | 45.8% | 43.8% | 43.2% | -5.6% | 115 | 278 | 41.4% |
| c. Inadequate prenatal weight gain | 22.6% | 20.6% | 27.9% | 23.5% | 86 | 278 | 30.9% |
| 5A. Percentage of women who participated in WIC who initiated breastfeeding. | 44.2% | 43.5% | 50.1% | 13.4% | 251 | 485 | 51.8% |
| 5B. Percentage of infants who participated in WIC who were: | | | | | | | |
| a. Breastfeeding at 6 weeks of age (1) | 24.2% | 27.3% | 31.6% | 30.7% | 141 | 440 | 32.0% |
| b. Breastfeeding at 6 months of age | 12.3% | 12.9% | 15.6% | 27.3% | 75 | 440 | 17.0% |

(1) Item appears also on other WCH Section Agreement Addenda. Data for 3A and 3B were not presented due to the limitation in recent trend datasets.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI >=30.0): Adapted from the National Academy of Sciences. Weight Gain During Pregnancy: Reexamining the Guidelines. Washington, DC : National Academy Press, 2009.

SFY 2014-2015 AGREEMENT ADDENDA DATA

WIC QUALITY ASSURANCE DELIVERABLES

Bladen County

| Process Outcome Objectives | 2004 to 2006 | 2007 to 2009 | 2010 to 2012 | % Change from 2004 to 2012 | 2012 Data | | |
|---|--------------------|--------------------|--------------------|----------------------------------|-----------|-------|---------|
| | | | | | Num | Den | Percent |
| 1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2) | 82.2% | 83.1% | 83.7% | 1.8% | 211 | 250 | 84.4% |
| 1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services. | 82.9% | 81.8% | 79.8% | -3.7% | 246 | 307 | 80.1% |
| 1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services. | 62.5% | 65.1% | 65.1% | 4.2% | 942 | 1,447 | 65.1% |
| 2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy. | 43.4% | 46.4% | 35.0% | -19.4% | 276 | 832 | 33.2% |
| 2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum. | 74.3% | 73.5% | 74.6% | 0.4% | 232 | 293 | 79.2% |
| 3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1) | | | | | | | |
| 3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1) | | | | | | | |
| 4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were: | | | | | | | |
| a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1) | 65.5% | 63.2% | 61.8% | -5.6% | 418 | 692 | 60.4% |
| b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile) | 14.5% | 16.6% | 16.2% | 12.2% | 103 | 692 | 14.9% |
| c. Obese (BMI at or greater than the 95th percentile). | 14.3% | 16.3% | 16.4% | 14.8% | 110 | 692 | 15.9% |
| d. Underweight (BMI less than the 5th percentile). | 5.8% | 4.0% | 5.6% | -3.7% | 61 | 692 | 8.8% |
| 4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3) | | | | | | | |
| a. Recommended prenatal weight gain | 29.0% | 28.4% | 26.3% | -9.6% | 63 | 226 | 27.9% |
| b. Excessive prenatal weight gain | 45.5% | 43.7% | 48.8% | 7.1% | 111 | 226 | 49.1% |
| c. Inadequate prenatal weight gain | 25.4% | 28.0% | 25.0% | -1.8% | 52 | 226 | 23.0% |
| 5A. Percentage of women who participated in WIC who initiated breastfeeding. | 50.8% | 58.5% | 53.1% | 4.5% | 204 | 366 | 55.7% |
| 5B. Percentage of infants who participated in WIC who were: | | | | | | | |
| a. Breastfeeding at 6 weeks of age (1) | 32.7% | 49.5% | 42.1% | 29.0% | 170 | 377 | 45.1% |
| b. Breastfeeding at 6 months of age | 19.7% | 36.5% | 32.8% | 66.7% | 140 | 377 | 37.1% |

(1) Item appears also on other WCH Section Agreement Addenda. Data for 3A and 3B were not presented due to the limitation in recent trend datasets.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI < 18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI ≥ 30.0): Adapted from the National Academy of Sciences. *Weight Gain During Pregnancy: Reexamining the Guidelines*. Washington, DC : National Academy Press, 2009.

SFY 2014-2015 AGREEMENT ADDENDA DATA

WIC QUALITY ASSURANCE DELIVERABLES

Brunswick County

| Process Outcome Objectives | 2004 to 2006 | 2007 to 2009 | 2010 to 2012 | % Change from 2004 to 2012 | 2012 Data | | |
|---|--------------------|--------------------|--------------------|----------------------------------|-----------|-------|---------|
| | | | | | Num | Den | Percent |
| 1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2) | 72.4% | 76.1% | 74.1% | 2.4% | 396 | 521 | 76.0% |
| 1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services. | 71.3% | 72.5% | 72.9% | 2.2% | 458 | 629 | 72.8% |
| 1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services. | 48.9% | 51.9% | 57.9% | 18.4% | 2,084 | 3,432 | 60.7% |
| 2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy. | 32.6% | 31.9% | 29.7% | -8.8% | 421 | 1,412 | 29.8% |
| 2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum. | 69.8% | 75.6% | 75.3% | 7.9% | 453 | 599 | 75.6% |
| 3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1) | | | | | | | |
| 3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1) | | | | | | | |
| 4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were: | | | | | | | |
| a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1) | 66.6% | 64.5% | 61.8% | -7.1% | 661 | 1,064 | 62.1% |
| b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile) | 16.1% | 17.4% | 17.3% | 7.5% | 184 | 1,064 | 17.3% |
| c. Obese (BMI at or greater than the 95th percentile). | 14.8% | 15.4% | 16.7% | 12.3% | 155 | 1,064 | 14.6% |
| d. Underweight (BMI less than the 5th percentile). | 2.5% | 2.8% | 4.3% | 68.0% | 64 | 1,064 | 6.0% |
| 4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3) | | | | | | | |
| a. Recommended prenatal weight gain | 33.0% | 31.9% | 28.4% | -14.0% | 149 | 518 | 28.8% |
| b. Excessive prenatal weight gain | 50.1% | 48.4% | 52.3% | 4.4% | 266 | 518 | 51.4% |
| c. Inadequate prenatal weight gain | 16.9% | 19.8% | 19.3% | 14.3% | 103 | 518 | 19.9% |
| 5A. Percentage of women who participated in WIC who initiated breastfeeding. | 41.4% | 34.0% | 42.0% | 1.3% | 353 | 777 | 45.4% |
| 5B. Percentage of infants who participated in WIC who were: | | | | | | | |
| a. Breastfeeding at 6 weeks of age (1) | 27.5% | 23.2% | 26.6% | -3.3% | 222 | 751 | 29.6% |
| b. Breastfeeding at 6 months of age | 16.5% | 11.8% | 15.3% | -7.2% | 138 | 751 | 18.4% |

(1) Item appears also on other WCH Section Agreement Addenda. Data for 3A and 3B were not presented due to the limitation in recent trend datasets.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI >=30.0): Adapted from the National Academy of Sciences. Weight Gain During Pregnancy: Reexamining the Guidelines. Washington, DC : National Academy Press, 2009.

SFY 2014-2015 AGREEMENT ADDENDA DATA

WIC QUALITY ASSURANCE DELIVERABLES

Buncombe County

| Process Outcome Objectives | 2004 to 2006 | 2007 to 2009 | 2010 to 2012 | % Change from 2004 to 2012 | 2012 Data | | |
|---|--------------------|--------------------|--------------------|----------------------------------|-----------|-------|---------|
| | | | | | Num | Den | Percent |
| 1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2) | 71.7% | 70.9% | 71.3% | -0.5% | 1,017 | 1,407 | 72.3% |
| 1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services. | 66.6% | 65.3% | 61.1% | -8.3% | 1,058 | 1,737 | 60.9% |
| 1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services. | 51.9% | 52.6% | 52.3% | 0.8% | 4,587 | 8,519 | 53.8% |
| 2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy. | 39.2% | 35.9% | 34.6% | -11.7% | 1,051 | 3,118 | 33.7% |
| 2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum. | 76.3% | 73.9% | 73.1% | -4.1% | 1,010 | 1,416 | 71.3% |
| 3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1) | | | | | | | |
| 3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1) | | | | | | | |
| 4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were: | | | | | | | |
| a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1) | 63.4% | 64.5% | 65.8% | 3.8% | 1,399 | 2,115 | 66.1% |
| b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile) | 17.3% | 16.0% | 16.1% | -6.9% | 333 | 2,115 | 15.7% |
| c. Obese (BMI at or greater than the 95th percentile). | 15.9% | 17.1% | 14.4% | -9.2% | 298 | 2,115 | 14.1% |
| d. Underweight (BMI less than the 5th percentile). | 3.4% | 2.4% | 3.7% | 6.7% | 85 | 2,115 | 4.0% |
| 4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3) | | | | | | | |
| a. Recommended prenatal weight gain | 33.3% | 33.2% | 30.1% | -9.5% | 331 | 1,087 | 30.5% |
| b. Excessive prenatal weight gain | 46.4% | 48.4% | 47.5% | 2.4% | 513 | 1,087 | 47.2% |
| c. Inadequate prenatal weight gain | 20.3% | 18.5% | 22.4% | 10.2% | 243 | 1,087 | 22.4% |
| 5A. Percentage of women who participated in WIC who initiated breastfeeding. | 67.9% | 70.7% | 73.2% | 7.8% | 1,247 | 1,692 | 73.7% |
| 5B. Percentage of infants who participated in WIC who were: | | | | | | | |
| a. Breastfeeding at 6 weeks of age (1) | 35.8% | 37.3% | 40.9% | 14.1% | 733 | 1,594 | 46.0% |
| b. Breastfeeding at 6 months of age | 19.7% | 21.2% | 23.8% | 20.7% | 421 | 1,594 | 26.4% |

(1) Item appears also on other WCH Section Agreement Addenda. Data for 3A and 3B were not presented due to the limitation in recent trend datasets.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI >=30.0): Adapted from the National Academy of Sciences. Weight Gain During Pregnancy: Reexamining the Guidelines. Washington, DC : National Academy Press, 2009.

SFY 2014-2015 AGREEMENT ADDENDA DATA

WIC QUALITY ASSURANCE DELIVERABLES

Burke County

| Process Outcome Objectives | 2004 to 2006 | 2007 to 2009 | 2010 to 2012 | % Change from 2004 to 2012 | 2012 Data | | |
|---|--------------------|--------------------|--------------------|----------------------------------|-----------|-------|---------|
| | | | | | Num | Den | Percent |
| 1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2) | 67.1% | 71.1% | 73.4% | 9.3% | 451 | 574 | 78.6% |
| 1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services. | 76.0% | 76.5% | 78.1% | 2.8% | 521 | 658 | 79.2% |
| 1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services. | 54.4% | 58.6% | 60.5% | 11.1% | 2,150 | 3,417 | 62.9% |
| 2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy. | 24.5% | 19.8% | 29.1% | 18.5% | 390 | 1,240 | 31.5% |
| 2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum. | 83.7% | 83.0% | 83.2% | -0.6% | 482 | 596 | 80.9% |
| 3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1) | | | | | | | |
| 3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1) | | | | | | | |
| 4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were: | | | | | | | |
| a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1) | 66.0% | 62.1% | 64.4% | -2.5% | 678 | 1,026 | 66.1% |
| b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile) | 17.0% | 17.8% | 16.7% | -1.4% | 165 | 1,026 | 16.1% |
| c. Obese (BMI at or greater than the 95th percentile). | 13.7% | 17.5% | 15.6% | 13.4% | 147 | 1,026 | 14.3% |
| d. Underweight (BMI less than the 5th percentile). | 3.2% | 2.5% | 3.3% | 1.0% | 36 | 1,026 | 3.5% |
| 4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3) | | | | | | | |
| a. Recommended prenatal weight gain | 34.6% | 31.5% | 30.8% | -11.0% | 165 | 484 | 34.1% |
| b. Excessive prenatal weight gain | 42.1% | 42.0% | 44.7% | 6.3% | 201 | 484 | 41.5% |
| c. Inadequate prenatal weight gain | 23.3% | 26.5% | 24.5% | 4.8% | 118 | 484 | 24.4% |
| 5A. Percentage of women who participated in WIC who initiated breastfeeding. | 61.3% | 59.5% | 67.8% | 10.6% | 515 | 726 | 70.9% |
| 5B. Percentage of infants who participated in WIC who were: | | | | | | | |
| a. Breastfeeding at 6 weeks of age (1) | 22.8% | 23.7% | 28.2% | 23.5% | 219 | 737 | 29.7% |
| b. Breastfeeding at 6 months of age | 9.5% | 11.1% | 13.8% | 44.6% | 117 | 737 | 15.9% |

(1) Item appears also on other WCH Section Agreement Addenda. Data for 3A and 3B were not presented due to the limitation in recent trend datasets.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI >=30.0): Adapted from the National Academy of Sciences. Weight Gain During Pregnancy: Reexamining the Guidelines. Washington, DC : National Academy Press, 2009.

SFY 2014-2015 AGREEMENT ADDENDA DATA

WIC QUALITY ASSURANCE DELIVERABLES

Cabarrus County

| Process Outcome Objectives | 2004 to 2006 | 2007 to 2009 | 2010 to 2012 | % Change from 2004 to 2012 | 2012 Data | | |
|---|--------------------|--------------------|--------------------|----------------------------------|-----------|-------|---------|
| | | | | | Num | Den | Percent |
| 1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2) | 69.8% | 70.0% | 66.1% | -5.3% | 653 | 968 | 67.5% |
| 1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services. | 72.8% | 71.3% | 69.5% | -4.5% | 883 | 1,296 | 68.1% |
| 1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services. | 54.4% | 56.3% | 52.4% | -3.6% | 3,534 | 6,642 | 53.2% |
| 2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy. | 28.0% | 21.0% | 26.9% | -4.1% | 657 | 2,448 | 26.8% |
| 2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum. | 80.1% | 81.3% | 78.1% | -2.5% | 854 | 1,104 | 77.4% |
| 3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1) | | | | | | | |
| 3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1) | | | | | | | |
| 4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were: | | | | | | | |
| a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1) | 64.5% | 61.4% | 62.8% | -2.7% | 1,369 | 2,196 | 62.3% |
| b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile) | 16.8% | 17.7% | 15.6% | -7.5% | 263 | 2,196 | 12.0% |
| c. Obese (BMI at or greater than the 95th percentile). | 15.0% | 18.6% | 14.7% | -2.1% | 268 | 2,196 | 12.2% |
| d. Underweight (BMI less than the 5th percentile). | 3.7% | 2.3% | 6.9% | 90.1% | 296 | 2,196 | 13.5% |
| 4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3) | | | | | | | |
| a. Recommended prenatal weight gain | 30.2% | 31.1% | 28.9% | -4.0% | 239 | 872 | 27.4% |
| b. Excessive prenatal weight gain | 46.3% | 45.4% | 47.8% | 3.2% | 440 | 872 | 50.5% |
| c. Inadequate prenatal weight gain | 23.6% | 23.5% | 23.3% | -1.1% | 193 | 872 | 22.1% |
| 5A. Percentage of women who participated in WIC who initiated breastfeeding. | 58.7% | 65.9% | 62.2% | 5.9% | 923 | 1,424 | 64.8% |
| 5B. Percentage of infants who participated in WIC who were: | | | | | | | |
| a. Breastfeeding at 6 weeks of age (1) | 36.7% | 38.8% | 37.9% | 3.4% | 562 | 1,447 | 38.8% |
| b. Breastfeeding at 6 months of age | 20.5% | 22.2% | 19.7% | -3.9% | 292 | 1,447 | 20.2% |

(1) Item appears also on other WCH Section Agreement Addenda. Data for 3A and 3B were not presented due to the limitation in recent trend datasets.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI >=30.0): Adapted from the National Academy of Sciences. *Weight Gain During Pregnancy: Reexamining the Guidelines*. Washington, DC : National Academy Press, 2009.

SFY 2014-2015 AGREEMENT ADDENDA DATA

WIC QUALITY ASSURANCE DELIVERABLES

Caldwell County

| Process Outcome Objectives | 2004 to 2006 | 2007 to 2009 | 2010 to 2012 | % Change from 2004 to 2012 | 2012 Data | | |
|---|--------------------|--------------------|--------------------|----------------------------------|-----------|-------|---------|
| | | | | | Num | Den | Percent |
| 1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2) | 78.5% | 72.5% | 79.4% | 1.2% | 449 | 552 | 81.3% |
| 1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services. | 77.8% | 78.8% | 79.9% | 2.8% | 492 | 613 | 80.3% |
| 1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services. | 57.9% | 60.1% | 58.8% | 1.6% | 1,872 | 3,107 | 60.3% |
| 2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy. | 32.4% | 33.3% | 26.7% | -17.5% | 421 | 1,756 | 24.0% |
| 2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum. | 78.3% | 78.0% | 78.8% | 0.5% | 437 | 557 | 78.5% |
| 3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1) | | | | | | | |
| 3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1) | | | | | | | |
| 4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were: | | | | | | | |
| a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1) | 62.5% | 57.7% | 63.3% | 1.2% | 467 | 720 | 64.9% |
| b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile) | 18.0% | 20.4% | 17.9% | -0.5% | 131 | 720 | 18.2% |
| c. Obese (BMI at or greater than the 95th percentile). | 16.5% | 19.6% | 15.3% | -7.3% | 93 | 720 | 12.9% |
| d. Underweight (BMI less than the 5th percentile). | 3.0% | 2.3% | 3.6% | 19.0% | 29 | 720 | 4.0% |
| 4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3) | | | | | | | |
| a. Recommended prenatal weight gain | 29.7% | 29.0% | 29.4% | -0.9% | 140 | 510 | 27.5% |
| b. Excessive prenatal weight gain | 51.8% | 51.8% | 48.8% | -5.9% | 252 | 510 | 49.4% |
| c. Inadequate prenatal weight gain | 18.5% | 19.2% | 21.8% | 17.9% | 118 | 510 | 23.1% |
| 5A. Percentage of women who participated in WIC who initiated breastfeeding. | 47.3% | 49.0% | 59.0% | 24.8% | 422 | 667 | 63.3% |
| 5B. Percentage of infants who participated in WIC who were: | | | | | | | |
| a. Breastfeeding at 6 weeks of age (1) | 25.2% | 27.3% | 29.3% | 16.2% | 220 | 670 | 32.8% |
| b. Breastfeeding at 6 months of age | 12.1% | 11.5% | 12.6% | 3.6% | 93 | 670 | 13.9% |

(1) Item appears also on other WCH Section Agreement Addenda. Data for 3A and 3B were not presented due to the limitation in recent trend datasets.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI >=30.0): Adapted from the National Academy of Sciences. *Weight Gain During Pregnancy: Reexamining the Guidelines*. Washington, DC : National Academy Press, 2009.

SFY 2014-2015 AGREEMENT ADDENDA DATA

WIC QUALITY ASSURANCE DELIVERABLES

Carteret County

| Process Outcome Objectives | 2004 to 2006 | 2007 to 2009 | 2010 to 2012 | % Change from 2004 to 2012 | 2012 Data | | |
|---|--------------------|--------------------|--------------------|----------------------------------|-----------|-------|---------|
| | | | | | Num | Den | Percent |
| 1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2) | 63.7% | 66.1% | 71.9% | 12.8% | 206 | 281 | 73.3% |
| 1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services. | 72.1% | 76.4% | 74.9% | 3.8% | 251 | 345 | 72.8% |
| 1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services. | 53.5% | 55.5% | 57.9% | 8.1% | 1,025 | 1,720 | 59.6% |
| 2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy. | 31.7% | 30.2% | 29.4% | -7.3% | 248 | 818 | 30.3% |
| 2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum. | 78.4% | 64.2% | 76.6% | -2.3% | 263 | 347 | 75.8% |
| 3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1) | | | | | | | |
| 3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1) | | | | | | | |
| 4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were: | | | | | | | |
| a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1) | 63.2% | 65.3% | 63.8% | 1.1% | 261 | 413 | 63.2% |
| b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile) | 17.1% | 15.3% | 16.5% | -3.5% | 64 | 413 | 15.5% |
| c. Obese (BMI at or greater than the 95th percentile). | 16.6% | 16.9% | 14.0% | -15.6% | 69 | 413 | 16.7% |
| d. Underweight (BMI less than the 5th percentile). | 3.1% | 2.5% | 5.6% | 80.4% | 19 | 413 | 4.6% |
| 4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3) | | | | | | | |
| a. Recommended prenatal weight gain | 33.5% | 34.2% | 28.1% | -16.3% | 75 | 292 | 25.7% |
| b. Excessive prenatal weight gain | 52.2% | 49.9% | 50.9% | -2.5% | 158 | 292 | 54.1% |
| c. Inadequate prenatal weight gain | 14.3% | 15.9% | 21.1% | 47.3% | 59 | 292 | 20.2% |
| 5A. Percentage of women who participated in WIC who initiated breastfeeding. | 52.9% | 60.3% | 64.9% | 22.7% | 306 | 447 | 68.5% |
| 5B. Percentage of infants who participated in WIC who were: | | | | | | | |
| a. Breastfeeding at 6 weeks of age (1) | 28.8% | 31.0% | 34.8% | 20.9% | 189 | 479 | 39.5% |
| b. Breastfeeding at 6 months of age | 11.8% | 13.7% | 15.8% | 34.1% | 81 | 479 | 16.9% |

(1) Item appears also on other WCH Section Agreement Addenda. Data for 3A and 3B were not presented due to the limitation in recent trend datasets.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI >=30.0): Adapted from the National Academy of Sciences. Weight Gain During Pregnancy: Reexamining the Guidelines. Washington, DC : National Academy Press, 2009.

SFY 2014-2015 AGREEMENT ADDENDA DATA

WIC QUALITY ASSURANCE DELIVERABLES

Caswell County

| Process Outcome Objectives | 2004 to 2006 | 2007 to 2009 | 2010 to 2012 | % Change from 2004 to 2012 | 2012 Data | | |
|---|--------------------|--------------------|--------------------|----------------------------------|-----------|-----|---------|
| | | | | | Num | Den | Percent |
| 1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2) | 81.3% | 82.7% | 69.4% | -14.6% | 83 | 129 | 64.3% |
| 1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services. | 75.4% | 76.7% | 71.5% | -5.1% | 102 | 146 | 69.9% |
| 1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services. | 57.5% | 61.7% | 60.1% | 4.6% | 520 | 829 | 62.7% |
| 2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy. | 39.6% | 41.9% | 35.9% | -9.3% | 112 | 329 | 34.0% |
| 2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum. | 77.1% | 78.9% | 76.4% | -0.8% | 98 | 131 | 74.8% |
| 3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1) | | | | | | | |
| 3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1) | | | | | | | |
| 4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were: | | | | | | | |
| a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1) | 68.0% | 69.4% | 67.1% | -1.3% | 179 | 289 | 61.9% |
| b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile) | 15.5% | 15.0% | 16.1% | 4.2% | 50 | 289 | 17.3% |
| c. Obese (BMI at or greater than the 95th percentile). | 14.0% | 11.3% | 12.6% | -10.0% | 41 | 289 | 14.2% |
| d. Underweight (BMI less than the 5th percentile). | 2.5% | 4.3% | 4.2% | 65.6% | 19 | 289 | 6.6% |
| 4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3) | | | | | | | |
| a. Recommended prenatal weight gain | 31.0% | 38.4% | 26.8% | -13.5% | 31 | 94 | 33.0% |
| b. Excessive prenatal weight gain | 46.5% | 41.6% | 46.1% | -0.8% | 44 | 94 | 46.8% |
| c. Inadequate prenatal weight gain | 22.6% | 20.0% | 27.1% | 20.2% | 19 | 94 | 20.2% |
| 5A. Percentage of women who participated in WIC who initiated breastfeeding. | 45.1% | 52.4% | 57.4% | 27.4% | 103 | 164 | 62.8% |
| 5B. Percentage of infants who participated in WIC who were: | | | | | | | |
| a. Breastfeeding at 6 weeks of age (1) | 22.3% | 28.2% | 32.2% | 44.3% | 61 | 168 | 36.3% |
| b. Breastfeeding at 6 months of age | 8.4% | 9.1% | 13.3% | 58.9% | 23 | 168 | 13.7% |

(1) Item appears also on other WCH Section Agreement Addenda. Data for 3A and 3B were not presented due to the limitation in recent trend datasets.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI >=30.0): Adapted from the National Academy of Sciences. Weight Gain During Pregnancy: Reexamining the Guidelines. Washington, DC : National Academy Press, 2009.

SFY 2014-2015 AGREEMENT ADDENDA DATA

WIC QUALITY ASSURANCE DELIVERABLES

Catawba County

| Process Outcome Objectives | 2004 to 2006 | 2007 to 2009 | 2010 to 2012 | % Change from 2004 to 2012 | 2012 Data | | |
|---|--------------------|--------------------|--------------------|----------------------------------|-----------|-------|---------|
| | | | | | Num | Den | Percent |
| 1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2) | 72.8% | 73.4% | 69.9% | -4.0% | 807 | 1,114 | 72.4% |
| 1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services. | 76.6% | 75.8% | 76.5% | -0.1% | 1,008 | 1,313 | 76.8% |
| 1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services. | 50.0% | 53.1% | 55.7% | 11.3% | 3,797 | 6,378 | 59.5% |
| 2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy. | 26.8% | 26.2% | 34.9% | 30.2% | 659 | 1,945 | 33.9% |
| 2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum. | 57.0% | 60.7% | 70.9% | 24.2% | 775 | 1,016 | 76.3% |
| 3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1) | | | | | | | |
| 3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1) | | | | | | | |
| 4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were: | | | | | | | |
| a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1) | 60.9% | 62.0% | 63.6% | 4.5% | 736 | 1,167 | 63.1% |
| b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile) | 18.6% | 19.0% | 17.1% | -7.9% | 208 | 1,167 | 17.8% |
| c. Obese (BMI at or greater than the 95th percentile). | 17.5% | 17.1% | 15.4% | -12.2% | 176 | 1,167 | 15.1% |
| d. Underweight (BMI less than the 5th percentile). | 3.0% | 1.9% | 3.9% | 28.8% | 47 | 1,167 | 4.0% |
| 4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3) | | | | | | | |
| a. Recommended prenatal weight gain | 32.7% | 31.1% | 30.6% | -6.5% | 241 | 846 | 28.5% |
| b. Excessive prenatal weight gain | 39.2% | 34.0% | 45.3% | 15.6% | 405 | 846 | 47.9% |
| c. Inadequate prenatal weight gain | 28.1% | 35.0% | 24.1% | -14.2% | 200 | 846 | 23.6% |
| 5A. Percentage of women who participated in WIC who initiated breastfeeding. | 52.1% | 62.5% | 61.5% | 18.0% | 787 | 1,293 | 60.9% |
| 5B. Percentage of infants who participated in WIC who were: | | | | | | | |
| a. Breastfeeding at 6 weeks of age (1) | 29.3% | 30.8% | 33.9% | 15.5% | 475 | 1,334 | 35.6% |
| b. Breastfeeding at 6 months of age | 13.5% | 14.2% | 18.2% | 34.8% | 266 | 1,334 | 19.9% |

(1) Item appears also on other WCH Section Agreement Addenda. Data for 3A and 3B were not presented due to the limitation in recent trend datasets.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI >=30.0): Adapted from the National Academy of Sciences. Weight Gain During Pregnancy: Reexamining the Guidelines. Washington, DC : National Academy Press, 2009.

SFY 2014-2015 AGREEMENT ADDENDA DATA

WIC QUALITY ASSURANCE DELIVERABLES

Chatham County

| Process Outcome Objectives | 2004 to 2006 | 2007 to 2009 | 2010 to 2012 | % Change from 2004 to 2012 | 2012 Data | | |
|---|--------------------|--------------------|--------------------|----------------------------------|-----------|-------|---------|
| | | | | | Num | Den | Percent |
| 1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2) | 64.8% | 70.5% | 70.7% | 9.1% | 143 | 195 | 73.3% |
| 1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services. | 47.7% | 49.4% | 63.3% | 32.7% | 226 | 342 | 66.1% |
| 1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services. | 41.9% | 43.9% | 51.8% | 23.7% | 1,027 | 1,807 | 56.8% |
| 2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy. | 26.4% | 29.4% | 27.0% | 2.3% | 676 | 2,451 | 27.6% |
| 2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum. | 59.5% | 62.6% | 69.4% | 16.7% | 820 | 1,175 | 69.8% |
| 3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1) | | | | | | | |
| 3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1) | | | | | | | |
| 4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were: | | | | | | | |
| a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1) | 61.0% | 62.3% | 61.9% | 1.5% | 680 | 1,080 | 63.0% |
| b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile) | 18.2% | 15.7% | 17.0% | -6.4% | 172 | 1,080 | 15.9% |
| c. Obese (BMI at or greater than the 95th percentile). | 17.9% | 19.3% | 17.4% | -3.1% | 183 | 1,080 | 16.9% |
| d. Underweight (BMI less than the 5th percentile). | 2.9% | 2.6% | 3.8% | 28.0% | 45 | 1,080 | 4.2% |
| 4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3) | | | | | | | |
| a. Recommended prenatal weight gain | 31.1% | 37.1% | 29.0% | -6.7% | 43 | 183 | 23.5% |
| b. Excessive prenatal weight gain | 47.8% | 43.1% | 45.5% | -4.8% | 95 | 183 | 51.9% |
| c. Inadequate prenatal weight gain | 21.1% | 19.8% | 25.5% | 20.7% | 45 | 183 | 24.6% |
| 5A. Percentage of women who participated in WIC who initiated breastfeeding. | 63.5% | 67.3% | 70.6% | 11.2% | 1,021 | 1,441 | 70.9% |
| 5B. Percentage of infants who participated in WIC who were: | | | | | | | |
| a. Breastfeeding at 6 weeks of age (1) | 50.2% | 53.5% | 53.4% | 6.4% | 753 | 1,386 | 54.3% |
| b. Breastfeeding at 6 months of age | 29.7% | 33.6% | 33.4% | 12.6% | 473 | 1,386 | 34.1% |

(1) Item appears also on other WCH Section Agreement Addenda. Data for 3A and 3B were not presented due to the limitation in recent trend datasets.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI >=30.0): Adapted from the National Academy of Sciences. Weight Gain During Pregnancy: Reexamining the Guidelines. Washington, DC : National Academy Press, 2009.

SFY 2014-2015 AGREEMENT ADDENDA DATA

WIC QUALITY ASSURANCE DELIVERABLES

Cherokee County

| Process Outcome Objectives | 2004 to 2006 | 2007 to 2009 | 2010 to 2012 | % Change from 2004 to 2012 | 2012 Data | | |
|---|--------------------|--------------------|--------------------|----------------------------------|-----------|-----|---------|
| | | | | | Num | Den | Percent |
| 1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2) | 75.6% | 81.3% | 80.3% | 6.3% | 129 | 151 | 85.4% |
| 1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services. | 76.2% | 73.5% | 73.9% | -3.0% | 127 | 166 | 76.5% |
| 1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services. | 62.9% | 65.6% | 65.0% | 3.4% | 639 | 964 | 66.3% |
| 2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy. | 39.6% | 47.0% | 44.9% | 13.5% | 151 | 366 | 41.3% |
| 2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum. | 80.1% | 82.0% | 79.8% | -0.4% | 137 | 173 | 79.2% |
| 3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1) | | | | | | | |
| 3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1) | | | | | | | |
| 4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were: | | | | | | | |
| a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1) | 68.5% | 68.4% | 65.3% | -4.6% | 173 | 258 | 67.1% |
| b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile) | 14.7% | 15.3% | 13.6% | -7.8% | 32 | 258 | 12.4% |
| c. Obese (BMI at or greater than the 95th percentile). | 12.2% | 13.8% | 16.6% | 36.0% | 43 | 258 | 16.7% |
| d. Underweight (BMI less than the 5th percentile). | 4.6% | 2.5% | 4.5% | -1.7% | 10 | 258 | 3.9% |
| 4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3) | | | | | | | |
| a. Recommended prenatal weight gain | 29.2% | 31.0% | 30.0% | 2.7% | 47 | 143 | 32.9% |
| b. Excessive prenatal weight gain | 47.0% | 47.0% | 46.5% | -0.9% | 64 | 143 | 44.8% |
| c. Inadequate prenatal weight gain | 23.8% | 22.0% | 23.5% | -1.4% | 32 | 143 | 22.4% |
| 5A. Percentage of women who participated in WIC who initiated breastfeeding. | 43.7% | 55.6% | 55.0% | 25.9% | 127 | 214 | 59.3% |
| 5B. Percentage of infants who participated in WIC who were: | | | | | | | |
| a. Breastfeeding at 6 weeks of age (1) | 27.7% | 33.4% | 31.3% | 12.7% | 57 | 182 | 31.3% |
| b. Breastfeeding at 6 months of age | 12.7% | 15.8% | 15.8% | 24.6% | 32 | 182 | 17.6% |

(1) Item appears also on other WCH Section Agreement Addenda. Data for 3A and 3B were not presented due to the limitation in recent trend datasets.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI >=30.0): Adapted from the National Academy of Sciences. Weight Gain During Pregnancy: Reexamining the Guidelines. Washington, DC : National Academy Press, 2009.

SFY 2014-2015 AGREEMENT ADDENDA DATA

WIC QUALITY ASSURANCE DELIVERABLES

Clay County

| Process Outcome Objectives | 2004 to 2006 | 2007 to 2009 | 2010 to 2012 | % Change from 2004 to 2012 | 2012 Data | | |
|---|--------------------|--------------------|--------------------|----------------------------------|-----------|-----|---------|
| | | | | | Num | Den | Percent |
| 1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2) | 80.4% | 84.2% | 76.4% | -5.0% | 44 | 55 | 80.0% |
| 1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services. | 74.6% | 73.1% | 72.4% | -3.1% | 50 | 79 | 63.3% |
| 1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services. | 68.7% | 68.2% | 62.8% | -8.7% | 246 | 413 | 59.6% |
| 2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy. | 47.3% | 54.7% | 43.1% | -8.7% | 79 | 200 | 39.5% |
| 2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum. | 75.8% | 71.2% | 68.4% | -9.7% | 56 | 87 | 64.4% |
| 3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1) | | | | | | | |
| 3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1) | | | | | | | |
| 4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were: | | | | | | | |
| a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1) | 61.2% | 64.8% | 69.1% | 12.9% | 140 | 195 | 71.8% |
| b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile) | 15.4% | 14.0% | 15.1% | -1.9% | 27 | 195 | 13.8% |
| c. Obese (BMI at or greater than the 95th percentile). | 18.8% | 17.6% | 13.6% | -27.3% | 23 | 195 | 11.8% |
| d. Underweight (BMI less than the 5th percentile). | 4.6% | 3.6% | 2.1% | -53.9% | 5 | 195 | 2.6% |
| 4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3) | | | | | | | |
| a. Recommended prenatal weight gain | 27.3% | 26.3% | 31.2% | 14.2% | 12 | 46 | 26.1% |
| b. Excessive prenatal weight gain | 51.8% | 51.8% | 51.6% | -0.4% | 25 | 46 | 54.3% |
| c. Inadequate prenatal weight gain | 20.9% | 21.9% | 17.2% | -17.6% | 9 | 46 | 19.6% |
| 5A. Percentage of women who participated in WIC who initiated breastfeeding. | 49.6% | 56.3% | 62.0% | 25.2% | 61 | 100 | 61.0% |
| 5B. Percentage of infants who participated in WIC who were: | | | | | | | |
| a. Breastfeeding at 6 weeks of age (1) | 31.4% | 36.7% | 42.7% | 36.3% | 32 | 72 | 44.4% |
| b. Breastfeeding at 6 months of age | 17.4% | 21.2% | 23.2% | 33.8% | 18 | 72 | 25.0% |

(1) Item appears also on other WCH Section Agreement Addenda. Data for 3A and 3B were not presented due to the limitation in recent trend datasets.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI >=30.0): Adapted from the National Academy of Sciences. Weight Gain During Pregnancy: Reexamining the Guidelines. Washington, DC : National Academy Press, 2009.

SFY 2014-2015 AGREEMENT ADDENDA DATA

WIC QUALITY ASSURANCE DELIVERABLES

Cleveland County

| Process Outcome Objectives | 2004 to 2006 | 2007 to 2009 | 2010 to 2012 | % Change from 2004 to 2012 | 2012 Data | | |
|---|--------------------|--------------------|--------------------|----------------------------------|-----------|-------|---------|
| | | | | | Num | Den | Percent |
| 1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2) | 81.7% | 83.6% | 84.8% | 3.8% | 649 | 766 | 84.7% |
| 1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services. | 77.6% | 75.4% | 76.7% | -1.2% | 648 | 828 | 78.3% |
| 1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services. | 49.2% | 50.8% | 53.6% | 9.0% | 2,447 | 4,299 | 56.9% |
| 2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy. | 34.0% | 36.3% | 32.7% | -3.9% | 571 | 1,920 | 29.7% |
| 2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum. | 13.4% | 31.6% | 58.1% | 334.8% | 461 | 806 | 57.2% |
| 3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1) | | | | | | | |
| 3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1) | | | | | | | |
| 4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were: | | | | | | | |
| a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1) | 70.3% | 67.7% | 66.0% | -6.1% | 855 | 1,257 | 68.0% |
| b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile) | 12.1% | 12.7% | 14.3% | 18.9% | 151 | 1,257 | 12.0% |
| c. Obese (BMI at or greater than the 95th percentile). | 10.8% | 12.3% | 13.7% | 27.1% | 152 | 1,257 | 12.1% |
| d. Underweight (BMI less than the 5th percentile). | 6.9% | 7.3% | 6.0% | -13.3% | 99 | 1,257 | 7.9% |
| 4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3) | | | | | | | |
| a. Recommended prenatal weight gain | 28.2% | 28.3% | 22.6% | -20.0% | 139 | 714 | 19.5% |
| b. Excessive prenatal weight gain | 48.5% | 40.9% | 35.1% | -27.7% | 226 | 714 | 31.7% |
| c. Inadequate prenatal weight gain | 23.3% | 30.8% | 42.4% | 82.1% | 349 | 714 | 48.9% |
| 5A. Percentage of women who participated in WIC who initiated breastfeeding. | 19.4% | 25.1% | 47.4% | 144.8% | 429 | 925 | 46.4% |
| 5B. Percentage of infants who participated in WIC who were: | | | | | | | |
| a. Breastfeeding at 6 weeks of age (1) | 11.4% | 11.9% | 33.1% | 189.9% | 380 | 868 | 43.8% |
| b. Breastfeeding at 6 months of age | 6.6% | 6.6% | 27.0% | 309.5% | 318 | 868 | 36.6% |

(1) Item appears also on other WCH Section Agreement Addenda. Data for 3A and 3B were not presented due to the limitation in recent trend datasets.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI >=30.0): Adapted from the National Academy of Sciences. Weight Gain During Pregnancy: Reexamining the Guidelines. Washington, DC : National Academy Press, 2009.

SFY 2014-2015 AGREEMENT ADDENDA DATA

WIC QUALITY ASSURANCE DELIVERABLES

Columbus County

| Process Outcome Objectives | 2004 to 2006 | 2007 to 2009 | 2010 to 2012 | % Change from 2004 to 2012 | 2012 Data | | |
|---|--------------------|--------------------|--------------------|----------------------------------|-----------|-------|---------|
| | | | | | Num | Den | Percent |
| 1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2) | 70.8% | 72.5% | 76.0% | 7.4% | 353 | 444 | 79.5% |
| 1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services. | 80.1% | 79.4% | 79.5% | -0.8% | 425 | 525 | 81.0% |
| 1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services. | 58.5% | 61.3% | 62.2% | 6.3% | 1,843 | 2,827 | 65.2% |
| 2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy. | 30.7% | 31.9% | 30.6% | -0.2% | 337 | 1,019 | 33.1% |
| 2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum. | 81.0% | 85.5% | 85.0% | 5.0% | 394 | 466 | 84.5% |
| 3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1) | | | | | | | |
| 3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1) | | | | | | | |
| 4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were: | | | | | | | |
| a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1) | 65.2% | 64.5% | 60.1% | -7.8% | 681 | 1,078 | 63.2% |
| b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile) | 15.9% | 15.8% | 16.5% | 3.5% | 165 | 1,078 | 15.3% |
| c. Obese (BMI at or greater than the 95th percentile). | 14.4% | 15.0% | 19.6% | 36.1% | 190 | 1,078 | 17.6% |
| d. Underweight (BMI less than the 5th percentile). | 4.5% | 4.8% | 3.8% | -14.4% | 42 | 1,078 | 3.9% |
| 4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3) | | | | | | | |
| a. Recommended prenatal weight gain | 26.6% | 28.1% | 29.2% | 9.8% | 131 | 415 | 31.6% |
| b. Excessive prenatal weight gain | 45.1% | 44.7% | 45.3% | 0.5% | 178 | 415 | 42.9% |
| c. Inadequate prenatal weight gain | 28.3% | 27.2% | 25.5% | -10.1% | 106 | 415 | 25.5% |
| 5A. Percentage of women who participated in WIC who initiated breastfeeding. | 41.2% | 35.4% | 37.3% | -9.3% | 238 | 591 | 40.3% |
| 5B. Percentage of infants who participated in WIC who were: | | | | | | | |
| a. Breastfeeding at 6 weeks of age (1) | 23.3% | 22.1% | 22.4% | -3.8% | 149 | 599 | 24.9% |
| b. Breastfeeding at 6 months of age | 11.7% | 14.2% | 17.3% | 47.4% | 106 | 599 | 17.7% |

(1) Item appears also on other WCH Section Agreement Addenda. Data for 3A and 3B were not presented due to the limitation in recent trend datasets.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI >=30.0): Adapted from the National Academy of Sciences. Weight Gain During Pregnancy: Reexamining the Guidelines. Washington, DC : National Academy Press, 2009.

SFY 2014-2015 AGREEMENT ADDENDA DATA

WIC QUALITY ASSURANCE DELIVERABLES

Craven County

| Process Outcome Objectives | 2004 to 2006 | 2007 to 2009 | 2010 to 2012 | % Change from 2004 to 2012 | 2012 Data | | |
|---|--------------------|--------------------|--------------------|----------------------------------|-----------|-------|---------|
| | | | | | Num | Den | Percent |
| 1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2) | 78.5% | 81.0% | 84.1% | 7.0% | 516 | 598 | 86.3% |
| 1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services. | 76.5% | 76.8% | 75.1% | -1.7% | 481 | 661 | 72.8% |
| 1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services. | 53.7% | 56.2% | 59.8% | 11.2% | 2,104 | 3,523 | 59.7% |
| 2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy. | 29.7% | 31.9% | 31.0% | 4.4% | 770 | 2,603 | 29.6% |
| 2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum. | 33.2% | 33.9% | 53.0% | 59.7% | 746 | 1,019 | 73.2% |
| 3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1) | | | | | | | |
| 3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1) | | | | | | | |
| 4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were: | | | | | | | |
| a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1) | 64.4% | 67.9% | 65.4% | 1.5% | 1,024 | 1,604 | 63.8% |
| b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile) | 16.7% | 15.3% | 15.4% | -7.5% | 257 | 1,604 | 16.0% |
| c. Obese (BMI at or greater than the 95th percentile). | 14.9% | 12.9% | 12.9% | -13.6% | 200 | 1,604 | 12.5% |
| d. Underweight (BMI less than the 5th percentile). | 4.0% | 4.0% | 6.3% | 57.4% | 123 | 1,604 | 7.7% |
| 4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3) | | | | | | | |
| a. Recommended prenatal weight gain | 30.5% | 33.9% | 28.4% | -7.0% | 239 | 854 | 28.0% |
| b. Excessive prenatal weight gain | 49.2% | 45.9% | 43.9% | -10.8% | 405 | 854 | 47.4% |
| c. Inadequate prenatal weight gain | 20.2% | 20.2% | 27.7% | 36.9% | 210 | 854 | 24.6% |
| 5A. Percentage of women who participated in WIC who initiated breastfeeding. | 47.9% | 41.8% | 38.7% | -19.3% | 563 | 1,206 | 46.7% |
| 5B. Percentage of infants who participated in WIC who were: | | | | | | | |
| a. Breastfeeding at 6 weeks of age (1) | 25.0% | 25.2% | 22.8% | -8.5% | 377 | 1,244 | 30.3% |
| b. Breastfeeding at 6 months of age | 12.8% | 13.3% | 14.1% | 9.7% | 259 | 1,244 | 20.8% |

(1) Item appears also on other WCH Section Agreement Addenda. Data for 3A and 3B were not presented due to the limitation in recent trend datasets.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI >=30.0): Adapted from the National Academy of Sciences. *Weight Gain During Pregnancy: Reexamining the Guidelines*. Washington, DC : National Academy Press, 2009.

SFY 2014-2015 AGREEMENT ADDENDA DATA

WIC QUALITY ASSURANCE DELIVERABLES

Cumberland County

| Process Outcome Objectives | 2004 to 2006 | 2007 to 2009 | 2010 to 2012 | % Change from 2004 to 2012 | 2012 Data | | |
|---|--------------------|--------------------|--------------------|----------------------------------|-----------|--------|---------|
| | | | | | Num | Den | Percent |
| 1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2) | 73.6% | 75.5% | 74.7% | 1.5% | 1,681 | 2,132 | 78.8% |
| 1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services. | 77.5% | 76.4% | 73.5% | -5.1% | 1,767 | 2,441 | 72.4% |
| 1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services. | 54.8% | 55.2% | 55.2% | 0.9% | 7,438 | 13,264 | 56.1% |
| 2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy. | 31.2% | 30.1% | 30.8% | -1.2% | 2,536 | 8,055 | 31.5% |
| 2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum. | 76.2% | 75.4% | 75.3% | -1.2% | 2,608 | 3,483 | 74.9% |
| 3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1) | | | | | | | |
| 3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1) | | | | | | | |
| 4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were: | | | | | | | |
| a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1) | 73.4% | 74.8% | 69.2% | -5.7% | 4,109 | 5,984 | 68.7% |
| b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile) | 11.6% | 9.3% | 11.7% | 1.1% | 673 | 5,984 | 11.2% |
| c. Obese (BMI at or greater than the 95th percentile). | 8.1% | 6.9% | 9.3% | 14.7% | 577 | 5,984 | 9.6% |
| d. Underweight (BMI less than the 5th percentile). | 6.9% | 9.0% | 9.8% | 41.8% | 625 | 5,984 | 10.4% |
| 4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3) | | | | | | | |
| a. Recommended prenatal weight gain | 31.5% | 29.3% | 27.8% | -11.9% | 852 | 3,028 | 28.1% |
| b. Excessive prenatal weight gain | 51.3% | 52.1% | 51.5% | 0.2% | 1,531 | 3,028 | 50.6% |
| c. Inadequate prenatal weight gain | 17.1% | 18.6% | 20.7% | 21.2% | 645 | 3,028 | 21.3% |
| 5A. Percentage of women who participated in WIC who initiated breastfeeding. | 54.6% | 59.1% | 64.8% | 18.8% | 2,865 | 4,578 | 62.6% |
| 5B. Percentage of infants who participated in WIC who were: | | | | | | | |
| a. Breastfeeding at 6 weeks of age (1) | 22.9% | 27.3% | 30.5% | 32.8% | 1,700 | 5,062 | 33.6% |
| b. Breastfeeding at 6 months of age | 9.8% | 11.5% | 14.8% | 50.7% | 894 | 5,062 | 17.7% |

(1) Item appears also on other WCH Section Agreement Addenda. Data for 3A and 3B were not presented due to the limitation in recent trend datasets.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI >=30.0): Adapted from the National Academy of Sciences. Weight Gain During Pregnancy: Reexamining the Guidelines. Washington, DC : National Academy Press, 2009.

SFY 2014-2015 AGREEMENT ADDENDA DATA

WIC QUALITY ASSURANCE DELIVERABLES

Dare County

| Process Outcome Objectives | 2004 to 2006 | 2007 to 2009 | 2010 to 2012 | % Change from 2004 to 2012 | 2012 Data | | |
|---|--------------------|--------------------|--------------------|----------------------------------|-----------|-------|---------|
| | | | | | Num | Den | Percent |
| 1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2) | 59.2% | 65.4% | 64.6% | 9.1% | 120 | 200 | 60.0% |
| 1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services. | 58.5% | 66.5% | 65.7% | 12.3% | 174 | 269 | 64.7% |
| 1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services. | 52.5% | 56.5% | 59.4% | 13.1% | 692 | 1,195 | 57.9% |
| 2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy. | 26.8% | 31.9% | 28.7% | 7.2% | 158 | 582 | 27.1% |
| 2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum. | 65.9% | 70.3% | 75.0% | 13.8% | 165 | 224 | 73.7% |
| 3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1) | | | | | | | |
| 3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1) | | | | | | | |
| 4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were: | | | | | | | |
| a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1) | 62.7% | 65.6% | 60.9% | -3.0% | 189 | 315 | 60.0% |
| b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile) | 18.4% | 15.1% | 20.6% | 12.0% | 67 | 315 | 21.3% |
| c. Obese (BMI at or greater than the 95th percentile). | 16.6% | 16.5% | 16.1% | -3.0% | 51 | 315 | 16.2% |
| d. Underweight (BMI less than the 5th percentile). | 2.3% | 2.7% | 2.4% | 6.9% | 8 | 315 | 2.5% |
| 4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3) | | | | | | | |
| a. Recommended prenatal weight gain | 37.1% | 37.8% | 31.1% | -16.2% | 38 | 155 | 24.5% |
| b. Excessive prenatal weight gain | 44.2% | 35.6% | 45.2% | 2.2% | 71 | 155 | 45.8% |
| c. Inadequate prenatal weight gain | 18.7% | 26.6% | 23.7% | 26.9% | 46 | 155 | 29.7% |
| 5A. Percentage of women who participated in WIC who initiated breastfeeding. | 57.2% | 67.4% | 73.5% | 28.4% | 210 | 286 | 73.4% |
| 5B. Percentage of infants who participated in WIC who were: | | | | | | | |
| a. Breastfeeding at 6 weeks of age (1) | 54.7% | 58.6% | 61.0% | 11.6% | 161 | 236 | 68.2% |
| b. Breastfeeding at 6 months of age | 28.6% | 32.3% | 41.6% | 45.1% | 112 | 236 | 47.5% |

(1) Item appears also on other WCH Section Agreement Addenda. Data for 3A and 3B were not presented due to the limitation in recent trend datasets.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI >=30.0): Adapted from the National Academy of Sciences. Weight Gain During Pregnancy: Reexamining the Guidelines. Washington, DC : National Academy Press, 2009.

SFY 2014-2015 AGREEMENT ADDENDA DATA

WIC QUALITY ASSURANCE DELIVERABLES

Davidson County

| Process Outcome Objectives | 2004 to 2006 | 2007 to 2009 | 2010 to 2012 | % Change from 2004 to 2012 | 2012 Data | | |
|---|--------------------|--------------------|--------------------|----------------------------------|-----------|-------|---------|
| | | | | | Num | Den | Percent |
| 1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2) | 69.3% | 70.8% | 74.0% | 6.7% | 774 | 1,008 | 76.8% |
| 1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services. | 70.6% | 73.2% | 74.2% | 5.2% | 867 | 1,154 | 75.1% |
| 1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services. | 49.5% | 53.3% | 55.8% | 12.6% | 3,493 | 6,056 | 57.7% |
| 2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy. | 31.7% | 30.2% | 32.6% | 2.9% | 664 | 2,018 | 32.9% |
| 2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum. | 73.4% | 78.8% | 76.8% | 4.6% | 836 | 1,071 | 78.1% |
| 3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1) | | | | | | | |
| 3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1) | | | | | | | |
| 4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were: | | | | | | | |
| a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1) | 70.5% | 67.3% | 65.2% | -7.4% | 1,123 | 1,685 | 66.6% |
| b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile) | 13.9% | 15.4% | 16.3% | 17.5% | 274 | 1,685 | 16.3% |
| c. Obese (BMI at or greater than the 95th percentile). | 11.5% | 14.7% | 15.0% | 30.4% | 228 | 1,685 | 13.5% |
| d. Underweight (BMI less than the 5th percentile). | 4.2% | 2.6% | 3.5% | -16.2% | 60 | 1,685 | 3.6% |
| 4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3) | | | | | | | |
| a. Recommended prenatal weight gain | 31.3% | 33.6% | 28.2% | -9.9% | 226 | 838 | 27.0% |
| b. Excessive prenatal weight gain | 46.4% | 42.1% | 50.1% | 8.1% | 426 | 838 | 50.8% |
| c. Inadequate prenatal weight gain | 22.3% | 24.2% | 21.6% | -3.0% | 186 | 838 | 22.2% |
| 5A. Percentage of women who participated in WIC who initiated breastfeeding. | 52.7% | 64.8% | 68.2% | 29.3% | 766 | 1,316 | 58.2% |
| 5B. Percentage of infants who participated in WIC who were: | | | | | | | |
| a. Breastfeeding at 6 weeks of age (1) | 28.4% | 29.1% | 30.4% | 7.2% | 438 | 1,279 | 34.2% |
| b. Breastfeeding at 6 months of age | 14.2% | 15.1% | 16.0% | 12.1% | 211 | 1,279 | 16.5% |

(1) Item appears also on other WCH Section Agreement Addenda. Data for 3A and 3B were not presented due to the limitation in recent trend datasets.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI >=30.0): Adapted from the National Academy of Sciences. *Weight Gain During Pregnancy: Reexamining the Guidelines*. Washington, DC : National Academy Press, 2009.

SFY 2014-2015 AGREEMENT ADDENDA DATA

WIC QUALITY ASSURANCE DELIVERABLES

Davie County

| Process Outcome Objectives | 2004 to 2006 | 2007 to 2009 | 2010 to 2012 | % Change from 2004 to 2012 | 2012 Data | | |
|---|--------------------|--------------------|--------------------|----------------------------------|-----------|-------|---------|
| | | | | | Num | Den | Percent |
| 1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2) | 72.0% | 71.0% | 70.0% | -2.8% | 151 | 214 | 70.6% |
| 1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services. | 71.6% | 69.2% | 75.3% | 5.1% | 206 | 267 | 77.2% |
| 1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services. | 60.0% | 57.7% | 59.6% | -0.7% | 753 | 1,234 | 61.0% |
| 2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy. | 25.0% | 23.5% | 26.2% | 4.9% | 113 | 532 | 21.2% |
| 2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum. | 67.6% | 67.5% | 68.6% | 1.5% | 142 | 209 | 67.9% |
| 3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1) | | | | | | | |
| 3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1) | | | | | | | |
| 4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were: | | | | | | | |
| a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1) | 67.3% | 66.6% | 66.8% | -0.7% | 313 | 450 | 69.6% |
| b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile) | 15.6% | 15.3% | 15.0% | -4.0% | 54 | 450 | 12.0% |
| c. Obese (BMI at or greater than the 95th percentile). | 15.2% | 15.6% | 14.9% | -2.1% | 63 | 450 | 14.0% |
| d. Underweight (BMI less than the 5th percentile). | 1.9% | 2.6% | 3.3% | 74.8% | 20 | 450 | 4.4% |
| 4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3) | | | | | | | |
| a. Recommended prenatal weight gain | 32.5% | 31.5% | 26.8% | -17.5% | 52 | 196 | 26.5% |
| b. Excessive prenatal weight gain | 47.9% | 47.1% | 51.8% | 8.2% | 99 | 196 | 50.5% |
| c. Inadequate prenatal weight gain | 19.6% | 21.4% | 21.4% | 9.0% | 45 | 196 | 23.0% |
| 5A. Percentage of women who participated in WIC who initiated breastfeeding. | 52.3% | 52.4% | 45.8% | -12.4% | 159 | 265 | 60.0% |
| 5B. Percentage of infants who participated in WIC who were: | | | | | | | |
| a. Breastfeeding at 6 weeks of age (1) | 31.9% | 33.3% | 28.1% | -11.8% | 66 | 274 | 24.1% |
| b. Breastfeeding at 6 months of age | 17.0% | 18.7% | 16.0% | -6.1% | 37 | 274 | 13.5% |

(1) Item appears also on other WCH Section Agreement Addenda. Data for 3A and 3B were not presented due to the limitation in recent trend datasets.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI >=30.0): Adapted from the National Academy of Sciences. Weight Gain During Pregnancy: Reexamining the Guidelines. Washington, DC : National Academy Press, 2009.

SFY 2014-2015 AGREEMENT ADDENDA DATA

WIC QUALITY ASSURANCE DELIVERABLES

Duplin County

| Process Outcome Objectives | 2004 to 2006 | 2007 to 2009 | 2010 to 2012 | % Change from 2004 to 2012 | 2012 Data | | |
|---|--------------------|--------------------|--------------------|----------------------------------|-----------|-------|---------|
| | | | | | Num | Den | Percent |
| 1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2) | 74.2% | 72.4% | 71.5% | -3.6% | 321 | 453 | 70.9% |
| 1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services. | 74.5% | 79.0% | 79.3% | 6.4% | 468 | 594 | 78.8% |
| 1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services. | 61.6% | 66.2% | 63.8% | 3.5% | 1,941 | 2,995 | 64.8% |
| 2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy. | 30.4% | 31.9% | 30.1% | -1.1% | 326 | 1,128 | 28.9% |
| 2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum. | 84.3% | 82.5% | 79.5% | -5.6% | 442 | 546 | 81.0% |
| 3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1) | | | | | | | |
| 3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1) | | | | | | | |
| 4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were: | | | | | | | |
| a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1) | 58.2% | 58.0% | 58.6% | 0.5% | 737 | 1,244 | 59.2% |
| b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile) | 13.6% | 17.4% | 16.0% | 17.9% | 185 | 1,244 | 14.9% |
| c. Obese (BMI at or greater than the 95th percentile). | 17.1% | 20.3% | 20.2% | 18.5% | 228 | 1,244 | 18.3% |
| d. Underweight (BMI less than the 5th percentile). | 11.1% | 4.3% | 5.2% | -53.2% | 94 | 1,244 | 7.6% |
| 4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3) | | | | | | | |
| a. Recommended prenatal weight gain | 34.1% | 32.5% | 25.0% | -26.7% | 96 | 379 | 25.3% |
| b. Excessive prenatal weight gain | 42.0% | 39.4% | 42.2% | 0.4% | 156 | 379 | 41.2% |
| c. Inadequate prenatal weight gain | 23.9% | 28.1% | 32.8% | 37.3% | 127 | 379 | 33.5% |
| 5A. Percentage of women who participated in WIC who initiated breastfeeding. | 55.5% | 50.9% | 52.4% | -5.5% | 424 | 719 | 59.0% |
| 5B. Percentage of infants who participated in WIC who were: | | | | | | | |
| a. Breastfeeding at 6 weeks of age (1) | 25.0% | 26.5% | 31.9% | 27.5% | 232 | 696 | 33.3% |
| b. Breastfeeding at 6 months of age | 12.5% | 12.6% | 18.0% | 44.4% | 110 | 696 | 15.8% |

(1) Item appears also on other WCH Section Agreement Addenda. Data for 3A and 3B were not presented due to the limitation in recent trend datasets.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI >=30.0): Adapted from the National Academy of Sciences. Weight Gain During Pregnancy: Reexamining the Guidelines. Washington, DC : National Academy Press, 2009.

SFY 2014-2015 AGREEMENT ADDENDA DATA

WIC QUALITY ASSURANCE DELIVERABLES

Durham County

| Process Outcome Objectives | 2004 to 2006 | 2007 to 2009 | 2010 to 2012 | % Change from 2004 to 2012 | 2012 Data | | |
|---|--------------------|--------------------|--------------------|----------------------------------|-----------|--------|---------|
| | | | | | Num | Den | Percent |
| 1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2) | 62.9% | 62.4% | 65.3% | 3.7% | 1,059 | 1,543 | 68.6% |
| 1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services. | 65.0% | 65.9% | 64.5% | -0.8% | 1,615 | 2,496 | 64.7% |
| 1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services. | 53.5% | 54.3% | 55.8% | 4.2% | 6,806 | 12,292 | 55.4% |
| 2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy. | 19.8% | 21.4% | 25.5% | 28.7% | 730 | 2,916 | 25.0% |
| 2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum. | 71.4% | 75.6% | 72.6% | 1.7% | 1,183 | 1,774 | 66.7% |
| 3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1) | | | | | | | |
| 3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1) | | | | | | | |
| 4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were: | | | | | | | |
| a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1) | 58.5% | 59.6% | 60.9% | 4.1% | 1,451 | 2,333 | 62.2% |
| b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile) | 15.3% | 15.7% | 16.2% | 6.4% | 351 | 2,333 | 15.0% |
| c. Obese (BMI at or greater than the 95th percentile). | 21.8% | 21.1% | 18.4% | -15.5% | 384 | 2,333 | 16.5% |
| d. Underweight (BMI less than the 5th percentile). | 4.5% | 3.6% | 4.5% | 0.3% | 147 | 2,333 | 6.3% |
| 4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3) | | | | | | | |
| a. Recommended prenatal weight gain | 36.5% | 33.6% | 28.2% | -22.9% | 394 | 1,384 | 28.5% |
| b. Excessive prenatal weight gain | 38.5% | 43.7% | 48.1% | 25.0% | 678 | 1,384 | 49.0% |
| c. Inadequate prenatal weight gain | 25.0% | 22.6% | 23.7% | -5.0% | 312 | 1,384 | 22.5% |
| 5A. Percentage of women who participated in WIC who initiated breastfeeding. | 59.4% | 58.6% | 66.4% | 11.7% | 1,568 | 2,355 | 66.6% |
| 5B. Percentage of infants who participated in WIC who were: | | | | | | | |
| a. Breastfeeding at 6 weeks of age (1) | 38.3% | 42.6% | 41.2% | 7.6% | 1,066 | 2,423 | 44.0% |
| b. Breastfeeding at 6 months of age | 20.4% | 23.9% | 24.3% | 18.9% | 632 | 2,423 | 26.1% |

(1) Item appears also on other WCH Section Agreement Addenda. Data for 3A and 3B were not presented due to the limitation in recent trend datasets.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI < 18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI ≥ 30.0): Adapted from the National Academy of Sciences. Weight Gain During Pregnancy: Reexamining the Guidelines. Washington, DC : National Academy Press, 2009.

SFY 2014-2015 AGREEMENT ADDENDA DATA

WIC QUALITY ASSURANCE DELIVERABLES

Edgecombe County

| Process Outcome Objectives | 2004 to 2006 | 2007 to 2009 | 2010 to 2012 | % Change from 2004 to 2012 | 2012 Data | | |
|---|--------------------|--------------------|--------------------|----------------------------------|-----------|-------|---------|
| | | | | | Num | Den | Percent |
| 1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2) | 83.4% | 84.4% | 85.0% | 2.0% | 476 | 547 | 87.0% |
| 1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services. | 74.4% | 73.0% | 73.2% | -1.7% | 444 | 600 | 74.0% |
| 1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services. | 56.4% | 53.7% | 53.0% | -6.0% | 1,662 | 3,044 | 54.6% |
| 2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy. | 27.6% | 28.4% | 24.2% | -12.3% | 298 | 1,224 | 24.3% |
| 2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum. | 71.9% | 69.6% | 69.0% | -4.1% | 342 | 515 | 66.4% |
| 3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1) | | | | | | | |
| 3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1) | | | | | | | |
| 4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were: | | | | | | | |
| a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1) | 66.0% | 66.6% | 67.9% | 2.9% | 252 | 367 | 68.7% |
| b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile) | 14.6% | 15.4% | 12.6% | -13.9% | 40 | 367 | 10.9% |
| c. Obese (BMI at or greater than the 95th percentile). | 16.2% | 14.6% | 13.5% | -16.6% | 51 | 367 | 13.9% |
| d. Underweight (BMI less than the 5th percentile). | 3.2% | 3.3% | 6.0% | 87.6% | 24 | 367 | 6.5% |
| 4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3) | | | | | | | |
| a. Recommended prenatal weight gain | 34.7% | 30.9% | 28.4% | -18.3% | 127 | 474 | 26.8% |
| b. Excessive prenatal weight gain | 38.2% | 42.2% | 44.9% | 17.3% | 228 | 474 | 48.1% |
| c. Inadequate prenatal weight gain | 27.0% | 26.9% | 26.8% | -1.0% | 119 | 474 | 25.1% |
| 5A. Percentage of women who participated in WIC who initiated breastfeeding. | 31.0% | 27.8% | 35.9% | 15.9% | 186 | 607 | 30.6% |
| 5B. Percentage of infants who participated in WIC who were: | | | | | | | |
| a. Breastfeeding at 6 weeks of age (1) | 16.8% | 15.1% | 12.4% | -26.0% | 75 | 601 | 12.5% |
| b. Breastfeeding at 6 months of age | 8.0% | 6.8% | 5.5% | -31.0% | 32 | 601 | 5.3% |

(1) Item appears also on other WCH Section Agreement Addenda. Data for 3A and 3B were not presented due to the limitation in recent trend datasets.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI >=30.0): Adapted from the National Academy of Sciences. Weight Gain During Pregnancy: Reexamining the Guidelines. Washington, DC : National Academy Press, 2009.

SFY 2014-2015 AGREEMENT ADDENDA DATA

WIC QUALITY ASSURANCE DELIVERABLES

Forsyth County

| Process Outcome Objectives | 2004 to 2006 | 2007 to 2009 | 2010 to 2012 | % Change from 2004 to 2012 | 2012 Data | | |
|---|--------------------|--------------------|--------------------|----------------------------------|-----------|--------|---------|
| | | | | | Num | Den | Percent |
| 1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2) | 66.0% | 67.0% | 72.5% | 9.8% | 1,662 | 2,251 | 73.8% |
| 1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services. | 66.5% | 70.3% | 65.6% | -1.4% | 1,997 | 3,105 | 64.3% |
| 1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services. | 51.9% | 55.1% | 58.4% | 12.6% | 8,953 | 15,423 | 58.0% |
| 2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy. | 23.3% | 34.0% | 32.4% | 39.4% | 1,391 | 4,605 | 30.2% |
| 2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum. | 81.8% | 83.6% | 80.5% | -1.5% | 2,099 | 2,609 | 80.5% |
| 3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1) | | | | | | | |
| 3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1) | | | | | | | |
| 4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were: | | | | | | | |
| a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1) | 65.9% | 64.7% | 65.9% | -0.1% | 3,186 | 4,805 | 66.3% |
| b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile) | 16.0% | 17.1% | 16.2% | 1.7% | 769 | 4,805 | 16.0% |
| c. Obese (BMI at or greater than the 95th percentile). | 14.8% | 15.2% | 14.2% | -4.1% | 659 | 4,805 | 13.7% |
| d. Underweight (BMI less than the 5th percentile). | 3.3% | 3.0% | 3.7% | 12.5% | 191 | 4,805 | 4.0% |
| 4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3) | | | | | | | |
| a. Recommended prenatal weight gain | 33.7% | 31.3% | 28.2% | -16.1% | 522 | 1,995 | 26.2% |
| b. Excessive prenatal weight gain | 42.2% | 45.1% | 47.9% | 13.6% | 969 | 1,995 | 48.6% |
| c. Inadequate prenatal weight gain | 24.2% | 23.7% | 23.9% | -1.3% | 504 | 1,995 | 25.3% |
| 5A. Percentage of women who participated in WIC who initiated breastfeeding. | 69.8% | 70.8% | 70.7% | 1.2% | 2,260 | 3,205 | 70.5% |
| 5B. Percentage of infants who participated in WIC who were: | | | | | | | |
| a. Breastfeeding at 6 weeks of age (1) | 42.2% | 45.8% | 47.9% | 13.4% | 1,520 | 3,084 | 49.3% |
| b. Breastfeeding at 6 months of age | 25.2% | 25.6% | 28.6% | 13.2% | 927 | 3,084 | 30.1% |

(1) Item appears also on other WCH Section Agreement Addenda. Data for 3A and 3B were not presented due to the limitation in recent trend datasets.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI >=30.0): Adapted from the National Academy of Sciences. Weight Gain During Pregnancy: Reexamining the Guidelines. Washington, DC : National Academy Press, 2009.

SFY 2014-2015 AGREEMENT ADDENDA DATA

WIC QUALITY ASSURANCE DELIVERABLES

Franklin County

| Process Outcome Objectives | 2004 to 2006 | 2007 to 2009 | 2010 to 2012 | % Change from 2004 to 2012 | 2012 Data | | |
|---|--------------------|--------------------|--------------------|----------------------------------|-----------|-------|---------|
| | | | | | Num | Den | Percent |
| 1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2) | 75.1% | 68.8% | 65.1% | -13.2% | 221 | 353 | 62.6% |
| 1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services. | 73.6% | 73.2% | 68.3% | -7.3% | 302 | 432 | 69.9% |
| 1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services. | 54.4% | 55.1% | 54.6% | 0.4% | 1,276 | 2,294 | 55.6% |
| 2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy. | 23.9% | 24.8% | 20.5% | -14.1% | 132 | 836 | 15.8% |
| 2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum. | 78.1% | 77.5% | 78.8% | 0.9% | 209 | 276 | 75.7% |
| 3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1) | | | | | | | |
| 3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1) | | | | | | | |
| 4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were: | | | | | | | |
| a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1) | 65.0% | 55.7% | 56.5% | -13.1% | 480 | 956 | 50.2% |
| b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile) | 16.7% | 20.5% | 16.6% | -0.4% | 146 | 956 | 15.3% |
| c. Obese (BMI at or greater than the 95th percentile). | 14.8% | 20.9% | 15.9% | 8.1% | 140 | 956 | 14.6% |
| d. Underweight (BMI less than the 5th percentile). | 3.6% | 2.8% | 11.0% | 203.7% | 190 | 956 | 19.9% |
| 4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3) | | | | | | | |
| a. Recommended prenatal weight gain | 32.0% | 32.2% | 29.4% | -7.9% | 67 | 252 | 26.6% |
| b. Excessive prenatal weight gain | 42.6% | 42.5% | 45.3% | 6.4% | 113 | 252 | 44.8% |
| c. Inadequate prenatal weight gain | 25.4% | 25.2% | 25.3% | -0.7% | 72 | 252 | 28.6% |
| 5A. Percentage of women who participated in WIC who initiated breastfeeding. | 41.0% | 40.9% | 45.1% | 10.1% | 170 | 379 | 44.9% |
| 5B. Percentage of infants who participated in WIC who were: | | | | | | | |
| a. Breastfeeding at 6 weeks of age (1) | 26.0% | 25.4% | 27.4% | 5.3% | 127 | 438 | 29.0% |
| b. Breastfeeding at 6 months of age | 13.0% | 12.8% | 13.4% | 2.8% | 59 | 438 | 13.5% |

(1) Item appears also on other WCH Section Agreement Addenda. Data for 3A and 3B were not presented due to the limitation in recent trend datasets.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI >=30.0): Adapted from the National Academy of Sciences. Weight Gain During Pregnancy: Reexamining the Guidelines. Washington, DC : National Academy Press, 2009.

SFY 2014-2015 AGREEMENT ADDENDA DATA

WIC QUALITY ASSURANCE DELIVERABLES

Gaston County

| Process Outcome Objectives | 2004 to 2006 | 2007 to 2009 | 2010 to 2012 | % Change from 2004 to 2012 | 2012 Data | | |
|---|--------------------|--------------------|--------------------|----------------------------------|-----------|-------|---------|
| | | | | | Num | Den | Percent |
| 1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2) | 71.0% | 72.8% | 70.7% | -0.4% | 1,073 | 1,499 | 71.6% |
| 1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services. | 74.6% | 74.2% | 75.8% | 1.6% | 1,323 | 1,757 | 75.3% |
| 1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services. | 48.9% | 50.9% | 51.8% | 5.8% | 4,791 | 8,981 | 53.3% |
| 2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy. | 19.5% | 27.3% | 32.8% | 67.7% | 775 | 2,598 | 29.8% |
| 2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum. | 74.1% | 82.0% | 82.2% | 10.9% | 1,121 | 1,379 | 81.3% |
| 3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1) | | | | | | | |
| 3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1) | | | | | | | |
| 4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were: | | | | | | | |
| a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1) | 66.0% | 67.5% | 66.6% | 0.8% | 1,520 | 2,238 | 67.9% |
| b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile) | 15.4% | 16.0% | 14.3% | -7.2% | 275 | 2,238 | 12.3% |
| c. Obese (BMI at or greater than the 95th percentile). | 14.9% | 13.2% | 14.0% | -5.9% | 313 | 2,238 | 14.0% |
| d. Underweight (BMI less than the 5th percentile). | 3.6% | 3.3% | 5.0% | 40.1% | 130 | 2,238 | 5.8% |
| 4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3) | | | | | | | |
| a. Recommended prenatal weight gain | 29.2% | 28.8% | 28.4% | -3.0% | 353 | 1,271 | 27.8% |
| b. Excessive prenatal weight gain | 46.3% | 44.0% | 45.6% | -1.6% | 596 | 1,271 | 46.9% |
| c. Inadequate prenatal weight gain | 24.4% | 27.2% | 26.1% | 6.7% | 322 | 1,271 | 25.3% |
| 5A. Percentage of women who participated in WIC who initiated breastfeeding. | 42.1% | 52.5% | 48.2% | 14.5% | 838 | 1,773 | 47.3% |
| 5B. Percentage of infants who participated in WIC who were: | | | | | | | |
| a. Breastfeeding at 6 weeks of age (1) | 23.0% | 24.7% | 27.8% | 20.8% | 560 | 1,846 | 30.3% |
| b. Breastfeeding at 6 months of age | 10.4% | 12.2% | 13.2% | 26.6% | 274 | 1,846 | 14.8% |

(1) Item appears also on other WCH Section Agreement Addenda. Data for 3A and 3B were not presented due to the limitation in recent trend datasets.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI >=30.0): Adapted from the National Academy of Sciences. Weight Gain During Pregnancy: Reexamining the Guidelines. Washington, DC : National Academy Press, 2009.

SFY 2014-2015 AGREEMENT ADDENDA DATA

WIC QUALITY ASSURANCE DELIVERABLES

Graham County

| Process Outcome Objectives | 2004 to 2006 | 2007 to 2009 | 2010 to 2012 | % Change from 2004 to 2012 | 2012 Data | | |
|---|--------------------|--------------------|--------------------|----------------------------------|-----------|-----|---------|
| | | | | | Num | Den | Percent |
| 1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2) | 80.3% | 77.0% | 87.0% | 8.3% | 69 | 76 | 90.8% |
| 1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services. | 73.8% | 80.4% | 74.4% | 0.9% | 64 | 82 | 78.0% |
| 1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services. | 67.7% | 71.9% | 76.1% | 12.4% | 295 | 400 | 73.8% |
| 2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy. | 64.6% | 63.3% | 61.0% | -5.5% | 88 | 152 | 57.9% |
| 2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum. | 78.4% | 77.0% | 78.5% | 0.2% | 61 | 74 | 82.4% |
| 3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1) | | | | | | | |
| 3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1) | | | | | | | |
| 4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were: | | | | | | | |
| a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1) | 65.4% | 64.9% | 57.7% | -11.8% | 139 | 245 | 56.7% |
| b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile) | 13.7% | 17.4% | 19.1% | 39.8% | 31 | 245 | 12.7% |
| c. Obese (BMI at or greater than the 95th percentile). | 17.1% | 15.3% | 19.0% | 10.9% | 63 | 245 | 25.7% |
| d. Underweight (BMI less than the 5th percentile). | 3.8% | 2.4% | 4.2% | 10.0% | 12 | 245 | 4.9% |
| 4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3) | | | | | | | |
| a. Recommended prenatal weight gain | 29.9% | 26.8% | 20.4% | -31.9% | 13 | 65 | 20.0% |
| b. Excessive prenatal weight gain | 51.9% | 57.9% | 55.0% | 6.0% | 32 | 65 | 49.2% |
| c. Inadequate prenatal weight gain | 18.2% | 15.3% | 24.6% | 35.5% | 20 | 65 | 30.8% |
| 5A. Percentage of women who participated in WIC who initiated breastfeeding. | 9.2% | 35.8% | 42.3% | 359.2% | 47 | 91 | 51.6% |
| 5B. Percentage of infants who participated in WIC who were: | | | | | | | |
| a. Breastfeeding at 6 weeks of age (1) | 22.8% | 18.8% | 18.9% | -16.9% | 10 | 72 | 13.9% |
| b. Breastfeeding at 6 months of age | 66.7% | 11.9% | 11.2% | -83.3% | 5 | 72 | 6.9% |

(1) Item appears also on other WCH Section Agreement Addenda. Data for 3A and 3B were not presented due to the limitation in recent trend datasets.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI < 18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI ≥ 30.0): Adapted from the National Academy of Sciences. Weight Gain During Pregnancy: Reexamining the Guidelines. Washington, DC : National Academy Press, 2009.

SFY 2014-2015 AGREEMENT ADDENDA DATA

WIC QUALITY ASSURANCE DELIVERABLES

Granville-Vance District

| Process Outcome Objectives | 2004 to 2006 | 2007 to 2009 | 2010 to 2012 | % Change from 2004 to 2012 | 2012 Data | | |
|---|--------------------|--------------------|--------------------|----------------------------------|-----------|-------|---------|
| | | | | | Num | Den | Percent |
| 1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2) | 72.2% | 75.9% | 72.9% | 0.9% | 520 | 714 | 72.8% |
| 1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services. | 76.6% | 75.3% | 75.7% | -1.2% | 674 | 901 | 74.8% |
| 1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services. | 56.7% | 58.3% | 57.5% | 1.3% | 2,701 | 4,638 | 58.2% |
| 2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy. | 31.0% | 33.2% | 27.7% | -10.5% | 430 | 1,554 | 27.7% |
| 2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum. | 80.2% | 75.2% | 76.7% | -4.4% | 556 | 707 | 78.6% |
| 3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1) | | | | | | | |
| 3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1) | | | | | | | |
| 4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were: | | | | | | | |
| a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1) | 65.3% | 65.3% | 66.0% | 1.0% | 891 | 1,343 | 66.3% |
| b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile) | 15.8% | 15.0% | 12.7% | -19.3% | 172 | 1,343 | 12.8% |
| c. Obese (BMI at or greater than the 95th percentile). | 14.7% | 14.0% | 13.8% | -6.0% | 181 | 1,343 | 13.5% |
| d. Underweight (BMI less than the 5th percentile). | 4.2% | 5.6% | 7.5% | 78.2% | 99 | 1,343 | 7.4% |
| 4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3) | | | | | | | |
| a. Recommended prenatal weight gain | 31.0% | 31.5% | 29.8% | -3.9% | 173 | 634 | 27.3% |
| b. Excessive prenatal weight gain | 41.1% | 42.9% | 43.5% | 5.7% | 295 | 634 | 46.5% |
| c. Inadequate prenatal weight gain | 27.8% | 25.6% | 26.7% | -4.0% | 166 | 634 | 26.2% |
| 5A. Percentage of women who participated in WIC who initiated breastfeeding. | 40.4% | 52.0% | 45.6% | 13.0% | 438 | 893 | 49.0% |
| 5B. Percentage of infants who participated in WIC who were: | | | | | | | |
| a. Breastfeeding at 6 weeks of age (1) | 28.0% | 38.1% | 35.4% | 26.2% | 337 | 928 | 36.3% |
| b. Breastfeeding at 6 months of age | 19.5% | 26.5% | 25.8% | 32.3% | 256 | 928 | 27.6% |

(1) Item appears also on other WCH Section Agreement Addenda. Data for 3A and 3B were not presented due to the limitation in recent trend datasets.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI >=30.0): Adapted from the National Academy of Sciences. Weight Gain During Pregnancy: Reexamining the Guidelines. Washington, DC : National Academy Press, 2009.

SFY 2014-2015 AGREEMENT ADDENDA DATA

WIC QUALITY ASSURANCE DELIVERABLES

Greene County

| Process Outcome Objectives | 2004 to 2006 | 2007 to 2009 | 2010 to 2012 | % Change from 2004 to 2012 | 2012 Data | | |
|---|--------------------|--------------------|--------------------|----------------------------------|-----------|-----|---------|
| | | | | | Num | Den | Percent |
| 1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2) | 79.8% | 82.3% | 77.6% | -2.7% | 101 | 138 | 73.2% |
| 1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services. | 78.5% | 76.9% | 73.6% | -6.3% | 129 | 167 | 77.2% |
| 1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services. | 63.4% | 61.8% | 63.6% | 0.4% | 599 | 923 | 64.9% |
| 2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy. | 32.5% | 29.3% | 30.4% | -6.4% | 142 | 456 | 31.1% |
| 2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum. | 78.7% | 85.2% | 83.2% | 5.8% | 136 | 171 | 79.5% |
| 3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1) | | | | | | | |
| 3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1) | | | | | | | |
| 4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were: | | | | | | | |
| a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1) | 65.9% | 65.5% | 60.1% | -8.7% | 312 | 525 | 59.4% |
| b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile) | 14.9% | 16.9% | 18.0% | 21.0% | 89 | 525 | 17.0% |
| c. Obese (BMI at or greater than the 95th percentile). | 15.4% | 15.1% | 16.7% | 9.0% | 82 | 525 | 15.6% |
| d. Underweight (BMI less than the 5th percentile). | 3.9% | 2.4% | 5.1% | 31.7% | 42 | 525 | 8.0% |
| 4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3) | | | | | | | |
| a. Recommended prenatal weight gain | 33.0% | 33.8% | 29.6% | -10.3% | 31 | 129 | 24.0% |
| b. Excessive prenatal weight gain | 38.8% | 39.8% | 45.3% | 16.7% | 65 | 129 | 50.4% |
| c. Inadequate prenatal weight gain | 28.2% | 26.4% | 25.1% | -10.9% | 33 | 129 | 25.6% |
| 5A. Percentage of women who participated in WIC who initiated breastfeeding. | 48.9% | 54.7% | 63.9% | 30.8% | 140 | 208 | 67.3% |
| 5B. Percentage of infants who participated in WIC who were: | | | | | | | |
| a. Breastfeeding at 6 weeks of age (1) | 32.9% | 31.7% | 36.8% | 12.1% | 83 | 217 | 38.2% |
| b. Breastfeeding at 6 months of age | 15.8% | 15.2% | 15.4% | -2.6% | 42 | 217 | 19.4% |

(1) Item appears also on other WCH Section Agreement Addenda. Data for 3A and 3B were not presented due to the limitation in recent trend datasets.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI >=30.0): Adapted from the National Academy of Sciences. *Weight Gain During Pregnancy: Reexamining the Guidelines*. Washington, DC : National Academy Press, 2009.

SFY 2014-2015 AGREEMENT ADDENDA DATA

WIC QUALITY ASSURANCE DELIVERABLES

Guilford County

| Process Outcome Objectives | 2004 to 2006 | 2007 to 2009 | 2010 to 2012 | % Change from 2004 to 2012 | 2012 Data | | |
|---|--------------------|--------------------|--------------------|----------------------------------|-----------|--------|---------|
| | | | | | Num | Den | Percent |
| 1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2) | 65.9% | 63.6% | 60.6% | -8.1% | 2,084 | 3,064 | 68.0% |
| 1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services. | 69.9% | 69.1% | 66.3% | -5.3% | 2,654 | 4,046 | 65.6% |
| 1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services. | 48.8% | 51.3% | 54.5% | 11.7% | 10,910 | 19,872 | 54.9% |
| 2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy. | 29.7% | 30.1% | 33.1% | 11.3% | 1,929 | 6,325 | 30.5% |
| 2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum. | 67.9% | 79.8% | 80.3% | 18.4% | 2,626 | 3,272 | 80.3% |
| 3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1) | | | | | | | |
| 3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1) | | | | | | | |
| 4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were: | | | | | | | |
| a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1) | 68.6% | 65.5% | 65.0% | -5.2% | 3,535 | 5,328 | 66.3% |
| b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile) | 14.5% | 16.8% | 15.2% | 5.4% | 783 | 5,328 | 14.7% |
| c. Obese (BMI at or greater than the 95th percentile). | 13.1% | 14.6% | 15.2% | 15.7% | 762 | 5,328 | 14.3% |
| d. Underweight (BMI less than the 5th percentile). | 3.8% | 3.1% | 4.6% | 18.3% | 248 | 5,328 | 4.7% |
| 4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3) | | | | | | | |
| a. Recommended prenatal weight gain | 31.5% | 30.7% | 28.5% | -9.5% | 754 | 2,661 | 28.3% |
| b. Excessive prenatal weight gain | 48.8% | 49.3% | 51.8% | 6.1% | 1,364 | 2,661 | 51.3% |
| c. Inadequate prenatal weight gain | 19.7% | 20.1% | 19.7% | 0.1% | 543 | 2,661 | 20.4% |
| 5A. Percentage of women who participated in WIC who initiated breastfeeding. | 59.9% | 64.6% | 63.6% | 6.1% | 2,788 | 4,269 | 65.3% |
| 5B. Percentage of infants who participated in WIC who were: | | | | | | | |
| a. Breastfeeding at 6 weeks of age (1) | 34.1% | 38.0% | 41.4% | 21.6% | 1,830 | 4,163 | 44.0% |
| b. Breastfeeding at 6 months of age | 18.3% | 22.8% | 24.7% | 34.9% | 1,041 | 4,163 | 25.0% |

(1) Item appears also on other WCH Section Agreement Addenda. Data for 3A and 3B were not presented due to the limitation in recent trend datasets.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI >=30.0): Adapted from the National Academy of Sciences. Weight Gain During Pregnancy: Reexamining the Guidelines. Washington, DC : National Academy Press, 2009.

SFY 2014-2015 AGREEMENT ADDENDA DATA

WIC QUALITY ASSURANCE DELIVERABLES

Halifax County

| Process Outcome Objectives | 2004 to 2006 | 2007 to 2009 | 2010 to 2012 | % Change from 2004 to 2012 | 2012 Data | | |
|---|--------------------|--------------------|--------------------|----------------------------------|-----------|-------|---------|
| | | | | | Num | Den | Percent |
| 1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2) | 81.3% | 82.5% | 83.1% | 2.3% | 386 | 452 | 85.4% |
| 1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services. | 74.5% | 73.4% | 70.6% | -5.1% | 326 | 472 | 69.1% |
| 1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services. | 60.6% | 61.2% | 56.5% | -6.8% | 1,386 | 2,450 | 56.6% |
| 2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy. | 29.0% | 28.5% | 27.5% | -5.3% | 293 | 992 | 29.5% |
| 2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum. | 71.7% | 71.4% | 66.7% | -7.0% | 308 | 457 | 67.4% |
| 3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1) | | | | | | | |
| 3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1) | | | | | | | |
| 4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were: | | | | | | | |
| a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1) | 66.5% | 65.8% | 64.0% | -3.8% | 570 | 904 | 63.1% |
| b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile) | 14.6% | 15.0% | 16.4% | 12.4% | 149 | 904 | 16.5% |
| c. Obese (BMI at or greater than the 95th percentile). | 14.7% | 14.9% | 14.2% | -3.5% | 121 | 904 | 13.4% |
| d. Underweight (BMI less than the 5th percentile). | 4.1% | 4.3% | 5.3% | 30.0% | 64 | 904 | 7.1% |
| 4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3) | | | | | | | |
| a. Recommended prenatal weight gain | 29.6% | 29.9% | 27.1% | -8.4% | 108 | 425 | 25.4% |
| b. Excessive prenatal weight gain | 47.0% | 46.8% | 45.0% | -4.2% | 200 | 425 | 47.1% |
| c. Inadequate prenatal weight gain | 23.4% | 23.4% | 27.8% | 19.0% | 117 | 425 | 27.5% |
| 5A. Percentage of women who participated in WIC who initiated breastfeeding. | 33.6% | 33.4% | 34.6% | 3.0% | 188 | 533 | 35.3% |
| 5B. Percentage of infants who participated in WIC who were: | | | | | | | |
| a. Breastfeeding at 6 weeks of age (1) | 13.0% | 13.7% | 14.3% | 9.6% | 96 | 568 | 16.9% |
| b. Breastfeeding at 6 months of age | 4.2% | 5.7% | 6.2% | 48.8% | 44 | 568 | 7.7% |

(1) Item appears also on other WCH Section Agreement Addenda. Data for 3A and 3B were not presented due to the limitation in recent trend datasets.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI >=30.0): Adapted from the National Academy of Sciences. Weight Gain During Pregnancy: Reexamining the Guidelines. Washington, DC : National Academy Press, 2009.

SFY 2014-2015 AGREEMENT ADDENDA DATA

WIC QUALITY ASSURANCE DELIVERABLES

Harnett County

| Process Outcome Objectives | 2004 to 2006 | 2007 to 2009 | 2010 to 2012 | % Change from 2004 to 2012 | 2012 Data | | |
|---|--------------------|--------------------|--------------------|----------------------------------|-----------|-------|---------|
| | | | | | Num | Den | Percent |
| 1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2) | 71.0% | 70.6% | 67.5% | -5.1% | 477 | 700 | 68.1% |
| 1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services. | 75.7% | 74.5% | 75.4% | -0.5% | 677 | 913 | 74.2% |
| 1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services. | 56.3% | 57.9% | 56.9% | 1.1% | 2,711 | 4,752 | 57.0% |
| 2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy. | 35.5% | 35.2% | 30.7% | -13.6% | 412 | 1,409 | 29.2% |
| 2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum. | 67.1% | 72.2% | 77.1% | 14.9% | 497 | 652 | 76.2% |
| 3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1) | | | | | | | |
| 3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1) | | | | | | | |
| 4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were: | | | | | | | |
| a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1) | 66.4% | 66.0% | 67.2% | 1.2% | 824 | 1,283 | 64.2% |
| b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile) | 16.5% | 15.4% | 13.2% | -19.8% | 198 | 1,283 | 15.4% |
| c. Obese (BMI at or greater than the 95th percentile). | 13.6% | 14.7% | 13.1% | -3.8% | 169 | 1,283 | 13.2% |
| d. Underweight (BMI less than the 5th percentile). | 3.5% | 3.9% | 6.5% | 86.8% | 92 | 1,283 | 7.2% |
| 4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3) | | | | | | | |
| a. Recommended prenatal weight gain | 31.6% | 30.9% | 29.5% | -6.5% | 194 | 644 | 30.1% |
| b. Excessive prenatal weight gain | 47.8% | 49.1% | 49.5% | 3.6% | 316 | 644 | 49.1% |
| c. Inadequate prenatal weight gain | 20.7% | 20.0% | 21.0% | 1.7% | 134 | 644 | 20.8% |
| 5A. Percentage of women who participated in WIC who initiated breastfeeding. | 41.5% | 49.6% | 48.9% | 17.7% | 458 | 907 | 50.5% |
| 5B. Percentage of infants who participated in WIC who were: | | | | | | | |
| a. Breastfeeding at 6 weeks of age (1) | 25.8% | 26.5% | 25.8% | -0.1% | 279 | 1,006 | 27.7% |
| b. Breastfeeding at 6 months of age | 11.4% | 11.3% | 12.0% | 4.8% | 135 | 1,006 | 13.4% |

(1) Item appears also on other WCH Section Agreement Addenda. Data for 3A and 3B were not presented due to the limitation in recent trend datasets.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI >=30.0): Adapted from the National Academy of Sciences. *Weight Gain During Pregnancy: Reexamining the Guidelines*. Washington, DC : National Academy Press, 2009.

SFY 2014-2015 AGREEMENT ADDENDA DATA

WIC QUALITY ASSURANCE DELIVERABLES

Haywood County

| Process Outcome Objectives | 2004 to 2006 | 2007 to 2009 | 2010 to 2012 | % Change from 2004 to 2012 | 2012 Data | | |
|---|--------------------|--------------------|--------------------|----------------------------------|-----------|-------|---------|
| | | | | | Num | Den | Percent |
| 1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2) | 73.8% | 70.0% | 78.7% | 6.6% | 281 | 345 | 81.4% |
| 1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services. | 69.6% | 71.9% | 71.4% | 2.6% | 295 | 414 | 71.3% |
| 1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services. | 50.8% | 50.9% | 59.5% | 17.0% | 1,298 | 2,084 | 62.3% |
| 2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy. | 45.7% | 44.4% | 47.1% | 3.0% | 296 | 721 | 41.1% |
| 2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum. | 61.0% | 69.6% | 77.2% | 26.7% | 294 | 377 | 78.0% |
| 3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1) | | | | | | | |
| 3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1) | | | | | | | |
| 4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were: | | | | | | | |
| a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1) | 63.7% | 66.1% | 63.9% | 0.3% | 361 | 557 | 64.8% |
| b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile) | 17.8% | 16.2% | 18.0% | 1.2% | 97 | 557 | 17.4% |
| c. Obese (BMI at or greater than the 95th percentile). | 14.7% | 13.7% | 14.3% | -2.5% | 78 | 557 | 14.0% |
| d. Underweight (BMI less than the 5th percentile). | 3.8% | 4.0% | 3.7% | -0.6% | 21 | 557 | 3.8% |
| 4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3) | | | | | | | |
| a. Recommended prenatal weight gain | 32.6% | 29.6% | 28.4% | -12.7% | 70 | 299 | 23.4% |
| b. Excessive prenatal weight gain | 48.8% | 51.5% | 50.6% | 3.5% | 164 | 299 | 54.8% |
| c. Inadequate prenatal weight gain | 18.6% | 18.9% | 21.0% | 12.9% | 65 | 299 | 21.7% |
| 5A. Percentage of women who participated in WIC who initiated breastfeeding. | 36.1% | 45.2% | 63.5% | 75.8% | 314 | 448 | 70.1% |
| 5B. Percentage of infants who participated in WIC who were: | | | | | | | |
| a. Breastfeeding at 6 weeks of age (1) | 25.4% | 25.3% | 19.2% | -24.3% | 91 | 454 | 20.0% |
| b. Breastfeeding at 6 months of age | 11.7% | 11.0% | 4.7% | -59.4% | 25 | 454 | 5.5% |

(1) Item appears also on other WCH Section Agreement Addenda. Data for 3A and 3B were not presented due to the limitation in recent trend datasets.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI >=30.0): Adapted from the National Academy of Sciences. Weight Gain During Pregnancy: Reexamining the Guidelines. Washington, DC : National Academy Press, 2009.

SFY 2014-2015 AGREEMENT ADDENDA DATA

WIC QUALITY ASSURANCE DELIVERABLES

Henderson County

| Process Outcome Objectives | 2004 to 2006 | 2007 to 2009 | 2010 to 2012 | % Change from 2004 to 2012 | 2012 Data | | |
|---|--------------------|--------------------|--------------------|----------------------------------|-----------|-------|---------|
| | | | | | Num | Den | Percent |
| 1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2) | 45.7% | 47.0% | 51.4% | 12.3% | 301 | 552 | 54.5% |
| 1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services. | 62.8% | 61.7% | 65.0% | 3.6% | 492 | 746 | 66.0% |
| 1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services. | 54.4% | 52.4% | 55.5% | 2.1% | 2,268 | 3,933 | 57.7% |
| 2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy. | 34.6% | 36.9% | 38.6% | 11.4% | 396 | 1,042 | 38.0% |
| 2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum. | 61.3% | 70.4% | 72.5% | 18.4% | 418 | 588 | 71.1% |
| 3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1) | | | | | | | |
| 3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1) | | | | | | | |
| 4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were: | | | | | | | |
| a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1) | 64.2% | 63.6% | 64.6% | 0.6% | 852 | 1,317 | 64.7% |
| b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile) | 17.8% | 18.3% | 17.6% | -1.1% | 212 | 1,317 | 16.1% |
| c. Obese (BMI at or greater than the 95th percentile). | 15.3% | 15.7% | 12.8% | -16.4% | 157 | 1,317 | 11.9% |
| d. Underweight (BMI less than the 5th percentile). | 2.7% | 2.4% | 5.0% | 84.3% | 96 | 1,317 | 7.3% |
| 4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3) | | | | | | | |
| a. Recommended prenatal weight gain | 36.1% | 33.1% | 30.5% | -15.6% | 114 | 413 | 27.6% |
| b. Excessive prenatal weight gain | 44.8% | 46.3% | 48.6% | 8.6% | 207 | 413 | 50.1% |
| c. Inadequate prenatal weight gain | 19.1% | 20.6% | 20.9% | 9.2% | 92 | 413 | 22.3% |
| 5A. Percentage of women who participated in WIC who initiated breastfeeding. | 70.5% | 80.3% | 80.7% | 14.5% | 596 | 766 | 77.8% |
| 5B. Percentage of infants who participated in WIC who were: | | | | | | | |
| a. Breastfeeding at 6 weeks of age (1) | 46.8% | 50.7% | 52.0% | 11.0% | 387 | 765 | 50.6% |
| b. Breastfeeding at 6 months of age | 27.8% | 30.0% | 32.2% | 15.8% | 253 | 765 | 33.1% |

(1) Item appears also on other WCH Section Agreement Addenda. Data for 3A and 3B were not presented due to the limitation in recent trend datasets.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI >=30.0): Adapted from the National Academy of Sciences. *Weight Gain During Pregnancy: Reexamining the Guidelines*. Washington, DC : National Academy Press, 2009.

SFY 2014-2015 AGREEMENT ADDENDA DATA

WIC QUALITY ASSURANCE DELIVERABLES

Hertford County

| Process Outcome Objectives | 2004 to 2006 | 2007 to 2009 | 2010 to 2012 | % Change from 2004 to 2012 | 2012 Data | | |
|---|--------------------|--------------------|--------------------|----------------------------------|-----------|-------|---------|
| | | | | | Num | Den | Percent |
| 1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2) | 79.1% | 78.4% | 75.0% | -5.3% | 130 | 171 | 76.0% |
| 1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services. | 74.8% | 71.8% | 69.4% | -7.3% | 135 | 181 | 74.6% |
| 1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services. | 55.9% | 55.0% | 55.8% | -0.2% | 594 | 1,036 | 57.3% |
| 2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy. | 28.2% | 27.0% | 24.5% | -13.2% | 98 | 422 | 23.2% |
| 2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum. | 77.7% | 86.1% | 85.2% | 9.6% | 170 | 196 | 86.7% |
| 3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1) | | | | | | | |
| 3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1) | | | | | | | |
| 4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were: | | | | | | | |
| a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1) | 65.6% | 68.8% | 63.6% | -3.1% | 256 | 414 | 61.8% |
| b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile) | 15.9% | 13.6% | 15.8% | -0.8% | 62 | 414 | 15.0% |
| c. Obese (BMI at or greater than the 95th percentile). | 14.6% | 12.1% | 14.1% | -3.5% | 63 | 414 | 15.2% |
| d. Underweight (BMI less than the 5th percentile). | 3.9% | 5.5% | 6.5% | 68.3% | 33 | 414 | 8.0% |
| 4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3) | | | | | | | |
| a. Recommended prenatal weight gain | 30.8% | 30.2% | 27.0% | -12.4% | 41 | 152 | 27.0% |
| b. Excessive prenatal weight gain | 50.8% | 45.2% | 42.2% | -17.0% | 63 | 152 | 41.4% |
| c. Inadequate prenatal weight gain | 18.4% | 24.6% | 30.8% | 67.8% | 48 | 152 | 31.6% |
| 5A. Percentage of women who participated in WIC who initiated breastfeeding. | 35.6% | 33.0% | 31.2% | -12.2% | 82 | 234 | 35.0% |
| 5B. Percentage of infants who participated in WIC who were: | | | | | | | |
| a. Breastfeeding at 6 weeks of age (1) | 15.7% | 16.8% | 19.2% | 21.8% | 42 | 229 | 18.3% |
| b. Breastfeeding at 6 months of age | 4.8% | 5.6% | 9.7% | 104.2% | 25 | 229 | 10.9% |

(1) Item appears also on other WCH Section Agreement Addenda. Data for 3A and 3B were not presented due to the limitation in recent trend datasets.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI >=30.0): Adapted from the National Academy of Sciences. Weight Gain During Pregnancy: Reexamining the Guidelines. Washington, DC : National Academy Press, 2009.

SFY 2014-2015 AGREEMENT ADDENDA DATA

WIC QUALITY ASSURANCE DELIVERABLES

Hoke County

| Process Outcome Objectives | 2004 to 2006 | 2007 to 2009 | 2010 to 2012 | % Change from 2004 to 2012 | 2012 Data | | |
|---|--------------------|--------------------|--------------------|----------------------------------|-----------|-------|---------|
| | | | | | Num | Den | Percent |
| 1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2) | 78.3% | 74.4% | 71.6% | -8.6% | 244 | 337 | 72.4% |
| 1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services. | 76.3% | 71.4% | 73.3% | -4.0% | 346 | 463 | 74.7% |
| 1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services. | 56.4% | 55.0% | 55.6% | -1.4% | 1,375 | 2,421 | 56.8% |
| 2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy. | 30.7% | 25.5% | 26.2% | -14.8% | 241 | 983 | 24.5% |
| 2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum. | 76.5% | 73.9% | 73.8% | -3.5% | 302 | 415 | 72.8% |
| 3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1) | | | | | | | |
| 3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1) | | | | | | | |
| 4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were: | | | | | | | |
| a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1) | 62.6% | 60.1% | 61.8% | -1.2% | 494 | 807 | 61.2% |
| b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile) | 14.9% | 18.5% | 16.4% | 9.9% | 115 | 807 | 14.3% |
| c. Obese (BMI at or greater than the 95th percentile). | 19.1% | 18.7% | 17.5% | -8.2% | 146 | 807 | 18.1% |
| d. Underweight (BMI less than the 5th percentile). | 3.4% | 2.7% | 4.2% | 26.2% | 52 | 807 | 6.4% |
| 4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3) | | | | | | | |
| a. Recommended prenatal weight gain | 31.8% | 32.1% | 28.5% | -10.4% | 102 | 351 | 29.1% |
| b. Excessive prenatal weight gain | 49.9% | 46.5% | 47.3% | -5.2% | 166 | 351 | 47.3% |
| c. Inadequate prenatal weight gain | 18.3% | 21.4% | 24.2% | 32.0% | 83 | 351 | 23.6% |
| 5A. Percentage of women who participated in WIC who initiated breastfeeding. | 47.8% | 46.7% | 53.1% | 11.1% | 305 | 572 | 53.3% |
| 5B. Percentage of infants who participated in WIC who were: | | | | | | | |
| a. Breastfeeding at 6 weeks of age (1) | 27.0% | 26.1% | 30.6% | 13.6% | 215 | 629 | 34.2% |
| b. Breastfeeding at 6 months of age | 13.5% | 13.8% | 18.0% | 33.9% | 119 | 629 | 18.9% |

(1) Item appears also on other WCH Section Agreement Addenda. Data for 3A and 3B were not presented due to the limitation in recent trend datasets.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI >=30.0): Adapted from the National Academy of Sciences. Weight Gain During Pregnancy: Reexamining the Guidelines. Washington, DC : National Academy Press, 2009.

SFY 2014-2015 AGREEMENT ADDENDA DATA

WIC QUALITY ASSURANCE DELIVERABLES

Hyde County

| Process Outcome Objectives | 2004 to 2006 | 2007 to 2009 | 2010 to 2012 | % Change from 2004 to 2012 | 2012 Data | | |
|---|--------------------|--------------------|--------------------|----------------------------------|-----------|-----|---------|
| | | | | | Num | Den | Percent |
| 1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2) | 75.3% | 71.2% | 79.1% | 5.1% | 23 | 26 | 88.5% |
| 1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services. | 65.9% | 75.9% | 75.6% | 14.7% | 27 | 36 | 75.0% |
| 1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services. | 66.5% | 70.6% | 69.8% | 5.0% | 134 | 196 | 68.4% |
| 2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy. | 38.0% | 33.5% | 28.3% | -25.5% | 28 | 91 | 30.8% |
| 2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum. | 70.0% | 71.8% | 59.0% | -15.7% | 18 | 34 | 52.9% |
| 3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1) | | | | | | | |
| 3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1) | | | | | | | |
| 4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were: | | | | | | | |
| a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1) | 63.5% | 66.2% | 64.3% | 1.2% | 45 | 71 | 63.4% |
| b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile) | 13.5% | 10.5% | 14.3% | 5.6% | 11 | 71 | 15.5% |
| c. Obese (BMI at or greater than the 95th percentile). | 18.0% | 12.7% | 16.8% | -6.8% | 10 | 71 | 14.1% |
| d. Underweight (BMI less than the 5th percentile). | 4.9% | 10.5% | 4.6% | -6.0% | 5 | 71 | 7.0% |
| 4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3) | | | | | | | |
| a. Recommended prenatal weight gain | 21.3% | 34.7% | 31.0% | 45.4% | 6 | 28 | 21.4% |
| b. Excessive prenatal weight gain | 56.2% | 44.4% | 46.0% | -18.2% | 9 | 28 | 32.1% |
| c. Inadequate prenatal weight gain | 22.5% | 20.8% | 23.0% | 2.3% | 13 | 28 | 46.4% |
| 5A. Percentage of women who participated in WIC who initiated breastfeeding. | 50.0% | 45.7% | 37.8% | -24.4% | 14 | 41 | 34.1% |
| 5B. Percentage of infants who participated in WIC who were: | | | | | | | |
| a. Breastfeeding at 6 weeks of age (1) | 32.9% | 34.0% | 25.7% | -21.8% | 14 | 49 | 28.6% |
| b. Breastfeeding at 6 months of age | 18.6% | 17.0% | 23.5% | 26.3% | 14 | 49 | 28.6% |

(1) Item appears also on other WCH Section Agreement Addenda. Data for 3A and 3B were not presented due to the limitation in recent trend datasets.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI >=30.0): Adapted from the National Academy of Sciences. Weight Gain During Pregnancy: Reexamining the Guidelines. Washington, DC : National Academy Press, 2009.

SFY 2014-2015 AGREEMENT ADDENDA DATA

WIC QUALITY ASSURANCE DELIVERABLES

Iredell County

| Process Outcome Objectives | 2004 to 2006 | 2007 to 2009 | 2010 to 2012 | % Change from 2004 to 2012 | 2012 Data | | |
|---|--------------------|--------------------|--------------------|----------------------------------|-----------|-------|---------|
| | | | | | Num | Den | Percent |
| 1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2) | 73.0% | 77.3% | 76.2% | 4.4% | 665 | 861 | 77.2% |
| 1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services. | 71.2% | 73.8% | 75.4% | 5.8% | 781 | 1,021 | 76.5% |
| 1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services. | 55.1% | 58.5% | 62.4% | 13.1% | 3,516 | 5,369 | 65.5% |
| 2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy. | 25.2% | 31.8% | 30.8% | 22.1% | 523 | 1,657 | 31.6% |
| 2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum. | 54.7% | 63.1% | 81.5% | 49.1% | 730 | 892 | 81.8% |
| 3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1) | | | | | | | |
| 3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1) | | | | | | | |
| 4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were: | | | | | | | |
| a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1) | 69.6% | 67.0% | 63.7% | -8.5% | 953 | 1,494 | 63.8% |
| b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile) | 14.6% | 14.9% | 16.9% | 15.8% | 235 | 1,494 | 15.7% |
| c. Obese (BMI at or greater than the 95th percentile). | 12.4% | 15.4% | 15.5% | 24.8% | 222 | 1,494 | 14.9% |
| d. Underweight (BMI less than the 5th percentile). | 3.3% | 2.7% | 3.9% | 15.7% | 84 | 1,494 | 5.6% |
| 4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3) | | | | | | | |
| a. Recommended prenatal weight gain | 31.6% | 30.0% | 32.0% | 1.3% | 224 | 731 | 30.6% |
| b. Excessive prenatal weight gain | 49.5% | 49.2% | 45.9% | -7.2% | 346 | 731 | 47.3% |
| c. Inadequate prenatal weight gain | 18.9% | 20.7% | 22.0% | 16.8% | 161 | 731 | 22.0% |
| 5A. Percentage of women who participated in WIC who initiated breastfeeding. | 50.8% | 57.1% | 59.5% | 17.0% | 671 | 1,132 | 59.3% |
| 5B. Percentage of infants who participated in WIC who were: | | | | | | | |
| a. Breastfeeding at 6 weeks of age (1) | 27.5% | 31.5% | 32.2% | 17.0% | 343 | 1,086 | 31.6% |
| b. Breastfeeding at 6 months of age | 14.7% | 16.8% | 18.2% | 23.8% | 185 | 1,086 | 17.0% |

(1) Item appears also on other WCH Section Agreement Addenda. Data for 3A and 3B were not presented due to the limitation in recent trend datasets.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI >=30.0): Adapted from the National Academy of Sciences. Weight Gain During Pregnancy: Reexamining the Guidelines. Washington, DC : National Academy Press, 2009.

SFY 2014-2015 AGREEMENT ADDENDA DATA

WIC QUALITY ASSURANCE DELIVERABLES

Jackson County

| Process Outcome Objectives | 2004 to 2006 | 2007 to 2009 | 2010 to 2012 | % Change from 2004 to 2012 | 2012 Data | | |
|---|--------------------|--------------------|--------------------|----------------------------------|-----------|-------|---------|
| | | | | | Num | Den | Percent |
| 1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2) | 81.5% | 80.2% | 78.1% | -4.2% | 200 | 249 | 80.3% |
| 1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services. | 66.5% | 70.2% | 72.3% | 8.8% | 214 | 301 | 71.1% |
| 1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services. | 65.0% | 64.2% | 66.0% | 1.5% | 1,009 | 1,514 | 66.6% |
| 2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy. | 46.9% | 52.0% | 42.6% | -9.1% | 196 | 515 | 38.1% |
| 2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum. | 62.7% | 65.5% | 62.2% | -0.8% | 156 | 256 | 60.9% |
| 3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1) | | | | | | | |
| 3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1) | | | | | | | |
| 4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were: | | | | | | | |
| a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1) | 60.5% | 59.4% | 67.0% | 10.8% | 277 | 398 | 69.6% |
| b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile) | 18.8% | 18.4% | 15.1% | -19.5% | 56 | 398 | 14.1% |
| c. Obese (BMI at or greater than the 95th percentile). | 17.8% | 19.0% | 14.4% | -19.2% | 52 | 398 | 13.1% |
| d. Underweight (BMI less than the 5th percentile). | 2.9% | 3.2% | 3.4% | 19.4% | 13 | 398 | 3.3% |
| 4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3) | | | | | | | |
| a. Recommended prenatal weight gain | 29.7% | 32.4% | 30.1% | 1.6% | 74 | 236 | 31.4% |
| b. Excessive prenatal weight gain | 53.5% | 51.5% | 51.5% | -3.7% | 124 | 236 | 52.5% |
| c. Inadequate prenatal weight gain | 16.9% | 16.1% | 18.4% | 9.1% | 38 | 236 | 16.1% |
| 5A. Percentage of women who participated in WIC who initiated breastfeeding. | 49.1% | 48.8% | 53.9% | 9.7% | 170 | 304 | 55.9% |
| 5B. Percentage of infants who participated in WIC who were: | | | | | | | |
| a. Breastfeeding at 6 weeks of age (1) | 36.3% | 33.1% | 40.2% | 10.8% | 106 | 259 | 40.9% |
| b. Breastfeeding at 6 months of age | 18.4% | 18.8% | 24.4% | 32.7% | 57 | 259 | 22.0% |

(1) Item appears also on other WCH Section Agreement Addenda. Data for 3A and 3B were not presented due to the limitation in recent trend datasets.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI >=30.0): Adapted from the National Academy of Sciences. Weight Gain During Pregnancy: Reexamining the Guidelines. Washington, DC : National Academy Press, 2009.

SFY 2014-2015 AGREEMENT ADDENDA DATA

WIC QUALITY ASSURANCE DELIVERABLES

Johnston County

| Process Outcome Objectives | 2004 to 2006 | 2007 to 2009 | 2010 to 2012 | % Change from 2004 to 2012 | 2012 Data | | |
|---|--------------------|--------------------|--------------------|----------------------------------|-----------|-------|---------|
| | | | | | Num | Den | Percent |
| 1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2) | 61.6% | 54.7% | 63.9% | 3.8% | 713 | 1,044 | 68.3% |
| 1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services. | 68.9% | 70.0% | 70.5% | 2.4% | 990 | 1,417 | 69.9% |
| 1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services. | 48.7% | 49.4% | 52.3% | 7.2% | 4,001 | 7,350 | 54.4% |
| 2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy. | 24.1% | 9.2% | 26.2% | 8.6% | 494 | 1,531 | 32.3% |
| 2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum. | 58.6% | 73.9% | 77.8% | 32.7% | 739 | 934 | 79.1% |
| 3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1) | | | | | | | |
| 3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1) | | | | | | | |
| 4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were: | | | | | | | |
| a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1) | 60.9% | 62.2% | 64.9% | 6.7% | 893 | 1,407 | 63.5% |
| b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile) | 17.2% | 17.0% | 14.5% | -16.1% | 215 | 1,407 | 15.3% |
| c. Obese (BMI at or greater than the 95th percentile). | 18.4% | 17.7% | 15.8% | -14.0% | 254 | 1,407 | 18.1% |
| d. Underweight (BMI less than the 5th percentile). | 3.5% | 3.1% | 4.8% | 36.4% | 45 | 1,407 | 3.2% |
| 4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3) | | | | | | | |
| a. Recommended prenatal weight gain | 32.6% | 38.6% | 28.8% | -11.5% | 230 | 804 | 28.6% |
| b. Excessive prenatal weight gain | 44.8% | 44.5% | 50.6% | 12.9% | 410 | 804 | 51.0% |
| c. Inadequate prenatal weight gain | 22.6% | 16.9% | 20.6% | -8.9% | 164 | 804 | 20.4% |
| 5A. Percentage of women who participated in WIC who initiated breastfeeding. | 38.1% | 50.3% | 52.5% | 37.8% | 730 | 1,228 | 59.4% |
| 5B. Percentage of infants who participated in WIC who were: | | | | | | | |
| a. Breastfeeding at 6 weeks of age (1) | 22.0% | 31.1% | 29.4% | 33.5% | 429 | 1,343 | 31.9% |
| b. Breastfeeding at 6 months of age | 10.8% | 17.1% | 17.2% | 59.7% | 275 | 1,343 | 20.5% |

(1) Item appears also on other WCH Section Agreement Addenda. Data for 3A and 3B were not presented due to the limitation in recent trend datasets.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI >=30.0): Adapted from the National Academy of Sciences. Weight Gain During Pregnancy: Reexamining the Guidelines. Washington, DC : National Academy Press, 2009.

SFY 2014-2015 AGREEMENT ADDENDA DATA

WIC QUALITY ASSURANCE DELIVERABLES

Jones County

| Process Outcome Objectives | 2004 to 2006 | 2007 to 2009 | 2010 to 2012 | % Change from 2004 to 2012 | 2012 Data | | |
|---|--------------------|--------------------|--------------------|----------------------------------|-----------|-----|---------|
| | | | | | Num | Den | Percent |
| 1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2) | 86.6% | 79.7% | 89.6% | 3.5% | 46 | 50 | 92.0% |
| 1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services. | 79.8% | 81.8% | 77.3% | -3.2% | 51 | 65 | 78.5% |
| 1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services. | 66.7% | 72.3% | 72.3% | 8.4% | 257 | 357 | 72.0% |
| 2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy. | 44.3% | 49.2% | 41.0% | -7.4% | 58 | 177 | 32.8% |
| 2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum. | 76.9% | 75.1% | 69.9% | -9.1% | 52 | 78 | 66.7% |
| 3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1) | | | | | | | |
| 3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1) | | | | | | | |
| 4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were: | | | | | | | |
| a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1) | 70.7% | 71.1% | 64.1% | -9.3% | 107 | 167 | 64.1% |
| b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile) | 14.9% | 14.4% | 15.9% | 7.2% | 26 | 167 | 15.6% |
| c. Obese (BMI at or greater than the 95th percentile). | 11.3% | 11.2% | 13.5% | 20.4% | 25 | 167 | 15.0% |
| d. Underweight (BMI less than the 5th percentile). | 3.2% | 3.3% | 6.4% | 100.2% | 9 | 167 | 5.4% |
| 4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3) | | | | | | | |
| a. Recommended prenatal weight gain | 35.5% | 33.2% | 22.0% | -37.9% | 16 | 65 | 24.6% |
| b. Excessive prenatal weight gain | 46.4% | 49.1% | 46.7% | 0.7% | 26 | 65 | 40.0% |
| c. Inadequate prenatal weight gain | 18.2% | 17.8% | 31.3% | 72.0% | 23 | 65 | 35.4% |
| 5A. Percentage of women who participated in WIC who initiated breastfeeding. | 48.6% | 48.7% | 44.7% | -8.0% | 45 | 93 | 48.4% |
| 5B. Percentage of infants who participated in WIC who were: | | | | | | | |
| a. Breastfeeding at 6 weeks of age (1) | 29.4% | 32.3% | 34.6% | 17.6% | 33 | 108 | 30.6% |
| b. Breastfeeding at 6 months of age | 14.7% | 15.1% | 19.6% | 33.5% | 18 | 108 | 16.7% |

(1) Item appears also on other WCH Section Agreement Addenda. Data for 3A and 3B were not presented due to the limitation in recent trend datasets.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI >=30.0): Adapted from the National Academy of Sciences. Weight Gain During Pregnancy: Reexamining the Guidelines. Washington, DC : National Academy Press, 2009.

SFY 2014-2015 AGREEMENT ADDENDA DATA

WIC QUALITY ASSURANCE DELIVERABLES

Lee County

| Process Outcome Objectives | 2004 to 2006 | 2007 to 2009 | 2010 to 2012 | % Change from 2004 to 2012 | 2012 Data | | |
|---|--------------------|--------------------|--------------------|----------------------------------|-----------|-------|---------|
| | | | | | Num | Den | Percent |
| 1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2) | 67.6% | 66.3% | 64.4% | -4.7% | 330 | 415 | 79.5% |
| 1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services. | 76.7% | 77.7% | 78.7% | 2.5% | 473 | 593 | 79.8% |
| 1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services. | 58.7% | 61.9% | 63.2% | 7.6% | 2,094 | 3,214 | 65.2% |
| 2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy. | 29.1% | 20.8% | 23.8% | -18.4% | 282 | 1,031 | 27.4% |
| 2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum. | 79.3% | 70.0% | 72.5% | -8.7% | 366 | 477 | 76.7% |
| 3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1) | | | | | | | |
| 3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1) | | | | | | | |
| 4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were: | | | | | | | |
| a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1) | 68.2% | 65.3% | 64.9% | -4.8% | 624 | 928 | 67.2% |
| b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile) | 14.8% | 15.7% | 15.6% | 5.8% | 136 | 928 | 14.7% |
| c. Obese (BMI at or greater than the 95th percentile). | 14.4% | 16.8% | 16.3% | 12.9% | 135 | 928 | 14.5% |
| d. Underweight (BMI less than the 5th percentile). | 2.6% | 2.2% | 3.2% | 21.7% | 33 | 928 | 3.6% |
| 4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3) | | | | | | | |
| a. Recommended prenatal weight gain | 29.8% | 32.3% | 30.2% | 1.2% | 101 | 377 | 26.8% |
| b. Excessive prenatal weight gain | 45.2% | 46.3% | 48.0% | 6.1% | 204 | 377 | 54.1% |
| c. Inadequate prenatal weight gain | 25.0% | 21.4% | 21.8% | -12.5% | 72 | 377 | 19.1% |
| 5A. Percentage of women who participated in WIC who initiated breastfeeding. | 51.2% | 54.9% | 51.0% | -0.4% | 306 | 578 | 52.9% |
| 5B. Percentage of infants who participated in WIC who were: | | | | | | | |
| a. Breastfeeding at 6 weeks of age (1) | 32.8% | 31.5% | 30.3% | -7.5% | 204 | 595 | 34.3% |
| b. Breastfeeding at 6 months of age | 16.1% | 17.4% | 16.4% | 2.1% | 106 | 595 | 17.8% |

(1) Item appears also on other WCH Section Agreement Addenda. Data for 3A and 3B were not presented due to the limitation in recent trend datasets.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI >=30.0): Adapted from the National Academy of Sciences. Weight Gain During Pregnancy: Reexamining the Guidelines. Washington, DC : National Academy Press, 2009.

SFY 2014-2015 AGREEMENT ADDENDA DATA

WIC QUALITY ASSURANCE DELIVERABLES

Lenoir County

| Process Outcome Objectives | 2004 to 2006 | 2007 to 2009 | 2010 to 2012 | % Change from 2004 to 2012 | 2012 Data | | |
|---|--------------------|--------------------|--------------------|----------------------------------|-----------|-------|---------|
| | | | | | Num | Den | Percent |
| 1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2) | 72.3% | 72.5% | 72.8% | 0.7% | 339 | 457 | 74.2% |
| 1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services. | 75.6% | 74.1% | 74.1% | -2.0% | 411 | 552 | 74.5% |
| 1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services. | 53.0% | 52.2% | 52.0% | -2.0% | 1,571 | 2,887 | 54.4% |
| 2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy. | 33.0% | 26.5% | 26.7% | -19.2% | 300 | 1,100 | 27.3% |
| 2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum. | 40.1% | 67.2% | 80.2% | 100.1% | 384 | 447 | 85.9% |
| 3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1) | | | | | | | |
| 3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1) | | | | | | | |
| 4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were: | | | | | | | |
| a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1) | 70.0% | 69.4% | 65.2% | -6.8% | 508 | 766 | 66.3% |
| b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile) | 10.3% | 10.8% | 15.1% | 46.0% | 108 | 766 | 14.1% |
| c. Obese (BMI at or greater than the 95th percentile). | 10.1% | 10.5% | 13.4% | 32.6% | 93 | 766 | 12.1% |
| d. Underweight (BMI less than the 5th percentile). | 9.6% | 9.3% | 6.3% | -34.5% | 57 | 766 | 7.4% |
| 4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3) | | | | | | | |
| a. Recommended prenatal weight gain | 29.2% | 27.0% | 26.4% | -9.5% | 91 | 394 | 23.1% |
| b. Excessive prenatal weight gain | 48.5% | 45.9% | 49.8% | 2.7% | 204 | 394 | 51.8% |
| c. Inadequate prenatal weight gain | 22.3% | 27.1% | 23.7% | 6.7% | 99 | 394 | 25.1% |
| 5A. Percentage of women who participated in WIC who initiated breastfeeding. | 33.5% | 43.2% | 47.9% | 43.0% | 257 | 565 | 45.5% |
| 5B. Percentage of infants who participated in WIC who were: | | | | | | | |
| a. Breastfeeding at 6 weeks of age (1) | 19.1% | 21.2% | 23.6% | 23.9% | 158 | 598 | 26.4% |
| b. Breastfeeding at 6 months of age | 8.7% | 10.1% | 10.9% | 24.8% | 76 | 598 | 12.7% |

(1) Item appears also on other WCH Section Agreement Addenda. Data for 3A and 3B were not presented due to the limitation in recent trend datasets.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI >=30.0): Adapted from the National Academy of Sciences. Weight Gain During Pregnancy: Reexamining the Guidelines. Washington, DC : National Academy Press, 2009.

SFY 2014-2015 AGREEMENT ADDENDA DATA

WIC QUALITY ASSURANCE DELIVERABLES

Lincoln County

| Process Outcome Objectives | 2004 to 2006 | 2007 to 2009 | 2010 to 2012 | % Change from 2004 to 2012 | 2012 Data | | |
|---|--------------------|--------------------|--------------------|----------------------------------|-----------|-------|---------|
| | | | | | Num | Den | Percent |
| 1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2) | 75.5% | 73.4% | 73.7% | -2.5% | 315 | 388 | 81.2% |
| 1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services. | 76.1% | 75.4% | 75.4% | -0.9% | 377 | 483 | 78.1% |
| 1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services. | 55.2% | 61.6% | 62.1% | 12.5% | 1,663 | 2,592 | 64.2% |
| 2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy. | 37.1% | 38.0% | 36.2% | -2.4% | 345 | 977 | 35.3% |
| 2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum. | 73.3% | 73.1% | 79.6% | 8.5% | 385 | 485 | 79.4% |
| 3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1) | | | | | | | |
| 3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1) | | | | | | | |
| 4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were: | | | | | | | |
| a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1) | 70.2% | 72.5% | 69.6% | -0.9% | 578 | 841 | 68.7% |
| b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile) | 7.7% | 10.1% | 10.6% | 38.1% | 89 | 841 | 10.6% |
| c. Obese (BMI at or greater than the 95th percentile). | 7.7% | 10.1% | 9.8% | 27.4% | 83 | 841 | 9.9% |
| d. Underweight (BMI less than the 5th percentile). | 14.4% | 7.3% | 10.0% | -30.5% | 91 | 841 | 10.8% |
| 4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3) | | | | | | | |
| a. Recommended prenatal weight gain | 29.4% | 30.9% | 29.5% | 0.5% | 106 | 384 | 27.6% |
| b. Excessive prenatal weight gain | 49.6% | 47.4% | 48.2% | -2.9% | 198 | 384 | 51.6% |
| c. Inadequate prenatal weight gain | 21.0% | 21.7% | 22.3% | 6.3% | 80 | 384 | 20.8% |
| 5A. Percentage of women who participated in WIC who initiated breastfeeding. | 51.0% | 53.5% | 60.3% | 18.2% | 354 | 589 | 60.1% |
| 5B. Percentage of infants who participated in WIC who were: | | | | | | | |
| a. Breastfeeding at 6 weeks of age (1) | 27.7% | 27.2% | 33.3% | 20.4% | 183 | 562 | 32.6% |
| b. Breastfeeding at 6 months of age | 11.1% | 11.3% | 15.2% | 37.4% | 80 | 562 | 14.2% |

(1) Item appears also on other WCH Section Agreement Addenda. Data for 3A and 3B were not presented due to the limitation in recent trend datasets.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI >=30.0): Adapted from the National Academy of Sciences. Weight Gain During Pregnancy: Reexamining the Guidelines. Washington, DC : National Academy Press, 2009.

SFY 2014-2015 AGREEMENT ADDENDA DATA

WIC QUALITY ASSURANCE DELIVERABLES

Macon County

| Process Outcome Objectives | 2004 to 2006 | 2007 to 2009 | 2010 to 2012 | % Change from 2004 to 2012 | 2012 Data | | |
|---|--------------------|--------------------|--------------------|----------------------------------|-----------|-------|---------|
| | | | | | Num | Den | Percent |
| 1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2) | 77.4% | 61.9% | 68.2% | -11.8% | 176 | 220 | 80.0% |
| 1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services. | 71.7% | 72.2% | 73.7% | 2.8% | 210 | 280 | 75.0% |
| 1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services. | 59.0% | 60.5% | 64.2% | 8.9% | 905 | 1,357 | 66.7% |
| 2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy. | 46.8% | 39.8% | 38.3% | -18.2% | 195 | 511 | 38.2% |
| 2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum. | 50.8% | 77.3% | 72.4% | 42.6% | 180 | 261 | 69.0% |
| 3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1) | | | | | | | |
| 3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1) | | | | | | | |
| 4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were: | | | | | | | |
| a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1) | 63.3% | 67.6% | 59.1% | -6.7% | 248 | 419 | 59.2% |
| b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile) | 18.1% | 16.9% | 17.2% | -4.8% | 70 | 419 | 16.7% |
| c. Obese (BMI at or greater than the 95th percentile). | 16.3% | 13.5% | 20.2% | 23.6% | 86 | 419 | 20.5% |
| d. Underweight (BMI less than the 5th percentile). | 2.2% | 2.0% | 3.5% | 56.8% | 15 | 419 | 3.6% |
| 4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3) | | | | | | | |
| a. Recommended prenatal weight gain | 34.0% | 37.7% | 28.9% | -14.9% | 54 | 202 | 26.7% |
| b. Excessive prenatal weight gain | 45.9% | 44.2% | 51.6% | 12.5% | 112 | 202 | 55.4% |
| c. Inadequate prenatal weight gain | 20.1% | 18.1% | 19.5% | -3.1% | 36 | 202 | 17.8% |
| 5A. Percentage of women who participated in WIC who initiated breastfeeding. | 65.1% | 69.4% | 62.1% | -4.6% | 179 | 312 | 57.4% |
| 5B. Percentage of infants who participated in WIC who were: | | | | | | | |
| a. Breastfeeding at 6 weeks of age (1) | 42.8% | 39.4% | 42.7% | -0.3% | 139 | 290 | 47.9% |
| b. Breastfeeding at 6 months of age | 22.5% | 23.8% | 29.1% | 29.0% | 96 | 290 | 33.1% |

(1) Item appears also on other WCH Section Agreement Addenda. Data for 3A and 3B were not presented due to the limitation in recent trend datasets.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI >=30.0): Adapted from the National Academy of Sciences. Weight Gain During Pregnancy: Reexamining the Guidelines. Washington, DC : National Academy Press, 2009.

SFY 2014-2015 AGREEMENT ADDENDA DATA

WIC QUALITY ASSURANCE DELIVERABLES

Madison County

| Process Outcome Objectives | 2004 to 2006 | 2007 to 2009 | 2010 to 2012 | % Change from 2004 to 2012 | 2012 Data | | |
|---|--------------------|--------------------|--------------------|----------------------------------|-----------|-----|---------|
| | | | | | Num | Den | Percent |
| 1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2) | 81.6% | 70.1% | 71.3% | -12.6% | 100 | 132 | 75.8% |
| 1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services. | 67.4% | 68.2% | 63.5% | -5.7% | 102 | 152 | 67.1% |
| 1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services. | 58.9% | 55.2% | 56.0% | -5.0% | 423 | 747 | 56.6% |
| 2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy. | 46.0% | 43.5% | 41.5% | -9.8% | 116 | 316 | 36.7% |
| 2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum. | 69.2% | 71.1% | 71.8% | 3.8% | 112 | 155 | 72.3% |
| 3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1) | | | | | | | |
| 3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1) | | | | | | | |
| 4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were: | | | | | | | |
| a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1) | 61.3% | 67.6% | 64.2% | 4.7% | 144 | 222 | 64.9% |
| b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile) | 16.3% | 15.0% | 13.4% | -17.4% | 27 | 222 | 12.2% |
| c. Obese (BMI at or greater than the 95th percentile). | 19.5% | 13.2% | 14.0% | -28.0% | 35 | 222 | 15.8% |
| d. Underweight (BMI less than the 5th percentile). | 3.0% | 4.2% | 8.4% | 182.7% | 16 | 222 | 7.2% |
| 4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3) | | | | | | | |
| a. Recommended prenatal weight gain | 27.8% | 34.0% | 29.7% | 7.0% | 41 | 118 | 34.7% |
| b. Excessive prenatal weight gain | 50.3% | 48.3% | 47.8% | -4.9% | 51 | 118 | 43.2% |
| c. Inadequate prenatal weight gain | 21.9% | 17.8% | 22.4% | 2.4% | 26 | 118 | 22.0% |
| 5A. Percentage of women who participated in WIC who initiated breastfeeding. | 39.4% | 58.1% | 65.7% | 66.5% | 125 | 181 | 69.1% |
| 5B. Percentage of infants who participated in WIC who were: | | | | | | | |
| a. Breastfeeding at 6 weeks of age (1) | 34.6% | 46.0% | 42.1% | 21.5% | 77 | 190 | 40.5% |
| b. Breastfeeding at 6 months of age | 19.1% | 29.7% | 25.8% | 35.3% | 49 | 190 | 25.8% |

(1) Item appears also on other WCH Section Agreement Addenda. Data for 3A and 3B were not presented due to the limitation in recent trend datasets.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI >=30.0): Adapted from the National Academy of Sciences. *Weight Gain During Pregnancy: Reexamining the Guidelines*. Washington, DC : National Academy Press, 2009.

SFY 2014-2015 AGREEMENT ADDENDA DATA

WIC QUALITY ASSURANCE DELIVERABLES

Martin-Tyrrell-Washington District

| Process Outcome Objectives | 2004 to 2006 | 2007 to 2009 | 2010 to 2012 | % Change from 2004 to 2012 | 2012 Data | | |
|---|--------------------|--------------------|--------------------|----------------------------------|-----------|-------|---------|
| | | | | | Num | Den | Percent |
| 1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2) | 79.4% | 86.6% | 83.2% | 4.9% | 247 | 299 | 82.6% |
| 1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services. | 73.3% | 71.7% | 82.8% | 13.0% | 285 | 330 | 86.4% |
| 1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services. | 62.9% | 62.6% | 62.3% | -1.0% | 1,220 | 1,880 | 64.9% |
| 2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy. | 35.3% | 33.0% | 28.0% | -20.7% | 269 | 995 | 27.0% |
| 2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum. | 81.3% | 82.0% | 80.5% | -1.0% | 268 | 320 | 83.8% |
| 3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1) | | | | | | | |
| 3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1) | | | | | | | |
| 4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were: | | | | | | | |
| a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1) | 66.5% | 64.6% | 63.2% | -4.9% | 587 | 1,031 | 56.9% |
| b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile) | 13.2% | 12.9% | 13.4% | 1.4% | 155 | 1,031 | 15.0% |
| c. Obese (BMI at or greater than the 95th percentile). | 12.5% | 15.4% | 16.4% | 31.5% | 211 | 1,031 | 20.5% |
| d. Underweight (BMI less than the 5th percentile). | 7.8% | 7.1% | 6.9% | -10.8% | 78 | 1,031 | 7.6% |
| 4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3) | | | | | | | |
| a. Recommended prenatal weight gain | 29.3% | 30.9% | 26.4% | -9.7% | 61 | 252 | 24.2% |
| b. Excessive prenatal weight gain | 43.8% | 43.5% | 44.0% | 0.5% | 113 | 252 | 44.8% |
| c. Inadequate prenatal weight gain | 27.0% | 25.7% | 29.6% | 9.7% | 78 | 252 | 31.0% |
| 5A. Percentage of women who participated in WIC who initiated breastfeeding. | 40.3% | 36.6% | 23.0% | -43.0% | 90 | 388 | 23.2% |
| 5B. Percentage of infants who participated in WIC who were: | | | | | | | |
| a. Breastfeeding at 6 weeks of age (1) | 25.5% | 21.6% | 11.2% | -56.3% | 32 | 351 | 9.1% |
| b. Breastfeeding at 6 months of age | 12.0% | 9.5% | 4.3% | -64.4% | 17 | 351 | 4.8% |

(1) Item appears also on other WCH Section Agreement Addenda. Data for 3A and 3B were not presented due to the limitation in recent trend datasets.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI >=30.0): Adapted from the National Academy of Sciences. *Weight Gain During Pregnancy: Reexamining the Guidelines*. Washington, DC : National Academy Press, 2009.

SFY 2014-2015 AGREEMENT ADDENDA DATA

WIC QUALITY ASSURANCE DELIVERABLES

Mecklenburg County

| Process Outcome Objectives | 2004 to 2006 | 2007 to 2009 | 2010 to 2012 | % Change from 2004 to 2012 | 2012 Data | | |
|---|--------------------|--------------------|--------------------|----------------------------------|-----------|--------|---------|
| | | | | | Num | Den | Percent |
| 1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2) | 57.2% | 64.0% | 64.7% | 13.2% | 3,563 | 5,309 | 67.1% |
| 1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services. | 58.7% | 63.1% | 65.1% | 10.8% | 5,078 | 7,724 | 65.7% |
| 1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services. | 42.5% | 45.5% | 47.8% | 12.6% | 19,694 | 40,124 | 49.1% |
| 2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy. | 14.7% | 17.0% | 22.8% | 55.1% | 1,534 | 7,617 | 20.1% |
| 2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum. | 50.5% | 43.6% | 53.3% | 5.6% | 3,237 | 4,972 | 65.1% |
| 3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1) | | | | | | | |
| 3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1) | | | | | | | |
| 4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were: | | | | | | | |
| a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1) | 62.7% | 62.8% | 61.4% | -2.1% | 4,484 | 7,162 | 62.6% |
| b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile) | 17.5% | 17.4% | 17.2% | -1.6% | 1,151 | 7,162 | 16.1% |
| c. Obese (BMI at or greater than the 95th percentile). | 16.3% | 16.8% | 17.7% | 8.7% | 1,198 | 7,162 | 16.7% |
| d. Underweight (BMI less than the 5th percentile). | 3.5% | 3.0% | 3.7% | 5.4% | 329 | 7,162 | 4.6% |
| 4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3) | | | | | | | |
| a. Recommended prenatal weight gain | 31.9% | 33.7% | 29.0% | -9.2% | 948 | 3,653 | 26.0% |
| b. Excessive prenatal weight gain | 49.0% | 47.7% | 47.0% | -4.1% | 1,813 | 3,653 | 49.6% |
| c. Inadequate prenatal weight gain | 19.0% | 18.6% | 24.0% | 26.1% | 892 | 3,653 | 24.4% |
| 5A. Percentage of women who participated in WIC who initiated breastfeeding. | 50.1% | 56.1% | 54.3% | 8.3% | 3,482 | 6,146 | 56.7% |
| 5B. Percentage of infants who participated in WIC who were: | | | | | | | |
| a. Breastfeeding at 6 weeks of age (1) | 29.0% | 34.0% | 32.9% | 13.3% | 2,380 | 7,235 | 32.9% |
| b. Breastfeeding at 6 months of age | 15.0% | 17.6% | 18.3% | 21.7% | 1,296 | 7,235 | 17.9% |

(1) Item appears also on other WCH Section Agreement Addenda. Data for 3A and 3B were not presented due to the limitation in recent trend datasets.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI >=30.0): Adapted from the National Academy of Sciences. Weight Gain During Pregnancy: Reexamining the Guidelines. Washington, DC : National Academy Press, 2009.

SFY 2014-2015 AGREEMENT ADDENDA DATA

WIC QUALITY ASSURANCE DELIVERABLES

Montgomery County

| Process Outcome Objectives | 2004 to 2006 | 2007 to 2009 | 2010 to 2012 | % Change from 2004 to 2012 | 2012 Data | | |
|---|--------------------|--------------------|--------------------|----------------------------------|-----------|-------|---------|
| | | | | | Num | Den | Percent |
| 1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2) | 70.0% | 73.7% | 77.1% | 10.2% | 162 | 199 | 81.4% |
| 1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services. | 75.3% | 80.1% | 79.9% | 6.2% | 215 | 274 | 78.5% |
| 1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services. | 57.8% | 66.2% | 66.1% | 14.3% | 930 | 1,374 | 67.7% |
| 2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy. | 37.8% | 38.5% | 33.8% | -10.5% | 237 | 771 | 30.7% |
| 2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum. | 80.1% | 81.0% | 82.3% | 2.7% | 244 | 298 | 81.9% |
| 3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1) | | | | | | | |
| 3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1) | | | | | | | |
| 4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were: | | | | | | | |
| a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1) | 60.4% | 62.8% | 63.5% | 5.2% | 352 | 546 | 64.5% |
| b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile) | 17.7% | 16.7% | 16.9% | -4.9% | 84 | 546 | 15.4% |
| c. Obese (BMI at or greater than the 95th percentile). | 18.6% | 18.7% | 15.6% | -16.0% | 84 | 546 | 15.4% |
| d. Underweight (BMI less than the 5th percentile). | 3.2% | 1.8% | 4.0% | 22.8% | 26 | 546 | 4.8% |
| 4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3) | | | | | | | |
| a. Recommended prenatal weight gain | 31.4% | 36.1% | 32.9% | 4.8% | 53 | 195 | 27.2% |
| b. Excessive prenatal weight gain | 46.7% | 43.1% | 42.9% | -8.2% | 98 | 195 | 50.3% |
| c. Inadequate prenatal weight gain | 21.9% | 20.8% | 24.2% | 10.5% | 44 | 195 | 22.6% |
| 5A. Percentage of women who participated in WIC who initiated breastfeeding. | 56.9% | 56.8% | 61.0% | 7.1% | 206 | 342 | 60.2% |
| 5B. Percentage of infants who participated in WIC who were: | | | | | | | |
| a. Breastfeeding at 6 weeks of age (1) | 34.4% | 34.5% | 32.6% | -5.3% | 116 | 313 | 37.1% |
| b. Breastfeeding at 6 months of age | 16.0% | 17.2% | 15.4% | -3.5% | 51 | 313 | 16.3% |

(1) Item appears also on other WCH Section Agreement Addenda. Data for 3A and 3B were not presented due to the limitation in recent trend datasets.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI < 18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI ≥ 30.0): Adapted from the National Academy of Sciences. *Weight Gain During Pregnancy: Reexamining the Guidelines*. Washington, DC : National Academy Press, 2009.

SFY 2014-2015 AGREEMENT ADDENDA DATA

WIC QUALITY ASSURANCE DELIVERABLES

Moore County

| Process Outcome Objectives | 2004 to 2006 | 2007 to 2009 | 2010 to 2012 | % Change from 2004 to 2012 | 2012 Data | | |
|---|--------------------|--------------------|--------------------|----------------------------------|-----------|-------|---------|
| | | | | | Num | Den | Percent |
| 1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2) | 67.0% | 50.1% | 66.8% | -0.2% | 332 | 430 | 77.2% |
| 1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services. | 79.8% | 80.6% | 79.1% | -0.9% | 411 | 524 | 78.4% |
| 1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services. | 62.6% | 61.0% | 61.1% | -2.5% | 1,758 | 2,795 | 62.9% |
| 2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy. | 30.4% | 28.7% | 29.3% | -3.7% | 353 | 1,160 | 30.4% |
| 2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum. | 75.3% | 68.9% | 82.4% | 9.5% | 414 | 511 | 81.0% |
| 3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1) | | | | | | | |
| 3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1) | | | | | | | |
| 4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were: | | | | | | | |
| a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1) | 65.9% | 65.4% | 64.3% | -2.4% | 455 | 699 | 65.1% |
| b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile) | 16.9% | 15.4% | 16.6% | -1.4% | 112 | 699 | 16.0% |
| c. Obese (BMI at or greater than the 95th percentile). | 13.8% | 16.0% | 15.3% | 10.9% | 103 | 699 | 14.7% |
| d. Underweight (BMI less than the 5th percentile). | 3.5% | 3.2% | 3.7% | 8.4% | 29 | 699 | 4.1% |
| 4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3) | | | | | | | |
| a. Recommended prenatal weight gain | 28.6% | 28.4% | 27.3% | -4.5% | 117 | 389 | 30.1% |
| b. Excessive prenatal weight gain | 51.4% | 53.9% | 47.6% | -7.3% | 176 | 389 | 45.2% |
| c. Inadequate prenatal weight gain | 20.0% | 17.7% | 25.1% | 25.3% | 96 | 389 | 24.7% |
| 5A. Percentage of women who participated in WIC who initiated breastfeeding. | 44.8% | 57.5% | 57.5% | 28.4% | 393 | 620 | 63.4% |
| 5B. Percentage of infants who participated in WIC who were: | | | | | | | |
| a. Breastfeeding at 6 weeks of age (1) | 26.0% | 26.0% | 29.6% | 14.0% | 208 | 589 | 35.3% |
| b. Breastfeeding at 6 months of age | 12.1% | 11.8% | 16.5% | 36.6% | 132 | 589 | 22.4% |

(1) Item appears also on other WCH Section Agreement Addenda. Data for 3A and 3B were not presented due to the limitation in recent trend datasets.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI >=30.0): Adapted from the National Academy of Sciences. Weight Gain During Pregnancy: Reexamining the Guidelines. Washington, DC : National Academy Press, 2009.

SFY 2014-2015 AGREEMENT ADDENDA DATA

WIC QUALITY ASSURANCE DELIVERABLES

Nash County

| Process Outcome Objectives | 2004 to 2006 | 2007 to 2009 | 2010 to 2012 | % Change from 2004 to 2012 | 2012 Data | | |
|---|--------------------|--------------------|--------------------|----------------------------------|-----------|-------|---------|
| | | | | | Num | Den | Percent |
| 1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2) | 75.0% | 78.1% | 78.5% | 4.7% | 572 | 715 | 80.0% |
| 1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services. | 73.8% | 73.7% | 73.7% | -0.2% | 645 | 854 | 75.5% |
| 1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services. | 49.8% | 49.4% | 51.7% | 3.8% | 2,349 | 4,349 | 54.0% |
| 2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy. | 31.8% | 32.0% | 27.4% | -13.8% | 487 | 1,873 | 26.0% |
| 2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum. | 78.4% | 81.0% | 81.3% | 3.8% | 674 | 824 | 81.8% |
| 3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1) | | | | | | | |
| 3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1) | | | | | | | |
| 4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were: | | | | | | | |
| a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1) | 65.5% | 65.3% | 62.9% | -3.9% | 808 | 1,275 | 63.4% |
| b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile) | 13.7% | 15.6% | 15.0% | 9.5% | 182 | 1,275 | 14.3% |
| c. Obese (BMI at or greater than the 95th percentile). | 13.6% | 14.5% | 16.7% | 22.9% | 208 | 1,275 | 16.3% |
| d. Underweight (BMI less than the 5th percentile). | 7.3% | 4.6% | 5.4% | -25.4% | 77 | 1,275 | 6.0% |
| 4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3) | | | | | | | |
| a. Recommended prenatal weight gain | 30.3% | 31.4% | 28.8% | -4.9% | 170 | 619 | 27.5% |
| b. Excessive prenatal weight gain | 44.1% | 40.2% | 46.3% | 5.0% | 272 | 619 | 43.9% |
| c. Inadequate prenatal weight gain | 25.6% | 28.4% | 24.9% | -2.8% | 177 | 619 | 28.6% |
| 5A. Percentage of women who participated in WIC who initiated breastfeeding. | 47.0% | 42.4% | 47.6% | 1.2% | 467 | 992 | 47.1% |
| 5B. Percentage of infants who participated in WIC who were: | | | | | | | |
| a. Breastfeeding at 6 weeks of age (1) | 19.5% | 21.4% | 23.7% | 21.0% | 224 | 971 | 23.1% |
| b. Breastfeeding at 6 months of age | 8.9% | 10.0% | 12.4% | 39.8% | 113 | 971 | 11.6% |

(1) Item appears also on other WCH Section Agreement Addenda. Data for 3A and 3B were not presented due to the limitation in recent trend datasets.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI >=30.0): Adapted from the National Academy of Sciences. *Weight Gain During Pregnancy: Reexamining the Guidelines*. Washington, DC : National Academy Press, 2009.

SFY 2014-2015 AGREEMENT ADDENDA DATA

WIC QUALITY ASSURANCE DELIVERABLES

New Hanover County

| Process Outcome Objectives | 2004 to 2006 | 2007 to 2009 | 2010 to 2012 | % Change from 2004 to 2012 | 2012 Data | | |
|---|--------------------|--------------------|--------------------|----------------------------------|-----------|-------|---------|
| | | | | | Num | Den | Percent |
| 1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2) | 70.3% | 73.4% | 72.8% | 3.5% | 706 | 974 | 72.5% |
| 1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services. | 65.1% | 67.6% | 66.4% | 2.0% | 887 | 1,325 | 66.9% |
| 1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services. | 44.9% | 48.1% | 51.1% | 13.8% | 3,386 | 6,709 | 50.5% |
| 2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy. | 30.4% | 33.7% | 27.0% | -11.4% | 662 | 2,631 | 25.2% |
| 2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum. | 71.3% | 65.3% | 73.0% | 2.3% | 799 | 1,037 | 77.0% |
| 3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1) | | | | | | | |
| 3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1) | | | | | | | |
| 4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were: | | | | | | | |
| a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1) | 66.9% | 64.8% | 66.6% | -0.5% | 828 | 1,232 | 67.2% |
| b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile) | 15.2% | 17.3% | 15.0% | -1.4% | 183 | 1,232 | 14.9% |
| c. Obese (BMI at or greater than the 95th percentile). | 14.5% | 15.7% | 13.0% | -10.6% | 155 | 1,232 | 12.6% |
| d. Underweight (BMI less than the 5th percentile). | 3.4% | 2.3% | 5.4% | 61.8% | 66 | 1,232 | 5.4% |
| 4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3) | | | | | | | |
| a. Recommended prenatal weight gain | 30.1% | 30.9% | 28.5% | -5.1% | 229 | 888 | 25.8% |
| b. Excessive prenatal weight gain | 49.4% | 48.8% | 50.1% | 1.6% | 443 | 888 | 49.9% |
| c. Inadequate prenatal weight gain | 20.6% | 20.3% | 21.3% | 3.7% | 216 | 888 | 24.3% |
| 5A. Percentage of women who participated in WIC who initiated breastfeeding. | 44.7% | 53.8% | 61.4% | 37.4% | 878 | 1,340 | 65.5% |
| 5B. Percentage of infants who participated in WIC who were: | | | | | | | |
| a. Breastfeeding at 6 weeks of age (1) | 29.8% | 31.5% | 34.3% | 15.3% | 485 | 1,331 | 36.4% |
| b. Breastfeeding at 6 months of age | 14.8% | 16.9% | 18.7% | 26.5% | 260 | 1,331 | 19.5% |

(1) Item appears also on other WCH Section Agreement Addenda. Data for 3A and 3B were not presented due to the limitation in recent trend datasets.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI >=30.0): Adapted from the National Academy of Sciences. Weight Gain During Pregnancy: Reexamining the Guidelines. Washington, DC : National Academy Press, 2009.

SFY 2014-2015 AGREEMENT ADDENDA DATA

WIC QUALITY ASSURANCE DELIVERABLES

Northampton County

| Process Outcome Objectives | 2004 to 2006 | 2007 to 2009 | 2010 to 2012 | % Change from 2004 to 2012 | 2012 Data | | |
|---|--------------------|--------------------|--------------------|----------------------------------|-----------|-------|---------|
| | | | | | Num | Den | Percent |
| 1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2) | 81.6% | 77.1% | 83.0% | 1.7% | 139 | 169 | 82.2% |
| 1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services. | 75.5% | 70.9% | 69.8% | -7.5% | 135 | 203 | 66.5% |
| 1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services. | 61.9% | 63.2% | 60.6% | -2.1% | 657 | 1,053 | 62.4% |
| 2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy. | 33.6% | 34.2% | 34.1% | 1.7% | 134 | 368 | 36.4% |
| 2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum. | 73.5% | 73.6% | 76.9% | 4.6% | 122 | 160 | 76.3% |
| 3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1) | | | | | | | |
| 3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1) | | | | | | | |
| 4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were: | | | | | | | |
| a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1) | 60.0% | 62.3% | 61.6% | 2.6% | 296 | 486 | 60.9% |
| b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile) | 17.4% | 14.2% | 17.6% | 1.5% | 93 | 486 | 19.1% |
| c. Obese (BMI at or greater than the 95th percentile). | 17.8% | 18.2% | 16.2% | -9.2% | 72 | 486 | 14.8% |
| d. Underweight (BMI less than the 5th percentile). | 4.8% | 5.3% | 4.6% | -4.2% | 25 | 486 | 5.1% |
| 4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3) | | | | | | | |
| a. Recommended prenatal weight gain | 25.6% | 27.2% | 28.8% | 12.5% | 45 | 157 | 28.7% |
| b. Excessive prenatal weight gain | 51.2% | 49.5% | 43.7% | -14.8% | 64 | 157 | 40.8% |
| c. Inadequate prenatal weight gain | 23.1% | 23.2% | 27.5% | 19.0% | 48 | 157 | 30.6% |
| 5A. Percentage of women who participated in WIC who initiated breastfeeding. | 16.8% | 29.1% | 36.9% | 118.8% | 70 | 198 | 35.4% |
| 5B. Percentage of infants who participated in WIC who were: | | | | | | | |
| a. Breastfeeding at 6 weeks of age (1) | 11.4% | 16.7% | 20.7% | 82.2% | 50 | 195 | 25.6% |
| b. Breastfeeding at 6 months of age | 4.8% | 6.8% | 9.3% | 91.3% | 28 | 195 | 14.4% |

(1) Item appears also on other WCH Section Agreement Addenda. Data for 3A and 3B were not presented due to the limitation in recent trend datasets.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI >=30.0): Adapted from the National Academy of Sciences. Weight Gain During Pregnancy: Reexamining the Guidelines. Washington, DC : National Academy Press, 2009.

SFY 2014-2015 AGREEMENT ADDENDA DATA

WIC QUALITY ASSURANCE DELIVERABLES

Onslow County

| Process Outcome Objectives | 2004 to 2006 | 2007 to 2009 | 2010 to 2012 | % Change from 2004 to 2012 | 2012 Data | | |
|---|--------------------|--------------------|--------------------|----------------------------------|-----------|-------|---------|
| | | | | | Num | Den | Percent |
| 1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2) | 74.1% | 74.8% | 76.2% | 2.7% | 648 | 849 | 76.3% |
| 1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services. | 76.0% | 75.8% | 74.5% | -2.0% | 817 | 1,112 | 73.5% |
| 1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services. | 57.2% | 55.8% | 52.6% | -8.1% | 3,049 | 5,678 | 53.7% |
| 2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy. | 31.1% | 26.8% | 25.2% | -18.8% | 1,502 | 6,287 | 23.9% |
| 2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum. | 41.0% | 39.5% | 44.7% | 9.0% | 1,547 | 2,448 | 63.2% |
| 3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1) | | | | | | | |
| 3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1) | | | | | | | |
| 4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were: | | | | | | | |
| a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1) | 72.0% | 69.9% | 68.4% | -5.0% | 1,605 | 2,344 | 68.5% |
| b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile) | 14.7% | 15.9% | 15.8% | 8.1% | 358 | 2,344 | 15.3% |
| c. Obese (BMI at or greater than the 95th percentile). | 9.6% | 11.6% | 11.6% | 20.6% | 275 | 2,344 | 11.7% |
| d. Underweight (BMI less than the 5th percentile). | 3.7% | 2.6% | 4.1% | 12.1% | 106 | 2,344 | 4.5% |
| 4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3) | | | | | | | |
| a. Recommended prenatal weight gain | 33.4% | 33.2% | 28.8% | -13.8% | 581 | 2,121 | 27.4% |
| b. Excessive prenatal weight gain | 48.8% | 49.5% | 52.3% | 7.3% | 1,107 | 2,121 | 52.2% |
| c. Inadequate prenatal weight gain | 17.8% | 17.3% | 18.9% | 6.0% | 433 | 2,121 | 20.4% |
| 5A. Percentage of women who participated in WIC who initiated breastfeeding. | 52.0% | 53.1% | 50.3% | -3.2% | 1,812 | 3,305 | 54.8% |
| 5B. Percentage of infants who participated in WIC who were: | | | | | | | |
| a. Breastfeeding at 6 weeks of age (1) | 31.6% | 32.1% | 30.7% | -2.6% | 1,078 | 3,308 | 32.6% |
| b. Breastfeeding at 6 months of age | 13.0% | 12.7% | 12.5% | -3.2% | 452 | 3,308 | 13.7% |

(1) Item appears also on other WCH Section Agreement Addenda. Data for 3A and 3B were not presented due to the limitation in recent trend datasets.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI >=30.0): Adapted from the National Academy of Sciences. *Weight Gain During Pregnancy: Reexamining the Guidelines*. Washington, DC : National Academy Press, 2009.

SFY 2014-2015 AGREEMENT ADDENDA DATA

WIC QUALITY ASSURANCE DELIVERABLES

Orange County

| Process Outcome Objectives | 2004 to 2006 | 2007 to 2009 | 2010 to 2012 | % Change from 2004 to 2012 | 2012 Data | | |
|---|--------------------|--------------------|--------------------|----------------------------------|-----------|-------|---------|
| | | | | | Num | Den | Percent |
| 1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2) | 62.0% | 68.8% | 67.9% | 9.4% | 274 | 403 | 68.0% |
| 1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services. | 61.1% | 59.3% | 59.5% | -2.6% | 414 | 674 | 61.4% |
| 1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services. | 52.1% | 53.9% | 55.0% | 5.7% | 1,792 | 3,250 | 55.1% |
| 2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy. | 26.4% | 29.4% | 27.0% | 2.3% | 676 | 2,451 | 27.6% |
| 2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum. | 59.5% | 62.6% | 69.4% | 16.7% | 820 | 1,175 | 69.8% |
| 3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1) | | | | | | | |
| 3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1) | | | | | | | |
| 4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were: | | | | | | | |
| a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1) | 59.5% | 65.4% | 63.8% | 7.2% | 940 | 1,498 | 62.8% |
| b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile) | 19.0% | 15.3% | 15.4% | -19.2% | 232 | 1,498 | 15.5% |
| c. Obese (BMI at or greater than the 95th percentile). | 18.7% | 14.8% | 15.4% | -17.6% | 218 | 1,498 | 14.6% |
| d. Underweight (BMI less than the 5th percentile). | 2.8% | 4.5% | 5.4% | 95.6% | 108 | 1,498 | 7.2% |
| 4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3) | | | | | | | |
| a. Recommended prenatal weight gain | 28.7% | 30.3% | 32.9% | 14.7% | 115 | 376 | 30.6% |
| b. Excessive prenatal weight gain | 52.5% | 49.1% | 44.6% | -15.2% | 173 | 376 | 46.0% |
| c. Inadequate prenatal weight gain | 18.8% | 20.6% | 22.5% | 20.1% | 88 | 376 | 23.4% |
| 5A. Percentage of women who participated in WIC who initiated breastfeeding. | 63.5% | 67.3% | 70.6% | 11.2% | 1,021 | 1,441 | 70.9% |
| 5B. Percentage of infants who participated in WIC who were: | | | | | | | |
| a. Breastfeeding at 6 weeks of age (1) | 50.2% | 53.5% | 53.4% | 6.4% | 753 | 1,386 | 54.3% |
| b. Breastfeeding at 6 months of age | 29.7% | 33.6% | 33.4% | 12.6% | 473 | 1,386 | 34.1% |

(1) Item appears also on other WCH Section Agreement Addenda. Data for 3A and 3B were not presented due to the limitation in recent trend datasets.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI >=30.0): Adapted from the National Academy of Sciences. Weight Gain During Pregnancy: Reexamining the Guidelines. Washington, DC : National Academy Press, 2009.

SFY 2014-2015 AGREEMENT ADDENDA DATA

WIC QUALITY ASSURANCE DELIVERABLES

Pamlico County

| Process Outcome Objectives | 2004 to 2006 | 2007 to 2009 | 2010 to 2012 | % Change from 2004 to 2012 | 2012 Data | | |
|---|--------------------|--------------------|--------------------|----------------------------------|-----------|-----|---------|
| | | | | | Num | Den | Percent |
| 1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2) | 80.3% | 90.4% | 81.8% | 1.8% | 55 | 66 | 83.3% |
| 1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services. | 78.5% | 75.2% | 78.3% | -0.3% | 69 | 89 | 77.5% |
| 1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services. | 59.1% | 62.4% | 68.5% | 15.8% | 298 | 443 | 67.3% |
| 2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy. | 53.1% | 52.2% | 43.8% | -17.5% | 80 | 185 | 43.2% |
| 2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum. | 71.0% | 74.3% | 74.5% | 4.9% | 70 | 96 | 72.9% |
| 3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1) | | | | | | | |
| 3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1) | | | | | | | |
| 4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were: | | | | | | | |
| a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1) | 61.3% | 63.6% | 63.2% | 3.0% | 110 | 172 | 64.0% |
| b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile) | 19.6% | 18.1% | 17.5% | -11.1% | 27 | 172 | 15.7% |
| c. Obese (BMI at or greater than the 95th percentile). | 17.4% | 14.7% | 14.4% | -17.3% | 21 | 172 | 12.2% |
| d. Underweight (BMI less than the 5th percentile). | 1.6% | 3.7% | 5.0% | 207.9% | 14 | 172 | 8.1% |
| 4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3) | | | | | | | |
| a. Recommended prenatal weight gain | 28.7% | 30.1% | 30.3% | 5.7% | 22 | 76 | 28.9% |
| b. Excessive prenatal weight gain | 53.2% | 47.2% | 37.2% | -30.2% | 29 | 76 | 38.2% |
| c. Inadequate prenatal weight gain | 18.1% | 22.7% | 32.5% | 79.9% | 25 | 76 | 32.9% |
| 5A. Percentage of women who participated in WIC who initiated breastfeeding. | 53.7% | 50.1% | 49.4% | -8.0% | 56 | 119 | 47.1% |
| 5B. Percentage of infants who participated in WIC who were: | | | | | | | |
| a. Breastfeeding at 6 weeks of age (1) | 35.2% | 41.9% | 29.6% | -16.0% | 33 | 89 | 37.1% |
| b. Breastfeeding at 6 months of age | 21.6% | 25.5% | 16.6% | -23.1% | 21 | 89 | 23.6% |

(1) Item appears also on other WCH Section Agreement Addenda. Data for 3A and 3B were not presented due to the limitation in recent trend datasets.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI >=30.0): Adapted from the National Academy of Sciences. *Weight Gain During Pregnancy: Reexamining the Guidelines*. Washington, DC : National Academy Press, 2009.

SFY 2014-2015 AGREEMENT ADDENDA DATA

WIC QUALITY ASSURANCE DELIVERABLES

Pender County

| Process Outcome Objectives | 2004 to 2006 | 2007 to 2009 | 2010 to 2012 | % Change from 2004 to 2012 | 2012 Data | | |
|---|--------------------|--------------------|--------------------|----------------------------------|-----------|-------|---------|
| | | | | | Num | Den | Percent |
| 1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2) | 68.6% | 70.0% | 65.9% | -4.0% | 184 | 254 | 72.4% |
| 1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services. | 72.1% | 77.0% | 71.2% | -1.3% | 274 | 373 | 73.5% |
| 1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services. | 56.5% | 59.5% | 56.7% | 0.4% | 1,187 | 2,098 | 56.6% |
| 2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy. | 33.7% | 31.8% | 28.7% | -14.7% | 203 | 699 | 29.0% |
| 2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum. | 75.9% | 77.0% | 75.4% | -0.6% | 239 | 320 | 74.7% |
| 3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1) | | | | | | | |
| 3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1) | | | | | | | |
| 4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were: | | | | | | | |
| a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1) | 63.1% | 58.1% | 58.2% | -7.8% | 439 | 723 | 60.7% |
| b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile) | 14.3% | 18.5% | 18.5% | 29.7% | 124 | 723 | 17.2% |
| c. Obese (BMI at or greater than the 95th percentile). | 16.7% | 21.4% | 18.2% | 8.9% | 111 | 723 | 15.4% |
| d. Underweight (BMI less than the 5th percentile). | 5.9% | 2.0% | 5.1% | -14.1% | 49 | 723 | 6.8% |
| 4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3) | | | | | | | |
| a. Recommended prenatal weight gain | 31.5% | 30.2% | 28.0% | -11.0% | 59 | 250 | 23.6% |
| b. Excessive prenatal weight gain | 45.3% | 48.6% | 46.1% | 1.9% | 132 | 250 | 52.8% |
| c. Inadequate prenatal weight gain | 23.2% | 21.3% | 25.8% | 11.2% | 59 | 250 | 23.6% |
| 5A. Percentage of women who participated in WIC who initiated breastfeeding. | 41.6% | 44.5% | 56.0% | 34.8% | 238 | 416 | 57.2% |
| 5B. Percentage of infants who participated in WIC who were: | | | | | | | |
| a. Breastfeeding at 6 weeks of age (1) | 21.4% | 28.0% | 26.1% | 22.2% | 109 | 431 | 25.3% |
| b. Breastfeeding at 6 months of age | 7.3% | 12.2% | 14.2% | 93.4% | 64 | 431 | 14.8% |

(1) Item appears also on other WCH Section Agreement Addenda. Data for 3A and 3B were not presented due to the limitation in recent trend datasets.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI >=30.0): Adapted from the National Academy of Sciences. Weight Gain During Pregnancy: Reexamining the Guidelines. Washington, DC : National Academy Press, 2009.

SFY 2014-2015 AGREEMENT ADDENDA DATA

WIC QUALITY ASSURANCE DELIVERABLES

Person County

| Process Outcome Objectives | 2004 to 2006 | 2007 to 2009 | 2010 to 2012 | % Change from 2004 to 2012 | 2012 Data | | |
|---|--------------------|--------------------|--------------------|----------------------------------|-----------|-------|---------|
| | | | | | Num | Den | Percent |
| 1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2) | 83.1% | 83.6% | 83.6% | 0.6% | 195 | 234 | 83.3% |
| 1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services. | 76.6% | 80.7% | 83.1% | 8.5% | 234 | 279 | 83.9% |
| 1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services. | 62.6% | 65.7% | 68.5% | 9.4% | 1,087 | 1,560 | 69.7% |
| 2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy. | 37.9% | 37.2% | 33.1% | -12.7% | 185 | 574 | 32.2% |
| 2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum. | 74.5% | 71.8% | 81.3% | 9.2% | 218 | 269 | 81.0% |
| 3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1) | | | | | | | |
| 3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1) | | | | | | | |
| 4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were: | | | | | | | |
| a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1) | 67.2% | 62.4% | 64.6% | -3.9% | 325 | 502 | 64.7% |
| b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile) | 15.7% | 17.7% | 15.4% | -1.6% | 83 | 502 | 16.5% |
| c. Obese (BMI at or greater than the 95th percentile). | 14.3% | 16.1% | 16.5% | 15.4% | 74 | 502 | 14.7% |
| d. Underweight (BMI less than the 5th percentile). | 2.9% | 3.8% | 3.6% | 25.0% | 20 | 502 | 4.0% |
| 4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3) | | | | | | | |
| a. Recommended prenatal weight gain | 32.8% | 34.3% | 26.6% | -18.9% | 55 | 215 | 25.6% |
| b. Excessive prenatal weight gain | 44.2% | 48.1% | 50.6% | 14.5% | 108 | 215 | 50.2% |
| c. Inadequate prenatal weight gain | 23.1% | 17.6% | 22.9% | -0.9% | 52 | 215 | 24.2% |
| 5A. Percentage of women who participated in WIC who initiated breastfeeding. | 36.7% | 40.5% | 43.3% | 18.0% | 143 | 324 | 44.1% |
| 5B. Percentage of infants who participated in WIC who were: | | | | | | | |
| a. Breastfeeding at 6 weeks of age (1) | 21.8% | 21.8% | 24.4% | 12.0% | 59 | 299 | 19.7% |
| b. Breastfeeding at 6 months of age | 9.7% | 10.0% | 14.8% | 51.7% | 33 | 299 | 11.0% |

(1) Item appears also on other WCH Section Agreement Addenda. Data for 3A and 3B were not presented due to the limitation in recent trend datasets.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI >=30.0): Adapted from the National Academy of Sciences. Weight Gain During Pregnancy: Reexamining the Guidelines. Washington, DC : National Academy Press, 2009.

SFY 2014-2015 AGREEMENT ADDENDA DATA

WIC QUALITY ASSURANCE DELIVERABLES

Pitt County

| Process Outcome Objectives | 2004 to 2006 | 2007 to 2009 | 2010 to 2012 | % Change from 2004 to 2012 | 2012 Data | | |
|---|--------------------|--------------------|--------------------|----------------------------------|-----------|-------|---------|
| | | | | | Num | Den | Percent |
| 1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2) | 75.1% | 75.5% | 75.0% | -0.1% | 844 | 1,158 | 72.9% |
| 1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services. | 75.6% | 77.9% | 78.3% | 3.5% | 1,141 | 1,447 | 78.9% |
| 1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services. | 58.0% | 58.1% | 61.3% | 5.8% | 4,454 | 7,120 | 62.6% |
| 2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy. | 30.4% | 31.0% | 27.2% | -10.6% | 710 | 3,043 | 23.3% |
| 2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum. | 81.8% | 81.4% | 80.1% | -2.2% | 936 | 1,170 | 80.0% |
| 3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1) | | | | | | | |
| 3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1) | | | | | | | |
| 4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were: | | | | | | | |
| a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1) | 65.0% | 64.5% | 62.7% | -3.5% | 1,118 | 1,706 | 65.5% |
| b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile) | 16.7% | 16.7% | 16.6% | -0.4% | 237 | 1,706 | 13.9% |
| c. Obese (BMI at or greater than the 95th percentile). | 15.6% | 16.2% | 16.8% | 7.6% | 256 | 1,706 | 15.0% |
| d. Underweight (BMI less than the 5th percentile). | 2.7% | 2.7% | 3.9% | 42.7% | 95 | 1,706 | 5.6% |
| 4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3) | | | | | | | |
| a. Recommended prenatal weight gain | 31.9% | 33.0% | 31.0% | -3.0% | 313 | 1,013 | 30.9% |
| b. Excessive prenatal weight gain | 45.4% | 47.2% | 46.6% | 2.7% | 471 | 1,013 | 46.5% |
| c. Inadequate prenatal weight gain | 22.7% | 19.8% | 22.4% | -1.3% | 229 | 1,013 | 22.6% |
| 5A. Percentage of women who participated in WIC who initiated breastfeeding. | 46.2% | 48.9% | 46.8% | 1.1% | 691 | 1,473 | 46.9% |
| 5B. Percentage of infants who participated in WIC who were: | | | | | | | |
| a. Breastfeeding at 6 weeks of age (1) | 26.1% | 28.5% | 30.3% | 16.3% | 435 | 1,467 | 29.7% |
| b. Breastfeeding at 6 months of age | 11.5% | 14.2% | 15.9% | 38.0% | 235 | 1,467 | 16.0% |

(1) Item appears also on other WCH Section Agreement Addenda. Data for 3A and 3B were not presented due to the limitation in recent trend datasets.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI >=30.0): Adapted from the National Academy of Sciences. Weight Gain During Pregnancy: Reexamining the Guidelines. Washington, DC : National Academy Press, 2009.

SFY 2014-2015 AGREEMENT ADDENDA DATA

WIC QUALITY ASSURANCE DELIVERABLES

Randolph County

| Process Outcome Objectives | 2004 to 2006 | 2007 to 2009 | 2010 to 2012 | % Change from 2004 to 2012 | 2012 Data | | |
|---|--------------------|--------------------|--------------------|----------------------------------|-----------|-------|---------|
| | | | | | Num | Den | Percent |
| 1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2) | 66.9% | 69.1% | 73.2% | 9.4% | 679 | 915 | 74.2% |
| 1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services. | 74.3% | 76.6% | 75.7% | 1.9% | 914 | 1,217 | 75.1% |
| 1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services. | 52.1% | 54.7% | 57.6% | 10.5% | 3,877 | 6,487 | 59.8% |
| 2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy. | 29.7% | 28.6% | 29.8% | 0.3% | 514 | 1,837 | 28.0% |
| 2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum. | 61.8% | 55.1% | 82.6% | 33.6% | 817 | 982 | 83.2% |
| 3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1) | | | | | | | |
| 3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1) | | | | | | | |
| 4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were: | | | | | | | |
| a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1) | 58.2% | 65.4% | 63.8% | 9.6% | 1,030 | 1,601 | 64.3% |
| b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile) | 18.2% | 16.0% | 16.0% | -12.0% | 243 | 1,601 | 15.2% |
| c. Obese (BMI at or greater than the 95th percentile). | 21.1% | 15.7% | 16.1% | -23.7% | 252 | 1,601 | 15.7% |
| d. Underweight (BMI less than the 5th percentile). | 2.5% | 2.9% | 4.1% | 61.8% | 76 | 1,601 | 4.7% |
| 4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3) | | | | | | | |
| a. Recommended prenatal weight gain | 30.7% | 32.9% | 29.5% | -3.6% | 200 | 819 | 24.4% |
| b. Excessive prenatal weight gain | 52.4% | 48.8% | 50.1% | -4.4% | 426 | 819 | 52.0% |
| c. Inadequate prenatal weight gain | 16.9% | 18.2% | 20.3% | 20.2% | 193 | 819 | 23.6% |
| 5A. Percentage of women who participated in WIC who initiated breastfeeding. | 61.9% | 57.8% | 62.3% | 0.7% | 796 | 1,220 | 65.2% |
| 5B. Percentage of infants who participated in WIC who were: | | | | | | | |
| a. Breastfeeding at 6 weeks of age (1) | 34.7% | 36.3% | 37.0% | 6.6% | 477 | 1,150 | 41.5% |
| b. Breastfeeding at 6 months of age | 17.5% | 18.2% | 17.3% | -1.0% | 219 | 1,150 | 19.0% |

(1) Item appears also on other WCH Section Agreement Addenda. Data for 3A and 3B were not presented due to the limitation in recent trend datasets.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI >=30.0): Adapted from the National Academy of Sciences. Weight Gain During Pregnancy: Reexamining the Guidelines. Washington, DC : National Academy Press, 2009.

SFY 2014-2015 AGREEMENT ADDENDA DATA

WIC QUALITY ASSURANCE DELIVERABLES

Richmond County

| Process Outcome Objectives | 2004 to 2006 | 2007 to 2009 | 2010 to 2012 | % Change from 2004 to 2012 | 2012 Data | | |
|---|--------------------|--------------------|--------------------|----------------------------------|-----------|-------|---------|
| | | | | | Num | Den | Percent |
| 1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2) | 80.7% | 77.7% | 79.3% | -1.8% | 340 | 403 | 84.4% |
| 1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services. | 83.2% | 81.6% | 80.8% | -2.9% | 386 | 483 | 79.9% |
| 1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services. | 69.6% | 67.8% | 60.5% | -13.1% | 1,719 | 2,657 | 64.7% |
| 2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy. | 31.5% | 33.7% | 31.3% | -0.6% | 358 | 1,146 | 31.2% |
| 2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum. | 72.4% | 69.1% | 68.7% | -5.1% | 297 | 417 | 71.2% |
| 3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1) | | | | | | | |
| 3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1) | | | | | | | |
| 4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were: | | | | | | | |
| a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1) | 68.0% | 68.9% | 68.1% | 0.2% | 627 | 899 | 69.7% |
| b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile) | 8.2% | 11.1% | 12.2% | 49.4% | 114 | 899 | 12.7% |
| c. Obese (BMI at or greater than the 95th percentile). | 8.5% | 11.5% | 12.2% | 43.7% | 118 | 899 | 13.1% |
| d. Underweight (BMI less than the 5th percentile). | 15.4% | 8.5% | 7.6% | -50.8% | 40 | 899 | 4.4% |
| 4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3) | | | | | | | |
| a. Recommended prenatal weight gain | 30.2% | 28.5% | 25.9% | -14.3% | 92 | 359 | 25.6% |
| b. Excessive prenatal weight gain | 49.5% | 48.1% | 49.7% | 0.3% | 177 | 359 | 49.3% |
| c. Inadequate prenatal weight gain | 20.2% | 23.5% | 24.4% | 20.6% | 90 | 359 | 25.1% |
| 5A. Percentage of women who participated in WIC who initiated breastfeeding. | 38.8% | 28.1% | 31.3% | -19.3% | 184 | 506 | 36.4% |
| 5B. Percentage of infants who participated in WIC who were: | | | | | | | |
| a. Breastfeeding at 6 weeks of age (1) | 19.9% | 17.4% | 17.0% | -14.6% | 107 | 518 | 20.7% |
| b. Breastfeeding at 6 months of age | 9.3% | 7.5% | 8.6% | -8.4% | 58 | 518 | 11.2% |

(1) Item appears also on other WCH Section Agreement Addenda. Data for 3A and 3B were not presented due to the limitation in recent trend datasets.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI >=30.0): Adapted from the National Academy of Sciences. Weight Gain During Pregnancy: Reexamining the Guidelines. Washington, DC : National Academy Press, 2009.

SFY 2014-2015 AGREEMENT ADDENDA DATA

WIC QUALITY ASSURANCE DELIVERABLES

Robeson County

| Process Outcome Objectives | 2004 to 2006 | 2007 to 2009 | 2010 to 2012 | % Change from 2004 to 2012 | 2012 Data | | |
|---|--------------------|--------------------|--------------------|----------------------------------|-----------|-------|---------|
| | | | | | Num | Den | Percent |
| 1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2) | 72.4% | 75.1% | 72.2% | -0.3% | 1,006 | 1,368 | 73.5% |
| 1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services. | 76.2% | 75.8% | 77.0% | 1.0% | 1,272 | 1,667 | 76.3% |
| 1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services. | 50.3% | 49.0% | 50.6% | 0.6% | 4,762 | 9,114 | 52.2% |
| 2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy. | 28.0% | 20.1% | 22.8% | -18.5% | 572 | 2,450 | 23.3% |
| 2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum. | 82.1% | 81.7% | 80.7% | -1.6% | 989 | 1,249 | 79.2% |
| 3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1) | | | | | | | |
| 3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1) | | | | | | | |
| 4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were: | | | | | | | |
| a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1) | 66.4% | 63.9% | 56.8% | -14.5% | 1,180 | 2,019 | 58.4% |
| b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile) | 13.8% | 16.4% | 19.1% | 38.3% | 370 | 2,019 | 18.3% |
| c. Obese (BMI at or greater than the 95th percentile). | 13.7% | 16.5% | 21.1% | 54.1% | 409 | 2,019 | 20.3% |
| d. Underweight (BMI less than the 5th percentile). | 6.1% | 3.2% | 3.0% | -50.4% | 60 | 2,019 | 3.0% |
| 4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3) | | | | | | | |
| a. Recommended prenatal weight gain | 31.0% | 30.8% | 27.3% | -12.1% | 293 | 1,113 | 26.3% |
| b. Excessive prenatal weight gain | 41.2% | 43.1% | 46.4% | 12.6% | 541 | 1,113 | 48.6% |
| c. Inadequate prenatal weight gain | 27.8% | 26.1% | 26.3% | -5.2% | 279 | 1,113 | 25.1% |
| 5A. Percentage of women who participated in WIC who initiated breastfeeding. | 34.7% | 35.1% | 38.3% | 10.3% | 679 | 1,664 | 40.8% |
| 5B. Percentage of infants who participated in WIC who were: | | | | | | | |
| a. Breastfeeding at 6 weeks of age (1) | 18.6% | 20.8% | 20.5% | 10.7% | 360 | 1,545 | 23.3% |
| b. Breastfeeding at 6 months of age | 9.3% | 10.9% | 9.5% | 2.5% | 160 | 1,545 | 10.4% |

(1) Item appears also on other WCH Section Agreement Addenda. Data for 3A and 3B were not presented due to the limitation in recent trend datasets.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI >=30.0): Adapted from the National Academy of Sciences. Weight Gain During Pregnancy: Reexamining the Guidelines. Washington, DC : National Academy Press, 2009.

SFY 2014-2015 AGREEMENT ADDENDA DATA

WIC QUALITY ASSURANCE DELIVERABLES

Rockingham County

| Process Outcome Objectives | 2004 to 2006 | 2007 to 2009 | 2010 to 2012 | % Change from 2004 to 2012 | 2012 Data | | |
|---|--------------------|--------------------|--------------------|----------------------------------|-----------|-------|---------|
| | | | | | Num | Den | Percent |
| 1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2) | 77.7% | 78.6% | 78.3% | 0.8% | 481 | 609 | 79.0% |
| 1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services. | 76.3% | 76.6% | 77.9% | 2.1% | 595 | 754 | 78.9% |
| 1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services. | 53.7% | 56.7% | 58.7% | 9.3% | 2,189 | 3,644 | 60.1% |
| 2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy. | 34.5% | 35.5% | 32.3% | -6.1% | 495 | 1,638 | 30.2% |
| 2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum. | 75.1% | 74.2% | 76.6% | 2.0% | 501 | 666 | 75.2% |
| 3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1) | | | | | | | |
| 3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1) | | | | | | | |
| 4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were: | | | | | | | |
| a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1) | 69.4% | 63.3% | 65.9% | -5.0% | 690 | 1,025 | 67.3% |
| b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile) | 12.4% | 16.9% | 14.9% | 19.8% | 144 | 1,025 | 14.0% |
| c. Obese (BMI at or greater than the 95th percentile). | 12.1% | 17.8% | 15.2% | 25.4% | 145 | 1,025 | 14.1% |
| d. Underweight (BMI less than the 5th percentile). | 6.1% | 2.0% | 4.0% | -33.5% | 46 | 1,025 | 4.5% |
| 4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3) | | | | | | | |
| a. Recommended prenatal weight gain | 30.4% | 31.1% | 27.8% | -8.4% | 123 | 518 | 23.7% |
| b. Excessive prenatal weight gain | 48.1% | 49.3% | 49.5% | 2.9% | 269 | 518 | 51.9% |
| c. Inadequate prenatal weight gain | 21.5% | 19.6% | 22.6% | 5.4% | 126 | 518 | 24.3% |
| 5A. Percentage of women who participated in WIC who initiated breastfeeding. | 41.4% | 42.8% | 50.4% | 21.7% | 429 | 790 | 54.3% |
| 5B. Percentage of infants who participated in WIC who were: | | | | | | | |
| a. Breastfeeding at 6 weeks of age (1) | 25.3% | 26.3% | 27.2% | 7.5% | 238 | 734 | 32.4% |
| b. Breastfeeding at 6 months of age | 12.9% | 12.3% | 13.1% | 0.9% | 119 | 734 | 16.2% |

(1) Item appears also on other WCH Section Agreement Addenda. Data for 3A and 3B were not presented due to the limitation in recent trend datasets.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI >=30.0): Adapted from the National Academy of Sciences. Weight Gain During Pregnancy: Reexamining the Guidelines. Washington, DC : National Academy Press, 2009.

SFY 2014-2015 AGREEMENT ADDENDA DATA

WIC QUALITY ASSURANCE DELIVERABLES

Rowan County

| Process Outcome Objectives | 2004 to 2006 | 2007 to 2009 | 2010 to 2012 | % Change from 2004 to 2012 | 2012 Data | | |
|---|--------------------|--------------------|--------------------|----------------------------------|-----------|-------|---------|
| | | | | | Num | Den | Percent |
| 1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2) | 73.9% | 69.2% | 68.9% | -6.7% | 662 | 956 | 69.2% |
| 1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services. | 71.6% | 73.3% | 73.0% | 1.9% | 849 | 1,193 | 71.2% |
| 1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services. | 55.1% | 56.0% | 54.0% | -2.0% | 3,434 | 6,344 | 54.1% |
| 2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy. | 29.4% | 20.5% | 29.8% | 1.3% | 433 | 1,407 | 30.8% |
| 2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum. | 77.6% | 79.8% | 76.1% | -2.0% | 584 | 814 | 71.7% |
| 3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1) | | | | | | | |
| 3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1) | | | | | | | |
| 4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were: | | | | | | | |
| a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1) | 62.7% | 68.7% | 66.5% | 6.0% | 906 | 1,357 | 66.8% |
| b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile) | 18.5% | 14.1% | 14.9% | -19.7% | 204 | 1,357 | 15.0% |
| c. Obese (BMI at or greater than the 95th percentile). | 16.7% | 12.7% | 13.8% | -17.8% | 178 | 1,357 | 13.1% |
| d. Underweight (BMI less than the 5th percentile). | 2.0% | 4.5% | 4.9% | 141.5% | 69 | 1,357 | 5.1% |
| 4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3) | | | | | | | |
| a. Recommended prenatal weight gain | 31.8% | 31.8% | 28.9% | -9.1% | 213 | 778 | 27.4% |
| b. Excessive prenatal weight gain | 45.5% | 45.6% | 48.6% | 6.8% | 371 | 778 | 47.7% |
| c. Inadequate prenatal weight gain | 22.6% | 22.6% | 22.5% | -0.8% | 194 | 778 | 24.9% |
| 5A. Percentage of women who participated in WIC who initiated breastfeeding. | 49.8% | 51.0% | 53.2% | 6.8% | 559 | 1,026 | 54.5% |
| 5B. Percentage of infants who participated in WIC who were: | | | | | | | |
| a. Breastfeeding at 6 weeks of age (1) | 29.7% | 28.7% | 30.1% | 1.3% | 365 | 1,088 | 33.5% |
| b. Breastfeeding at 6 months of age | 12.2% | 12.0% | 12.5% | 2.7% | 162 | 1,088 | 14.9% |

(1) Item appears also on other WCH Section Agreement Addenda. Data for 3A and 3B were not presented due to the limitation in recent trend datasets.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI >=30.0): Adapted from the National Academy of Sciences. Weight Gain During Pregnancy: Reexamining the Guidelines. Washington, DC : National Academy Press, 2009.

SFY 2014-2015 AGREEMENT ADDENDA DATA

WIC QUALITY ASSURANCE DELIVERABLES

Rutherford-Polk-McDowell District

| Process Outcome Objectives | 2004 to 2006 | 2007 to 2009 | 2010 to 2012 | % Change from 2004 to 2012 | 2012 Data | | |
|---|--------------------|--------------------|--------------------|----------------------------------|-----------|-------|---------|
| | | | | | Num | Den | Percent |
| 1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2) | 64.4% | 71.7% | 78.9% | 22.5% | 719 | 847 | 84.9% |
| 1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services. | 74.7% | 76.5% | 75.2% | 0.6% | 771 | 1,026 | 75.1% |
| 1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services. | 60.7% | 61.5% | 60.4% | -0.5% | 3,321 | 5,436 | 61.1% |
| 2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy. | 32.9% | 35.4% | 33.2% | 1.0% | 847 | 2,654 | 31.9% |
| 2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum. | 71.0% | 73.2% | 78.0% | 9.8% | 785 | 990 | 79.3% |
| 3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1) | | | | | | | |
| 3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1) | | | | | | | |
| 4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were: | | | | | | | |
| a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1) | 67.6% | 67.2% | 65.7% | -2.9% | 1,056 | 1,630 | 64.8% |
| b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile) | 14.9% | 16.0% | 14.8% | -0.6% | 236 | 1,630 | 14.5% |
| c. Obese (BMI at or greater than the 95th percentile). | 13.1% | 13.1% | 14.2% | 8.1% | 261 | 1,630 | 16.0% |
| d. Underweight (BMI less than the 5th percentile). | 4.4% | 3.8% | 5.3% | 22.4% | 77 | 1,630 | 4.7% |
| 4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3) | | | | | | | |
| a. Recommended prenatal weight gain | 32.1% | 30.6% | 27.7% | -13.8% | 218 | 792 | 27.5% |
| b. Excessive prenatal weight gain | 37.0% | 41.9% | 50.2% | 35.6% | 396 | 792 | 50.0% |
| c. Inadequate prenatal weight gain | 30.9% | 27.5% | 22.2% | -28.3% | 178 | 792 | 22.5% |
| 5A. Percentage of women who participated in WIC who initiated breastfeeding. | 49.2% | 56.9% | 59.1% | 20.2% | 670 | 1,190 | 56.3% |
| 5B. Percentage of infants who participated in WIC who were: | | | | | | | |
| a. Breastfeeding at 6 weeks of age (1) | 30.0% | 29.7% | 32.2% | 7.2% | 374 | 1,081 | 34.6% |
| b. Breastfeeding at 6 months of age | 15.2% | 14.0% | 17.3% | 13.9% | 185 | 1,081 | 17.1% |

(1) Item appears also on other WCH Section Agreement Addenda. Data for 3A and 3B were not presented due to the limitation in recent trend datasets.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI >=30.0): Adapted from the National Academy of Sciences. Weight Gain During Pregnancy: Reexamining the Guidelines. Washington, DC : National Academy Press, 2009.

SFY 2014-2015 AGREEMENT ADDENDA DATA

WIC QUALITY ASSURANCE DELIVERABLES

Sampson County

| Process Outcome Objectives | 2004 to 2006 | 2007 to 2009 | 2010 to 2012 | % Change from 2004 to 2012 | 2012 Data | | |
|---|--------------------|--------------------|--------------------|----------------------------------|-----------|-------|---------|
| | | | | | Num | Den | Percent |
| 1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2) | 75.9% | 78.7% | 71.9% | -5.3% | 356 | 505 | 70.5% |
| 1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services. | 76.5% | 78.2% | 79.2% | 3.5% | 598 | 772 | 77.5% |
| 1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services. | 54.8% | 57.6% | 60.2% | 9.9% | 2,471 | 4,011 | 61.6% |
| 2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy. | 39.7% | 30.7% | 26.7% | -32.7% | 336 | 1,298 | 25.9% |
| 2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum. | 72.6% | 79.3% | 80.0% | 10.2% | 408 | 520 | 78.5% |
| 3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1) | | | | | | | |
| 3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1) | | | | | | | |
| 4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were: | | | | | | | |
| a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1) | 59.5% | 54.1% | 57.1% | -4.1% | 729 | 1,216 | 60.0% |
| b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile) | 16.5% | 19.8% | 18.5% | 12.0% | 194 | 1,216 | 16.0% |
| c. Obese (BMI at or greater than the 95th percentile). | 19.4% | 23.8% | 21.3% | 9.6% | 254 | 1,216 | 20.9% |
| d. Underweight (BMI less than the 5th percentile). | 4.6% | 2.3% | 3.2% | -30.2% | 39 | 1,216 | 3.2% |
| 4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3) | | | | | | | |
| a. Recommended prenatal weight gain | 30.9% | 31.6% | 28.2% | -8.5% | 112 | 434 | 25.8% |
| b. Excessive prenatal weight gain | 48.6% | 50.3% | 44.0% | -9.5% | 209 | 434 | 48.2% |
| c. Inadequate prenatal weight gain | 20.5% | 18.2% | 27.8% | 35.2% | 113 | 434 | 26.0% |
| 5A. Percentage of women who participated in WIC who initiated breastfeeding. | 26.4% | 27.2% | 31.9% | 21.1% | 273 | 664 | 41.1% |
| 5B. Percentage of infants who participated in WIC who were: | | | | | | | |
| a. Breastfeeding at 6 weeks of age (1) | 15.6% | 20.7% | 18.6% | 19.0% | 173 | 668 | 25.9% |
| b. Breastfeeding at 6 months of age | 7.4% | 13.0% | 12.1% | 62.4% | 111 | 668 | 16.6% |

(1) Item appears also on other WCH Section Agreement Addenda. Data for 3A and 3B were not presented due to the limitation in recent trend datasets.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI >=30.0): Adapted from the National Academy of Sciences. *Weight Gain During Pregnancy: Reexamining the Guidelines*. Washington, DC : National Academy Press, 2009.

SFY 2014-2015 AGREEMENT ADDENDA DATA

WIC QUALITY ASSURANCE DELIVERABLES

Scotland County

| Process Outcome Objectives | 2004 to 2006 | 2007 to 2009 | 2010 to 2012 | % Change from 2004 to 2012 | 2012 Data | | |
|---|--------------------|--------------------|--------------------|----------------------------------|-----------|-------|---------|
| | | | | | Num | Den | Percent |
| 1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2) | 85.2% | 84.3% | 84.5% | -0.8% | 305 | 348 | 87.6% |
| 1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services. | 80.0% | 82.1% | 80.4% | 0.4% | 328 | 407 | 80.6% |
| 1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services. | 63.8% | 62.3% | 63.3% | -0.7% | 1,462 | 2,257 | 64.8% |
| 2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy. | 55.4% | 48.9% | 40.5% | -26.8% | 373 | 978 | 38.1% |
| 2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum. | 79.3% | 80.1% | 81.0% | 2.2% | 323 | 393 | 82.2% |
| 3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1) | | | | | | | |
| 3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1) | | | | | | | |
| 4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were: | | | | | | | |
| a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1) | 68.8% | 67.1% | 67.7% | -1.5% | 531 | 749 | 70.9% |
| b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile) | 12.9% | 16.1% | 12.9% | 0.0% | 78 | 749 | 10.4% |
| c. Obese (BMI at or greater than the 95th percentile). | 13.1% | 12.4% | 13.6% | 3.8% | 96 | 749 | 12.8% |
| d. Underweight (BMI less than the 5th percentile). | 5.2% | 4.4% | 5.7% | 10.7% | 44 | 749 | 5.9% |
| 4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3) | | | | | | | |
| a. Recommended prenatal weight gain | 29.3% | 27.5% | 23.6% | -19.5% | 84 | 353 | 23.8% |
| b. Excessive prenatal weight gain | 47.6% | 46.5% | 43.7% | -8.1% | 147 | 353 | 41.6% |
| c. Inadequate prenatal weight gain | 23.1% | 26.0% | 32.7% | 41.5% | 122 | 353 | 34.6% |
| 5A. Percentage of women who participated in WIC who initiated breastfeeding. | 32.9% | 24.5% | 32.2% | -2.0% | 169 | 474 | 35.7% |
| 5B. Percentage of infants who participated in WIC who were: | | | | | | | |
| a. Breastfeeding at 6 weeks of age (1) | 11.1% | 8.3% | 15.3% | 38.3% | 97 | 495 | 19.6% |
| b. Breastfeeding at 6 months of age | 3.0% | 2.4% | 7.0% | 131.5% | 39 | 495 | 7.9% |

(1) Item appears also on other WCH Section Agreement Addenda. Data for 3A and 3B were not presented due to the limitation in recent trend datasets.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI >=30.0): Adapted from the National Academy of Sciences. Weight Gain During Pregnancy: Reexamining the Guidelines. Washington, DC : National Academy Press, 2009.

SFY 2014-2015 AGREEMENT ADDENDA DATA

WIC QUALITY ASSURANCE DELIVERABLES

Stanly County

| Process Outcome Objectives | 2004 to 2006 | 2007 to 2009 | 2010 to 2012 | % Change from 2004 to 2012 | 2012 Data | | |
|---|--------------------|--------------------|--------------------|----------------------------------|-----------|-------|---------|
| | | | | | Num | Den | Percent |
| 1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2) | 76.3% | 81.7% | 76.7% | 0.6% | 300 | 379 | 79.2% |
| 1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services. | 76.5% | 75.8% | 76.5% | -0.1% | 334 | 432 | 77.3% |
| 1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services. | 57.0% | 55.3% | 55.5% | -2.6% | 1,354 | 2,367 | 57.2% |
| 2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy. | 37.1% | 38.3% | 35.0% | -5.7% | 303 | 856 | 35.4% |
| 2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum. | 61.2% | 39.0% | 78.0% | 27.4% | 299 | 362 | 82.6% |
| 3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1) | | | | | | | |
| 3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1) | | | | | | | |
| 4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were: | | | | | | | |
| a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1) | 70.8% | 67.8% | 64.2% | -9.4% | 389 | 583 | 66.7% |
| b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile) | 12.0% | 14.6% | 15.3% | 27.3% | 94 | 583 | 16.1% |
| c. Obese (BMI at or greater than the 95th percentile). | 11.1% | 13.1% | 15.8% | 41.9% | 77 | 583 | 13.2% |
| d. Underweight (BMI less than the 5th percentile). | 6.0% | 4.5% | 4.7% | -22.1% | 23 | 583 | 3.9% |
| 4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3) | | | | | | | |
| a. Recommended prenatal weight gain | 32.3% | 28.5% | 28.2% | -12.6% | 94 | 353 | 26.6% |
| b. Excessive prenatal weight gain | 41.2% | 44.4% | 47.9% | 16.3% | 165 | 353 | 46.7% |
| c. Inadequate prenatal weight gain | 26.5% | 27.1% | 23.9% | -10.0% | 94 | 353 | 26.6% |
| 5A. Percentage of women who participated in WIC who initiated breastfeeding. | 63.1% | 64.8% | 66.2% | 5.0% | 291 | 461 | 63.1% |
| 5B. Percentage of infants who participated in WIC who were: | | | | | | | |
| a. Breastfeeding at 6 weeks of age (1) | 15.6% | 20.6% | 26.7% | 71.4% | 150 | 490 | 30.6% |
| b. Breastfeeding at 6 months of age | 6.5% | 10.2% | 14.7% | 127.2% | 93 | 490 | 19.0% |

(1) Item appears also on other WCH Section Agreement Addenda. Data for 3A and 3B were not presented due to the limitation in recent trend datasets.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI >=30.0): Adapted from the National Academy of Sciences. Weight Gain During Pregnancy: Reexamining the Guidelines. Washington, DC : National Academy Press, 2009.

SFY 2014-2015 AGREEMENT ADDENDA DATA

WIC QUALITY ASSURANCE DELIVERABLES

Stokes County

| Process Outcome Objectives | 2004 to 2006 | 2007 to 2009 | 2010 to 2012 | % Change from 2004 to 2012 | 2012 Data | | |
|---|--------------------|--------------------|--------------------|----------------------------------|-----------|-------|---------|
| | | | | | Num | Den | Percent |
| 1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2) | 75.9% | 76.0% | 69.3% | -8.6% | 167 | 249 | 67.1% |
| 1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services. | 75.0% | 75.5% | 72.0% | -3.9% | 196 | 287 | 68.3% |
| 1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services. | 55.5% | 57.8% | 55.7% | 0.3% | 818 | 1,487 | 55.0% |
| 2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy. | 34.4% | 39.3% | 34.2% | -0.5% | 208 | 566 | 36.7% |
| 2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum. | 81.1% | 75.9% | 78.1% | -3.7% | 177 | 226 | 78.3% |
| 3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1) | | | | | | | |
| 3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1) | | | | | | | |
| 4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were: | | | | | | | |
| a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1) | 65.5% | 61.7% | 66.9% | 2.1% | 242 | 381 | 63.5% |
| b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile) | 16.8% | 18.4% | 15.8% | -5.8% | 64 | 381 | 16.8% |
| c. Obese (BMI at or greater than the 95th percentile). | 14.2% | 17.2% | 13.5% | -5.4% | 57 | 381 | 15.0% |
| d. Underweight (BMI less than the 5th percentile). | 3.5% | 2.7% | 3.8% | 10.5% | 18 | 381 | 4.7% |
| 4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3) | | | | | | | |
| a. Recommended prenatal weight gain | 32.7% | 28.5% | 25.7% | -21.4% | 51 | 219 | 23.3% |
| b. Excessive prenatal weight gain | 52.3% | 54.1% | 46.3% | -11.4% | 102 | 219 | 46.6% |
| c. Inadequate prenatal weight gain | 15.0% | 17.4% | 28.0% | 86.0% | 66 | 219 | 30.1% |
| 5A. Percentage of women who participated in WIC who initiated breastfeeding. | 49.1% | 49.0% | 53.1% | 8.0% | 180 | 313 | 57.5% |
| 5B. Percentage of infants who participated in WIC who were: | | | | | | | |
| a. Breastfeeding at 6 weeks of age (1) | 21.8% | 23.5% | 24.9% | 14.5% | 85 | 304 | 28.0% |
| b. Breastfeeding at 6 months of age | 9.1% | 11.3% | 10.3% | 13.2% | 35 | 304 | 11.5% |

(1) Item appears also on other WCH Section Agreement Addenda. Data for 3A and 3B were not presented due to the limitation in recent trend datasets.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI >=30.0): Adapted from the National Academy of Sciences. Weight Gain During Pregnancy: Reexamining the Guidelines. Washington, DC : National Academy Press, 2009.

SFY 2014-2015 AGREEMENT ADDENDA DATA

WIC QUALITY ASSURANCE DELIVERABLES

Surry County

| Process Outcome Objectives | 2004 to 2006 | 2007 to 2009 | 2010 to 2012 | % Change from 2004 to 2012 | 2012 Data | | |
|---|--------------------|--------------------|--------------------|----------------------------------|-----------|-------|---------|
| | | | | | Num | Den | Percent |
| 1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2) | 70.2% | 68.4% | 70.7% | 0.7% | 362 | 478 | 75.7% |
| 1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services. | 76.3% | 76.8% | 75.4% | -1.1% | 485 | 619 | 78.4% |
| 1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services. | 52.5% | 54.1% | 57.4% | 9.4% | 1,909 | 3,224 | 59.2% |
| 2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy. | 46.2% | 43.4% | 41.0% | -11.2% | 381 | 934 | 40.8% |
| 2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum. | 80.3% | 79.9% | 80.9% | 0.8% | 370 | 469 | 78.9% |
| 3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1) | | | | | | | |
| 3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1) | | | | | | | |
| 4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were: | | | | | | | |
| a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1) | 67.8% | 62.7% | 59.7% | -11.8% | 558 | 962 | 58.0% |
| b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile) | 14.1% | 17.4% | 14.4% | 1.9% | 125 | 962 | 13.0% |
| c. Obese (BMI at or greater than the 95th percentile). | 12.9% | 17.1% | 17.2% | 33.3% | 152 | 962 | 15.8% |
| d. Underweight (BMI less than the 5th percentile). | 5.2% | 2.8% | 8.6% | 66.2% | 127 | 962 | 13.2% |
| 4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3) | | | | | | | |
| a. Recommended prenatal weight gain | 29.8% | 29.3% | 27.9% | -6.3% | 101 | 392 | 25.8% |
| b. Excessive prenatal weight gain | 52.3% | 53.0% | 49.7% | -5.1% | 196 | 392 | 50.0% |
| c. Inadequate prenatal weight gain | 17.9% | 17.7% | 22.4% | 25.2% | 95 | 392 | 24.2% |
| 5A. Percentage of women who participated in WIC who initiated breastfeeding. | 50.8% | 44.8% | 48.3% | -4.8% | 242 | 593 | 40.8% |
| 5B. Percentage of infants who participated in WIC who were: | | | | | | | |
| a. Breastfeeding at 6 weeks of age (1) | 34.3% | 26.5% | 32.1% | -6.4% | 185 | 626 | 29.6% |
| b. Breastfeeding at 6 months of age | 17.5% | 11.3% | 18.5% | 5.3% | 116 | 626 | 18.5% |

(1) Item appears also on other WCH Section Agreement Addenda. Data for 3A and 3B were not presented due to the limitation in recent trend datasets.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI >=30.0): Adapted from the National Academy of Sciences. Weight Gain During Pregnancy: Reexamining the Guidelines. Washington, DC : National Academy Press, 2009.

SFY 2014-2015 AGREEMENT ADDENDA DATA

WIC QUALITY ASSURANCE DELIVERABLES

Swain County

| Process Outcome Objectives | 2004 to 2006 | 2007 to 2009 | 2010 to 2012 | % Change from 2004 to 2012 | 2012 Data | | |
|---|--------------------|--------------------|--------------------|----------------------------------|-----------|-----|---------|
| | | | | | Num | Den | Percent |
| 1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2) | 87.0% | 85.4% | 77.9% | -10.4% | 124 | 152 | 81.6% |
| 1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services. | 75.5% | 69.9% | 69.1% | -8.4% | 117 | 167 | 70.1% |
| 1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services. | 75.1% | 75.9% | 72.6% | -3.3% | 626 | 858 | 73.0% |
| 2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy. | 48.7% | 45.7% | 38.5% | -20.8% | 83 | 192 | 43.2% |
| 2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum. | 75.7% | 74.1% | 61.0% | -19.4% | 57 | 77 | 74.0% |
| 3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1) | | | | | | | |
| 3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1) | | | | | | | |
| 4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were: | | | | | | | |
| a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1) | 71.9% | 68.0% | 65.7% | -8.6% | 124 | 187 | 66.3% |
| b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile) | 14.3% | 17.8% | 15.8% | 10.2% | 28 | 187 | 15.0% |
| c. Obese (BMI at or greater than the 95th percentile). | 11.6% | 12.2% | 13.1% | 13.8% | 24 | 187 | 12.8% |
| d. Underweight (BMI less than the 5th percentile). | 2.2% | 2.0% | 5.3% | 143.2% | 11 | 187 | 5.9% |
| 4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3) | | | | | | | |
| a. Recommended prenatal weight gain | 31.2% | 27.5% | 25.9% | -16.8% | 28 | 120 | 23.3% |
| b. Excessive prenatal weight gain | 49.5% | 57.8% | 51.6% | 4.2% | 62 | 120 | 51.7% |
| c. Inadequate prenatal weight gain | 19.3% | 14.7% | 22.4% | 16.3% | 30 | 120 | 25.0% |
| 5A. Percentage of women who participated in WIC who initiated breastfeeding. | 76.1% | 74.5% | 57.7% | -24.2% | 27 | 90 | 30.0% |
| 5B. Percentage of infants who participated in WIC who were: | | | | | | | |
| a. Breastfeeding at 6 weeks of age (1) | 36.0% | 32.4% | 45.1% | 25.3% | 27 | 86 | 31.4% |
| b. Breastfeeding at 6 months of age | 19.9% | 18.2% | 25.3% | 26.7% | 17 | 86 | 19.8% |

(1) Item appears also on other WCH Section Agreement Addenda. Data for 3A and 3B were not presented due to the limitation in recent trend datasets.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI >=30.0): Adapted from the National Academy of Sciences. Weight Gain During Pregnancy: Reexamining the Guidelines. Washington, DC : National Academy Press, 2009.

SFY 2014-2015 AGREEMENT ADDENDA DATA

WIC QUALITY ASSURANCE DELIVERABLES

Toe River District

| Process Outcome Objectives | 2004 to 2006 | 2007 to 2009 | 2010 to 2012 | % Change from 2004 to 2012 | 2012 Data | | |
|---|--------------------|--------------------|--------------------|----------------------------------|-----------|-------|---------|
| | | | | | Num | Den | Percent |
| 1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2) | 73.8% | 72.0% | 73.5% | -0.3% | 231 | 297 | 77.8% |
| 1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services. | 63.3% | 64.6% | 69.4% | 9.6% | 253 | 349 | 72.5% |
| 1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services. | 62.9% | 61.5% | 65.5% | 4.2% | 1,177 | 1,797 | 65.5% |
| 2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy. | 40.9% | 43.9% | 37.9% | -7.2% | 303 | 845 | 35.9% |
| 2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum. | 68.0% | 68.8% | 69.4% | 2.1% | 232 | 316 | 73.4% |
| 3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1) | | | | | | | |
| 3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1) | | | | | | | |
| 4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were: | | | | | | | |
| a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1) | 67.0% | 66.9% | 65.0% | -2.9% | 534 | 818 | 65.3% |
| b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile) | 15.0% | 16.1% | 15.7% | 4.9% | 115 | 818 | 14.1% |
| c. Obese (BMI at or greater than the 95th percentile). | 15.1% | 13.3% | 13.4% | -11.4% | 94 | 818 | 11.5% |
| d. Underweight (BMI less than the 5th percentile). | 3.0% | 3.7% | 5.9% | 97.7% | 75 | 818 | 9.2% |
| 4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3) | | | | | | | |
| a. Recommended prenatal weight gain | 34.3% | 31.0% | 30.1% | -12.3% | 67 | 238 | 28.2% |
| b. Excessive prenatal weight gain | 42.7% | 46.3% | 45.8% | 7.3% | 119 | 238 | 50.0% |
| c. Inadequate prenatal weight gain | 23.0% | 22.7% | 24.1% | 4.8% | 52 | 238 | 21.8% |
| 5A. Percentage of women who participated in WIC who initiated breastfeeding. | 59.2% | 67.5% | 63.4% | 7.0% | 268 | 389 | 68.9% |
| 5B. Percentage of infants who participated in WIC who were: | | | | | | | |
| a. Breastfeeding at 6 weeks of age (1) | 36.6% | 39.8% | 42.0% | 14.8% | 180 | 411 | 43.8% |
| b. Breastfeeding at 6 months of age | 24.6% | 24.2% | 28.1% | 14.3% | 113 | 411 | 27.5% |

(1) Item appears also on other WCH Section Agreement Addenda. Data for 3A and 3B were not presented due to the limitation in recent trend datasets.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI >=30.0): Adapted from the National Academy of Sciences. Weight Gain During Pregnancy: Reexamining the Guidelines. Washington, DC : National Academy Press, 2009.

SFY 2014-2015 AGREEMENT ADDENDA DATA

WIC QUALITY ASSURANCE DELIVERABLES

Transylvania County

| Process Outcome Objectives | 2004 to 2006 | 2007 to 2009 | 2010 to 2012 | % Change from 2004 to 2012 | 2012 Data | | |
|---|--------------------|--------------------|--------------------|----------------------------------|-----------|-------|---------|
| | | | | | Num | Den | Percent |
| 1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2) | 73.5% | 71.4% | 68.1% | -7.3% | 108 | 159 | 67.9% |
| 1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services. | 69.2% | 65.2% | 65.0% | -6.0% | 122 | 188 | 64.9% |
| 1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services. | 55.5% | 55.3% | 56.7% | 2.2% | 610 | 1,078 | 56.6% |
| 2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy. | 50.4% | 50.8% | 45.9% | -8.9% | 170 | 393 | 43.3% |
| 2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum. | 76.9% | 76.1% | 72.4% | -5.9% | 121 | 160 | 75.6% |
| 3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1) | | | | | | | |
| 3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1) | | | | | | | |
| 4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were: | | | | | | | |
| a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1) | 66.6% | 67.8% | 64.5% | -3.1% | 175 | 273 | 64.1% |
| b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile) | 16.2% | 15.5% | 17.9% | 10.6% | 54 | 273 | 19.8% |
| c. Obese (BMI at or greater than the 95th percentile). | 14.3% | 13.7% | 14.5% | 1.2% | 39 | 273 | 14.3% |
| d. Underweight (BMI less than the 5th percentile). | 2.9% | 2.9% | 3.1% | 5.9% | 5 | 273 | 1.8% |
| 4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3) | | | | | | | |
| a. Recommended prenatal weight gain | 31.1% | 33.3% | 26.6% | -14.5% | 27 | 133 | 20.3% |
| b. Excessive prenatal weight gain | 51.0% | 47.8% | 50.3% | -1.3% | 73 | 133 | 54.9% |
| c. Inadequate prenatal weight gain | 17.9% | 19.0% | 23.1% | 28.9% | 33 | 133 | 24.8% |
| 5A. Percentage of women who participated in WIC who initiated breastfeeding. | 80.1% | 78.2% | 78.6% | -1.8% | 156 | 204 | 76.5% |
| 5B. Percentage of infants who participated in WIC who were: | | | | | | | |
| a. Breastfeeding at 6 weeks of age (1) | 51.1% | 54.4% | 47.4% | -7.3% | 105 | 220 | 47.7% |
| b. Breastfeeding at 6 months of age | 30.2% | 30.7% | 30.6% | 1.0% | 70 | 220 | 31.8% |

(1) Item appears also on other WCH Section Agreement Addenda. Data for 3A and 3B were not presented due to the limitation in recent trend datasets.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI >=30.0): Adapted from the National Academy of Sciences. Weight Gain During Pregnancy: Reexamining the Guidelines. Washington, DC : National Academy Press, 2009.

SFY 2014-2015 AGREEMENT ADDENDA DATA

WIC QUALITY ASSURANCE DELIVERABLES

Union County

| Process Outcome Objectives | 2004 to 2006 | 2007 to 2009 | 2010 to 2012 | % Change from 2004 to 2012 | 2012 Data | | |
|---|--------------------|--------------------|--------------------|----------------------------------|-----------|-------|---------|
| | | | | | Num | Den | Percent |
| 1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2) | 62.0% | 54.8% | 59.5% | -4.2% | 533 | 865 | 61.6% |
| 1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services. | 61.9% | 65.2% | 68.0% | 9.8% | 837 | 1,276 | 65.6% |
| 1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services. | 50.2% | 51.6% | 55.4% | 10.2% | 3,829 | 6,813 | 56.2% |
| 2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy. | 26.5% | 12.6% | 21.6% | -18.6% | 353 | 1,279 | 27.6% |
| 2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum. | 78.8% | 76.3% | 83.8% | 6.4% | 699 | 830 | 84.2% |
| 3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1) | | | | | | | |
| 3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1) | | | | | | | |
| 4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were: | | | | | | | |
| a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1) | 66.3% | 65.7% | 56.5% | -14.9% | 1,201 | 2,194 | 54.7% |
| b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile) | 16.6% | 16.0% | 16.9% | 2.1% | 282 | 2,194 | 12.9% |
| c. Obese (BMI at or greater than the 95th percentile). | 14.2% | 15.1% | 17.6% | 24.3% | 262 | 2,194 | 11.9% |
| d. Underweight (BMI less than the 5th percentile). | 2.9% | 3.2% | 9.0% | 206.3% | 449 | 2,194 | 20.5% |
| 4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3) | | | | | | | |
| a. Recommended prenatal weight gain | 35.1% | 38.0% | 30.2% | -14.0% | 200 | 666 | 30.0% |
| b. Excessive prenatal weight gain | 43.0% | 40.0% | 45.1% | 4.8% | 309 | 666 | 46.4% |
| c. Inadequate prenatal weight gain | 21.9% | 22.1% | 24.8% | 13.0% | 157 | 666 | 23.6% |
| 5A. Percentage of women who participated in WIC who initiated breastfeeding. | 66.0% | 70.5% | 72.4% | 9.6% | 861 | 1,153 | 74.7% |
| 5B. Percentage of infants who participated in WIC who were: | | | | | | | |
| a. Breastfeeding at 6 weeks of age (1) | 36.9% | 35.7% | 40.0% | 8.5% | 534 | 1,230 | 43.4% |
| b. Breastfeeding at 6 months of age | 19.2% | 19.2% | 24.1% | 26.1% | 304 | 1,230 | 24.7% |

(1) Item appears also on other WCH Section Agreement Addenda. Data for 3A and 3B were not presented due to the limitation in recent trend datasets.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI >=30.0): Adapted from the National Academy of Sciences. Weight Gain During Pregnancy: Reexamining the Guidelines. Washington, DC : National Academy Press, 2009.

SFY 2014-2015 AGREEMENT ADDENDA DATA

WIC QUALITY ASSURANCE DELIVERABLES

Wake County

| Process Outcome Objectives | 2004 to 2006 | 2007 to 2009 | 2010 to 2012 | % Change from 2004 to 2012 | 2012 Data | | |
|---|--------------------|--------------------|--------------------|----------------------------------|-----------|--------|---------|
| | | | | | Num | Den | Percent |
| 1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2) | 68.2% | 68.7% | 69.0% | 1.1% | 2,572 | 3,548 | 72.5% |
| 1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services. | 66.6% | 68.2% | 67.0% | 0.6% | 3,551 | 5,361 | 66.2% |
| 1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services. | 50.2% | 52.8% | 56.1% | 11.7% | 15,605 | 28,088 | 55.6% |
| 2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy. | 18.5% | 19.2% | 17.8% | -3.7% | 1,910 | 10,879 | 17.6% |
| 2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum. | 70.9% | 70.6% | 74.5% | 5.1% | 3,139 | 4,229 | 74.2% |
| 3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1) | | | | | | | |
| 3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1) | | | | | | | |
| 4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were: | | | | | | | |
| a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1) | 63.9% | 63.5% | 63.2% | -1.2% | 5,954 | 9,381 | 63.5% |
| b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile) | 16.6% | 17.6% | 16.2% | -2.4% | 1,378 | 9,381 | 14.7% |
| c. Obese (BMI at or greater than the 95th percentile). | 16.6% | 16.7% | 15.6% | -5.8% | 1,363 | 9,381 | 14.5% |
| d. Underweight (BMI less than the 5th percentile). | 2.9% | 2.3% | 5.0% | 71.0% | 686 | 9,381 | 7.3% |
| 4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3) | | | | | | | |
| a. Recommended prenatal weight gain | 32.5% | 33.3% | 30.5% | -6.2% | 939 | 3,133 | 30.0% |
| b. Excessive prenatal weight gain | 42.1% | 40.0% | 47.7% | 13.4% | 1,497 | 3,133 | 47.8% |
| c. Inadequate prenatal weight gain | 25.4% | 26.7% | 21.8% | -14.3% | 697 | 3,133 | 22.2% |
| 5A. Percentage of women who participated in WIC who initiated breastfeeding. | 63.6% | 68.8% | 70.2% | 10.3% | 3,936 | 5,420 | 72.6% |
| 5B. Percentage of infants who participated in WIC who were: | | | | | | | |
| a. Breastfeeding at 6 weeks of age (1) | 46.7% | 55.6% | 55.4% | 18.5% | 2,753 | 4,970 | 55.4% |
| b. Breastfeeding at 6 months of age | 30.1% | 33.0% | 29.4% | -2.5% | 1,428 | 4,970 | 28.7% |

(1) Item appears also on other WCH Section Agreement Addenda. Data for 3A and 3B were not presented due to the limitation in recent trend datasets.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI >=30.0): Adapted from the National Academy of Sciences. Weight Gain During Pregnancy: Reexamining the Guidelines. Washington, DC : National Academy Press, 2009.

SFY 2014-2015 AGREEMENT ADDENDA DATA

WIC QUALITY ASSURANCE DELIVERABLES

Warren County

| Process Outcome Objectives | 2004 to 2006 | 2007 to 2009 | 2010 to 2012 | % Change from 2004 to 2012 | 2012 Data | | |
|---|--------------------|--------------------|--------------------|----------------------------------|-----------|-----|---------|
| | | | | | Num | Den | Percent |
| 1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2) | 84.4% | 85.8% | 81.4% | -3.5% | 111 | 131 | 84.7% |
| 1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services. | 76.4% | 77.0% | 76.9% | 0.6% | 116 | 159 | 73.0% |
| 1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services. | 67.7% | 67.5% | 65.7% | -2.9% | 592 | 903 | 65.6% |
| 2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy. | 51.6% | 43.9% | 39.9% | -22.6% | 120 | 307 | 39.1% |
| 2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum. | 78.9% | 77.5% | 70.9% | -10.2% | 108 | 149 | 72.5% |
| 3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1) | | | | | | | |
| 3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1) | | | | | | | |
| 4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were: | | | | | | | |
| a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1) | 71.7% | 70.5% | 64.4% | -10.2% | 259 | 427 | 60.7% |
| b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile) | 11.5% | 11.1% | 12.6% | 9.3% | 55 | 427 | 12.9% |
| c. Obese (BMI at or greater than the 95th percentile). | 10.1% | 9.7% | 12.4% | 22.9% | 54 | 427 | 12.6% |
| d. Underweight (BMI less than the 5th percentile). | 6.7% | 8.8% | 10.6% | 58.4% | 59 | 427 | 13.8% |
| 4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3) | | | | | | | |
| a. Recommended prenatal weight gain | 27.7% | 32.0% | 26.6% | -4.2% | 29 | 117 | 24.8% |
| b. Excessive prenatal weight gain | 45.5% | 40.7% | 48.6% | 6.9% | 54 | 117 | 46.2% |
| c. Inadequate prenatal weight gain | 26.8% | 27.3% | 24.8% | -7.4% | 34 | 117 | 29.1% |
| 5A. Percentage of women who participated in WIC who initiated breastfeeding. | 34.9% | 39.4% | 49.7% | 42.4% | 80 | 171 | 46.8% |
| 5B. Percentage of infants who participated in WIC who were: | | | | | | | |
| a. Breastfeeding at 6 weeks of age (1) | 24.0% | 27.6% | 25.9% | 7.9% | 40 | 163 | 24.5% |
| b. Breastfeeding at 6 months of age | 11.8% | 13.9% | 15.9% | 34.7% | 24 | 163 | 14.7% |

(1) Item appears also on other WCH Section Agreement Addenda. Data for 3A and 3B were not presented due to the limitation in recent trend datasets.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI >=30.0): Adapted from the National Academy of Sciences. Weight Gain During Pregnancy: Reexamining the Guidelines. Washington, DC : National Academy Press, 2009.

SFY 2014-2015 AGREEMENT ADDENDA DATA

WIC QUALITY ASSURANCE DELIVERABLES

Wayne County

| Process Outcome Objectives | 2004 to 2006 | 2007 to 2009 | 2010 to 2012 | % Change from 2004 to 2012 | 2012 Data | | |
|---|--------------------|--------------------|--------------------|----------------------------------|-----------|-------|---------|
| | | | | | Num | Den | Percent |
| 1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2) | 73.8% | 77.0% | 73.0% | -1.0% | 781 | 967 | 80.8% |
| 1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services. | 74.1% | 76.7% | 70.5% | -4.8% | 911 | 1,292 | 70.5% |
| 1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services. | 53.2% | 54.1% | 53.3% | 0.2% | 3,353 | 6,279 | 53.4% |
| 2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy. | 32.9% | 36.1% | 28.5% | -13.2% | 676 | 2,726 | 24.8% |
| 2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum. | 73.0% | 71.2% | 70.0% | -4.1% | 830 | 1,175 | 70.6% |
| 3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1) | | | | | | | |
| 3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1) | | | | | | | |
| 4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were: | | | | | | | |
| a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1) | 63.2% | 65.9% | 65.5% | 3.6% | 1,035 | 1,534 | 67.5% |
| b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile) | 18.0% | 15.9% | 14.6% | -19.2% | 208 | 1,534 | 13.6% |
| c. Obese (BMI at or greater than the 95th percentile). | 16.7% | 15.2% | 15.4% | -7.8% | 208 | 1,534 | 13.6% |
| d. Underweight (BMI less than the 5th percentile). | 2.0% | 3.0% | 4.6% | 122.4% | 83 | 1,534 | 5.4% |
| 4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3) | | | | | | | |
| a. Recommended prenatal weight gain | 30.6% | 32.9% | 28.8% | -5.9% | 247 | 946 | 26.1% |
| b. Excessive prenatal weight gain | 46.8% | 43.4% | 46.4% | -0.8% | 466 | 946 | 49.3% |
| c. Inadequate prenatal weight gain | 22.6% | 23.7% | 24.8% | 9.5% | 233 | 946 | 24.6% |
| 5A. Percentage of women who participated in WIC who initiated breastfeeding. | 52.2% | 48.0% | 54.3% | 4.0% | 795 | 1,396 | 56.9% |
| 5B. Percentage of infants who participated in WIC who were: | | | | | | | |
| a. Breastfeeding at 6 weeks of age (1) | 27.2% | 28.9% | 32.1% | 17.9% | 531 | 1,429 | 37.2% |
| b. Breastfeeding at 6 months of age | 11.4% | 12.4% | 16.3% | 42.7% | 297 | 1,429 | 20.8% |

(1) Item appears also on other WCH Section Agreement Addenda. Data for 3A and 3B were not presented due to the limitation in recent trend datasets.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI < 18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI ≥ 30.0): Adapted from the National Academy of Sciences. Weight Gain During Pregnancy: Reexamining the Guidelines. Washington, DC : National Academy Press, 2009.

SFY 2014-2015 AGREEMENT ADDENDA DATA

WIC QUALITY ASSURANCE DELIVERABLES

Wilkes County

| Process Outcome Objectives | 2004 to 2006 | 2007 to 2009 | 2010 to 2012 | % Change from 2004 to 2012 | 2012 Data | | |
|---|--------------------|--------------------|--------------------|----------------------------------|-----------|-------|---------|
| | | | | | Num | Den | Percent |
| 1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2) | 78.5% | 81.1% | 77.7% | -1.0% | 334 | 436 | 76.6% |
| 1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services. | 79.2% | 80.4% | 76.0% | -4.0% | 416 | 550 | 75.6% |
| 1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services. | 60.8% | 62.9% | 60.4% | -0.7% | 1,583 | 2,758 | 57.4% |
| 2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy. | 43.7% | 37.1% | 42.0% | -3.9% | 340 | 806 | 42.2% |
| 2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum. | 80.8% | 79.3% | 78.5% | -2.9% | 349 | 443 | 78.8% |
| 3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1) | | | | | | | |
| 3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1) | | | | | | | |
| 4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were: | | | | | | | |
| a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1) | 60.5% | 64.4% | 59.8% | -1.2% | 456 | 788 | 57.9% |
| b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile) | 17.9% | 17.9% | 17.8% | -0.7% | 140 | 788 | 17.8% |
| c. Obese (BMI at or greater than the 95th percentile). | 19.5% | 15.8% | 18.0% | -7.7% | 155 | 788 | 19.7% |
| d. Underweight (BMI less than the 5th percentile). | 2.0% | 2.0% | 4.4% | 116.6% | 37 | 788 | 4.7% |
| 4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3) | | | | | | | |
| a. Recommended prenatal weight gain | 29.9% | 32.9% | 29.3% | -2.0% | 104 | 371 | 28.0% |
| b. Excessive prenatal weight gain | 46.0% | 37.7% | 49.0% | 6.5% | 190 | 371 | 51.2% |
| c. Inadequate prenatal weight gain | 24.1% | 29.5% | 21.7% | -9.8% | 77 | 371 | 20.8% |
| 5A. Percentage of women who participated in WIC who initiated breastfeeding. | 49.3% | 57.3% | 56.9% | 15.4% | 290 | 533 | 54.4% |
| 5B. Percentage of infants who participated in WIC who were: | | | | | | | |
| a. Breastfeeding at 6 weeks of age (1) | 24.5% | 24.9% | 29.0% | 18.4% | 168 | 503 | 33.4% |
| b. Breastfeeding at 6 months of age | 11.6% | 10.7% | 15.8% | 35.9% | 96 | 503 | 19.1% |

(1) Item appears also on other WCH Section Agreement Addenda. Data for 3A and 3B were not presented due to the limitation in recent trend datasets.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI >=30.0): Adapted from the National Academy of Sciences. Weight Gain During Pregnancy: Reexamining the Guidelines. Washington, DC : National Academy Press, 2009.

SFY 2014-2015 AGREEMENT ADDENDA DATA

WIC QUALITY ASSURANCE DELIVERABLES

Wilson County

| Process Outcome Objectives | 2004 to 2006 | 2007 to 2009 | 2010 to 2012 | % Change from 2004 to 2012 | 2012 Data | | |
|---|--------------------|--------------------|--------------------|----------------------------------|-----------|-------|---------|
| | | | | | Num | Den | Percent |
| 1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2) | 83.8% | 84.3% | 83.6% | -0.2% | 436 | 535 | 81.5% |
| 1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services. | 77.3% | 76.1% | 75.5% | -2.3% | 520 | 695 | 74.8% |
| 1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services. | 56.7% | 58.2% | 56.7% | 0.0% | 2,393 | 4,210 | 56.8% |
| 2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy. | 28.0% | 22.8% | 20.7% | -26.1% | 346 | 1,743 | 19.9% |
| 2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum. | 67.1% | 69.5% | 73.9% | 10.2% | 547 | 667 | 82.0% |
| 3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1) | | | | | | | |
| 3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1) | | | | | | | |
| 4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were: | | | | | | | |
| a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1) | 62.1% | 62.4% | 62.9% | 1.3% | 914 | 1,474 | 62.0% |
| b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile) | 17.6% | 18.3% | 15.5% | -11.8% | 233 | 1,474 | 15.8% |
| c. Obese (BMI at or greater than the 95th percentile). | 17.8% | 16.5% | 16.4% | -8.0% | 221 | 1,474 | 15.0% |
| d. Underweight (BMI less than the 5th percentile). | 2.5% | 2.7% | 5.2% | 104.8% | 106 | 1,474 | 7.2% |
| 4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3) | | | | | | | |
| a. Recommended prenatal weight gain | 31.5% | 31.6% | 32.9% | 4.5% | 150 | 494 | 30.4% |
| b. Excessive prenatal weight gain | 40.6% | 39.1% | 44.3% | 9.0% | 219 | 494 | 44.3% |
| c. Inadequate prenatal weight gain | 27.9% | 29.3% | 22.9% | -18.2% | 125 | 494 | 25.3% |
| 5A. Percentage of women who participated in WIC who initiated breastfeeding. | 41.4% | 38.5% | 42.3% | 2.1% | 379 | 826 | 45.9% |
| 5B. Percentage of infants who participated in WIC who were: | | | | | | | |
| a. Breastfeeding at 6 weeks of age (1) | 23.6% | 25.3% | 24.6% | 4.1% | 220 | 812 | 27.1% |
| b. Breastfeeding at 6 months of age | 12.0% | 13.6% | 16.6% | 38.2% | 125 | 812 | 15.4% |

(1) Item appears also on other WCH Section Agreement Addenda. Data for 3A and 3B were not presented due to the limitation in recent trend datasets.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI >=30.0): Adapted from the National Academy of Sciences. Weight Gain During Pregnancy: Reexamining the Guidelines. Washington, DC : National Academy Press, 2009.

SFY 2014-2015 AGREEMENT ADDENDA DATA

WIC QUALITY ASSURANCE DELIVERABLES

Yadkin County

| Process Outcome Objectives | 2004 to 2006 | 2007 to 2009 | 2010 to 2012 | % Change from 2004 to 2012 | 2012 Data | | |
|---|--------------------|--------------------|--------------------|----------------------------------|-----------|-------|---------|
| | | | | | Num | Den | Percent |
| 1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2) | 72.9% | 73.8% | 77.2% | 5.8% | 186 | 235 | 79.1% |
| 1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services. | 68.4% | 73.2% | 79.7% | 16.4% | 253 | 314 | 80.6% |
| 1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services. | 53.4% | 62.4% | 64.3% | 20.3% | 1,068 | 1,620 | 65.9% |
| 2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy. | 46.6% | 45.1% | 41.6% | -10.7% | 192 | 503 | 38.2% |
| 2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum. | 74.3% | 77.7% | 76.7% | 3.3% | 207 | 270 | 76.7% |
| 3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1) | | | | | | | |
| 3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1) | | | | | | | |
| 4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were: | | | | | | | |
| a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1) | 62.2% | 64.8% | 60.9% | -2.1% | 361 | 614 | 58.8% |
| b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile) | 16.7% | 15.3% | 17.1% | 2.3% | 104 | 614 | 16.9% |
| c. Obese (BMI at or greater than the 95th percentile). | 17.8% | 18.3% | 16.5% | -7.3% | 86 | 614 | 14.0% |
| d. Underweight (BMI less than the 5th percentile). | 3.4% | 1.7% | 5.6% | 65.7% | 63 | 614 | 10.3% |
| 4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3) | | | | | | | |
| a. Recommended prenatal weight gain | 33.3% | 33.5% | 30.3% | -9.0% | 63 | 224 | 28.1% |
| b. Excessive prenatal weight gain | 47.5% | 47.0% | 46.9% | -1.3% | 110 | 224 | 49.1% |
| c. Inadequate prenatal weight gain | 19.2% | 19.5% | 22.8% | 18.7% | 51 | 224 | 22.8% |
| 5A. Percentage of women who participated in WIC who initiated breastfeeding. | 66.1% | 71.0% | 67.6% | 2.2% | 209 | 325 | 64.3% |
| 5B. Percentage of infants who participated in WIC who were: | | | | | | | |
| a. Breastfeeding at 6 weeks of age (1) | 37.8% | 44.5% | 41.5% | 9.8% | 133 | 335 | 39.7% |
| b. Breastfeeding at 6 months of age | 15.7% | 23.0% | 22.0% | 40.4% | 67 | 335 | 20.0% |

(1) Item appears also on other WCH Section Agreement Addenda. Data for 3A and 3B were not presented due to the limitation in recent trend datasets.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI >=30.0): Adapted from the National Academy of Sciences. Weight Gain During Pregnancy: Reexamining the Guidelines. Washington, DC : National Academy Press, 2009.

SFY 2014-2015 AGREEMENT ADDENDA DATA
WIC QUALITY ASSURANCE DELIVERABLES

Allegheny

| Process Outcome Objectives | 2004 to 2006 | 2007 to 2009 | 2010 to 2012 | % Change from 2004 to 2012 | 2012 Data | | |
|---|--------------------|--------------------|--------------------|----------------------------------|-----------|-----|---------|
| | | | | | Num | Den | Percent |
| 1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2) | 79.7% | 79.9% | 70.6% | -11.4% | 45 | 59 | 76.3% |
| 1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services. | 79.9% | 78.2% | 77.5% | -3.0% | 48 | 74 | 64.9% |
| 1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services. | 69.4% | 69.1% | 70.6% | 1.8% | 284 | 391 | 72.6% |
| 2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy. | 46.1% | 45.8% | 41.4% | -10.1% | 60 | 139 | 43.2% |
| 2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum. | 74.0% | 75.2% | 73.6% | -0.5% | 44 | 64 | 68.8% |
| 3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1) | | | | | | | |
| 3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1) | | | | | | | |
| 4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were: | | | | | | | |
| a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1) | 58.0% | 54.6% | 56.6% | -2.3% | 132 | 216 | 61.1% |
| b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile) | 19.2% | 19.1% | 18.0% | -6.0% | 28 | 216 | 13.0% |
| c. Obese (BMI at or greater than the 95th percentile). | 21.6% | 24.3% | 21.7% | 0.3% | 42 | 216 | 19.4% |
| d. Underweight (BMI less than the 5th percentile). | 1.2% | 1.9% | 3.6% | 200.2% | 14 | 216 | 6.5% |
| 4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3) | | | | | | | |
| a. Recommended prenatal weight gain | 38.2% | 30.8% | 27.4% | -28.3% | 9 | 45 | 20.0% |
| b. Excessive prenatal weight gain | 43.5% | 55.9% | 47.0% | 8.1% | 24 | 45 | 53.3% |
| c. Inadequate prenatal weight gain | 18.3% | 13.3% | 25.6% | 39.7% | 12 | 45 | 26.7% |
| 5A. Percentage of women who participated in WIC who initiated breastfeeding. | 60.4% | 52.3% | 46.8% | -22.4% | 56 | 91 | 61.5% |
| 5B. Percentage of infants who participated in WIC who were: | | | | | | | |
| a. Breastfeeding at 6 weeks of age (1) | 18.5% | 25.3% | 16.2% | -12.1% | 19 | 86 | 22.1% |
| b. Breastfeeding at 6 months of age | 8.0% | 13.6% | 9.7% | 22.3% | 15 | 86 | 17.4% |

(1) Item appears also on other WCH Section Agreement Addenda. Data for 3A and 3B are not presented due to limitation in recent trend data.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI >=30.0); Adapted from the National Academy of Sciences. Weight Gain During Pregnancy: Reexamining the Guidelines. Washington, DC : National Academy Press, 2009.

SFY 2014-2015 AGREEMENT ADDENDA DATA

WIC QUALITY ASSURANCE DELIVERABLES

Ashe

| Process Outcome Objectives | 2004 to 2006 | 2007 to 2009 | 2010 to 2012 | % Change from 2004 to 2012 | 2012 Data | | |
|---|--------------------|--------------------|--------------------|----------------------------------|-----------|-----|---------|
| | | | | | Num | Den | Percent |
| 1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2) | 75.8% | 74.9% | 74.8% | -1.3% | 137 | 168 | 81.5% |
| 1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services. | 71.5% | 73.0% | 71.5% | -0.1% | 151 | 196 | 77.0% |
| 1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services. | 65.6% | 63.5% | 61.1% | -7.0% | 573 | 914 | 62.7% |
| 2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy. | 53.4% | 49.0% | 40.5% | -24.2% | 112 | 302 | 37.1% |
| 2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum. | 78.9% | 80.0% | 77.5% | -1.8% | 126 | 153 | 82.4% |
| 3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1) | | | | | | | |
| 3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1) | | | | | | | |
| 4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were: | | | | | | | |
| a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1) | 68.6% | 63.9% | 66.1% | -3.7% | 177 | 272 | 65.1% |
| b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile) | 11.7% | 16.4% | 14.7% | 25.2% | 38 | 272 | 14.0% |
| c. Obese (BMI at or greater than the 95th percentile). | 13.7% | 14.4% | 13.6% | -0.5% | 37 | 272 | 13.6% |
| d. Underweight (BMI less than the 5th percentile). | 6.0% | 5.3% | 5.6% | -6.0% | 20 | 272 | 7.4% |
| 4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3) | | | | | | | |
| a. Recommended prenatal weight gain | 31.0% | 28.4% | 32.1% | 3.5% | 42 | 137 | 30.7% |
| b. Excessive prenatal weight gain | 45.3% | 45.9% | 47.2% | 4.2% | 67 | 137 | 48.9% |
| c. Inadequate prenatal weight gain | 23.6% | 25.8% | 20.7% | -12.5% | 28 | 137 | 20.4% |
| 5A. Percentage of women who participated in WIC who initiated breastfeeding. | 48.8% | 51.9% | 52.9% | 8.3% | 78 | 191 | 40.8% |
| 5B. Percentage of infants who participated in WIC who were: | | | | | | | |
| a. Breastfeeding at 6 weeks of age (1) | 30.6% | 28.1% | 31.6% | 3.3% | 64 | 189 | 33.9% |
| b. Breastfeeding at 6 months of age | 12.9% | 14.1% | 21.5% | 67.2% | 46 | 189 | 24.3% |

(1) Item appears also on other WCH Section Agreement Addenda. Data for 3A and 3B are not presented due to limitation in recent trend data.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI >=30.0); Adapted from the National Academy of Sciences. Weight Gain During Pregnancy: Reexamining the Guidelines. Washington, DC : National Academy Press, 2009.

SFY 2014-2015 AGREEMENT ADDENDA DATA

WIC QUALITY ASSURANCE DELIVERABLES

Avery

| Process Outcome Objectives | 2004 to 2006 | 2007 to 2009 | 2010 to 2012 | % Change from 2004 to 2012 | 2012 Data | | |
|---|--------------------|--------------------|--------------------|----------------------------------|-----------|-----|---------|
| | | | | | Num | Den | Percent |
| 1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2) | 73.8% | 73.7% | 72.8% | -1.4% | 73 | 91 | 80.2% |
| 1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services. | 65.1% | 66.8% | 68.9% | 5.9% | 78 | 110 | 70.9% |
| 1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services. | 61.4% | 62.7% | 65.5% | 6.6% | 374 | 563 | 66.4% |
| 2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy. | 44.8% | 47.9% | 41.8% | -6.7% | 100 | 250 | 40.0% |
| 2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum. | 68.7% | 69.8% | 74.6% | 8.6% | 86 | 102 | 84.3% |
| 3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1) | | | | | | | |
| 3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1) | | | | | | | |
| 4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were: | | | | | | | |
| a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1) | 64.7% | 63.3% | 66.8% | 3.2% | 167 | 258 | 64.7% |
| b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile) | 16.1% | 19.4% | 14.1% | -12.5% | 37 | 258 | 14.3% |
| c. Obese (BMI at or greater than the 95th percentile). | 17.3% | 14.0% | 12.0% | -30.4% | 30 | 258 | 11.6% |
| d. Underweight (BMI less than the 5th percentile). | 1.8% | 3.2% | 7.1% | 281.7% | 24 | 258 | 9.3% |
| 4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3) | | | | | | | |
| a. Recommended prenatal weight gain | 30.1% | 31.5% | 33.2% | 10.4% | 24 | 71 | 33.8% |
| b. Excessive prenatal weight gain | 44.9% | 45.3% | 46.5% | 3.5% | 29 | 71 | 40.8% |
| c. Inadequate prenatal weight gain | 25.0% | 23.2% | 20.3% | -18.8% | 18 | 71 | 25.4% |
| 5A. Percentage of women who participated in WIC who initiated breastfeeding. | 49.3% | 60.7% | 61.7% | 25.2% | 92 | 127 | 72.4% |
| 5B. Percentage of infants who participated in WIC who were: | | | | | | | |
| a. Breastfeeding at 6 weeks of age (1) | 34.4% | 35.9% | 41.6% | 21.0% | 55 | 111 | 49.5% |
| b. Breastfeeding at 6 months of age | 26.4% | 24.0% | 27.3% | 3.3% | 37 | 111 | 33.3% |

(1) Item appears also on other WCH Section Agreement Addenda. Data for 3A and 3B are not presented due to limitation in recent trend data.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI >=30.0); Adapted from the National Academy of Sciences. Weight Gain During Pregnancy: Reexamining the Guidelines. Washington, DC : National Academy Press, 2009.

SFY 2014-2015 AGREEMENT ADDENDA DATA

WIC QUALITY ASSURANCE DELIVERABLES

Bertie

| Process Outcome Objectives | 2004 to 2006 | 2007 to 2009 | 2010 to 2012 | % Change from 2004 to 2012 | 2012 Data | | |
|---|--------------------|--------------------|--------------------|----------------------------------|-----------|-----|---------|
| | | | | | Num | Den | Percent |
| 1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2) | 82.8% | 80.3% | 80.8% | -2.4% | 119 | 139 | 85.6% |
| 1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services. | 74.9% | 70.4% | 72.7% | -2.9% | 109 | 146 | 74.7% |
| 1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services. | 62.7% | 60.8% | 58.7% | -6.3% | 493 | 828 | 59.5% |
| 2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy. | 36.2% | 24.7% | 26.4% | -27.0% | 96 | 346 | 27.7% |
| 2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum. | 33.9% | 24.9% | 45.5% | 34.3% | 100 | 146 | 68.5% |
| 3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1) | | | | | | | |
| 3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1) | | | | | | | |
| 4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were: | | | | | | | |
| a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1) | 65.1% | 63.4% | 64.8% | -0.3% | 405 | 630 | 64.3% |
| b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile) | 13.2% | 13.3% | 11.6% | -12.2% | 68 | 630 | 10.8% |
| c. Obese (BMI at or greater than the 95th percentile). | 14.5% | 17.3% | 14.4% | -0.7% | 97 | 630 | 15.4% |
| d. Underweight (BMI less than the 5th percentile). | 7.3% | 6.0% | 9.2% | 26.5% | 60 | 630 | 9.5% |
| 4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3) | | | | | | | |
| a. Recommended prenatal weight gain | 28.1% | 28.1% | 25.1% | -10.7% | 22 | 119 | 18.5% |
| b. Excessive prenatal weight gain | 49.3% | 47.1% | 26.9% | -45.4% | 33 | 119 | 27.7% |
| c. Inadequate prenatal weight gain | 22.7% | 24.7% | 48.0% | 112.0% | 64 | 119 | 53.8% |
| 5A. Percentage of women who participated in WIC who initiated breastfeeding. | 15.5% | 13.5% | 40.6% | 162.2% | 90 | 173 | 52.0% |
| 5B. Percentage of infants who participated in WIC who were: | | | | | | | |
| a. Breastfeeding at 6 weeks of age (1) | 7.8% | 6.2% | 8.6% | 9.4% | 18 | 182 | 9.9% |
| b. Breastfeeding at 6 months of age | 2.4% | 1.8% | 1.6% | -33.3% | 2 | 182 | 1.1% |

(1) Item appears also on other WCH Section Agreement Addenda. Data for 3A and 3B are not presented due to limitation in recent trend data.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI >=30.0); Adapted from the National Academy of Sciences. Weight Gain During Pregnancy: Reexamining the Guidelines. Washington, DC : National Academy Press, 2009.

SFY 2014-2015 AGREEMENT ADDENDA DATA
WIC QUALITY ASSURANCE DELIVERABLES

Camden

| Process Outcome Objectives | 2004 to 2006 | 2007 to 2009 | 2010 to 2012 | % Change from 2004 to 2012 | 2012 Data | | |
|---|--------------------|--------------------|--------------------|----------------------------------|-----------|-------|---------|
| | | | | | Num | Den | Percent |
| 1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2) | 75.0% | 85.4% | 71.4% | -4.8% | 18 | 27 | 66.7% |
| 1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services. | 69.2% | 75.8% | 72.8% | 5.2% | 22 | 30 | 73.3% |
| 1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services. | 56.4% | 59.2% | 52.6% | -6.6% | 95 | 178 | 53.4% |
| 2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy. | 36.3% | 35.1% | 27.0% | -25.6% | 463 | 1,750 | 26.5% |
| 2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum. | 33.5% | 41.9% | 58.0% | 73.3% | 509 | 681 | 74.7% |
| 3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1) | | | | | | | |
| 3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1) | | | | | | | |
| 4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were: | | | | | | | |
| a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1) | 69.3% | 66.3% | 63.0% | -9.1% | 255 | 419 | 60.9% |
| b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile) | 6.0% | 10.4% | 11.1% | 84.4% | 48 | 419 | 11.5% |
| c. Obese (BMI at or greater than the 95th percentile). | 10.0% | 13.9% | 16.6% | 65.9% | 71 | 419 | 16.9% |
| d. Underweight (BMI less than the 5th percentile). | 14.6% | 9.4% | 9.2% | -36.7% | 45 | 419 | 10.7% |
| 4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3) | | | | | | | |
| a. Recommended prenatal weight gain | 34.5% | 27.6% | 27.4% | -20.6% | 7 | 24 | 29.2% |
| b. Excessive prenatal weight gain | 33.3% | 48.0% | 48.8% | 46.4% | 11 | 24 | 45.8% |
| c. Inadequate prenatal weight gain | 32.2% | 24.5% | 23.8% | -26.0% | 6 | 24 | 25.0% |
| 5A. Percentage of women who participated in WIC who initiated breastfeeding. | 20.8% | 23.9% | 32.6% | 56.9% | 275 | 857 | 32.1% |
| 5B. Percentage of infants who participated in WIC who were: | | | | | | | |
| a. Breastfeeding at 6 weeks of age (1) | 8.7% | 10.8% | 17.3% | 97.5% | 169 | 714 | 23.7% |
| b. Breastfeeding at 6 months of age | 3.4% | 3.4% | 9.2% | 172.6% | 85 | 714 | 11.9% |

(1) Item appears also on other WCH Section Agreement Addenda. Data for 3A and 3B are not presented due to limitation in recent trend data.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI >=30.0); Adapted from the National Academy of Sciences. Weight Gain During Pregnancy: Reexamining the Guidelines. Washington, DC : National Academy Press, 2009.

SFY 2014-2015 AGREEMENT ADDENDA DATA
WIC QUALITY ASSURANCE DELIVERABLES

Chowan

| Process Outcome Objectives | 2004 to 2006 | 2007 to 2009 | 2010 to 2012 | % Change from 2004 to 2012 | 2012 Data | | |
|---|--------------------|--------------------|--------------------|----------------------------------|-----------|-------|---------|
| | | | | | Num | Den | Percent |
| 1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2) | 82.6% | 78.9% | 78.0% | -5.6% | 68 | 93 | 73.1% |
| 1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services. | 84.5% | 83.6% | 84.5% | 0.0% | 85 | 101 | 84.2% |
| 1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services. | 66.9% | 67.0% | 69.8% | 4.3% | 393 | 546 | 72.0% |
| 2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy. | 36.3% | 35.1% | 27.0% | -25.6% | 463 | 1,750 | 26.5% |
| 2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum. | 33.5% | 41.9% | 58.0% | 73.3% | 509 | 681 | 74.7% |
| 3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1) | | | | | | | |
| 3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1) | | | | | | | |
| 4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were: | | | | | | | |
| a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1) | 62.0% | 68.5% | 58.9% | -5.0% | 300 | 508 | 59.1% |
| b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile) | 17.6% | 14.2% | 13.9% | -21.3% | 69 | 508 | 13.6% |
| c. Obese (BMI at or greater than the 95th percentile). | 18.1% | 13.9% | 18.3% | 0.8% | 87 | 508 | 17.1% |
| d. Underweight (BMI less than the 5th percentile). | 2.2% | 3.4% | 8.9% | 299.8% | 52 | 508 | 10.2% |
| 4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3) | | | | | | | |
| a. Recommended prenatal weight gain | 31.6% | 33.2% | 29.8% | -5.6% | 25 | 74 | 33.8% |
| b. Excessive prenatal weight gain | 43.8% | 43.0% | 45.1% | 3.1% | 27 | 74 | 36.5% |
| c. Inadequate prenatal weight gain | 24.7% | 23.8% | 25.1% | 1.8% | 22 | 74 | 29.7% |
| 5A. Percentage of women who participated in WIC who initiated breastfeeding. | 20.8% | 23.9% | 32.6% | 56.9% | 275 | 857 | 32.1% |
| 5B. Percentage of infants who participated in WIC who were: | | | | | | | |
| a. Breastfeeding at 6 weeks of age (1) | 8.7% | 10.8% | 17.3% | 97.5% | 169 | 714 | 23.7% |
| b. Breastfeeding at 6 months of age | 3.4% | 3.4% | 9.2% | 172.6% | 85 | 714 | 11.9% |

(1) Item appears also on other WCH Section Agreement Addenda. Data for 3A and 3B are not presented due to limitation in recent trend data.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI >=30.0); Adapted from the National Academy of Sciences. Weight Gain During Pregnancy: Reexamining the Guidelines. Washington, DC : National Academy Press, 2009.

SFY 2014-2015 AGREEMENT ADDENDA DATA

WIC QUALITY ASSURANCE DELIVERABLES

Currituck

| Process Outcome Objectives | 2004 to 2006 | 2007 to 2009 | 2010 to 2012 | % Change from 2004 to 2012 | 2012 Data | | |
|---|--------------------|--------------------|--------------------|----------------------------------|-----------|-------|---------|
| | | | | | Num | Den | Percent |
| 1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2) | 73.6% | 73.9% | 58.6% | -20.4% | 47 | 95 | 49.5% |
| 1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services. | 71.8% | 66.2% | 69.5% | -3.2% | 79 | 110 | 71.8% |
| 1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services. | 54.5% | 50.7% | 51.4% | -5.7% | 327 | 596 | 54.9% |
| 2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy. | 36.3% | 35.1% | 27.0% | -25.6% | 463 | 1,750 | 26.5% |
| 2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum. | 33.5% | 41.9% | 58.0% | 73.3% | 509 | 681 | 74.7% |
| 3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1) | | | | | | | |
| 3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1) | | | | | | | |
| 4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were: | | | | | | | |
| a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1) | 66.3% | 67.8% | 62.6% | -5.6% | 285 | 478 | 59.6% |
| b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile) | 17.6% | 14.3% | 15.1% | -14.4% | 66 | 478 | 13.8% |
| c. Obese (BMI at or greater than the 95th percentile). | 12.6% | 15.7% | 15.3% | 21.5% | 84 | 478 | 17.6% |
| d. Underweight (BMI less than the 5th percentile). | 3.5% | 2.2% | 7.0% | 102.9% | 43 | 478 | 9.0% |
| 4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3) | | | | | | | |
| a. Recommended prenatal weight gain | 36.3% | 33.9% | 22.9% | -37.0% | 11 | 64 | 17.2% |
| b. Excessive prenatal weight gain | 47.9% | 45.0% | 51.1% | 6.5% | 40 | 64 | 62.5% |
| c. Inadequate prenatal weight gain | 15.8% | 21.1% | 26.1% | 65.4% | 13 | 64 | 20.3% |
| 5A. Percentage of women who participated in WIC who initiated breastfeeding. | 20.8% | 23.9% | 32.6% | 56.9% | 275 | 857 | 32.1% |
| 5B. Percentage of infants who participated in WIC who were: | | | | | | | |
| a. Breastfeeding at 6 weeks of age (1) | 8.7% | 10.8% | 17.3% | 97.5% | 169 | 714 | 23.7% |
| b. Breastfeeding at 6 months of age | 3.4% | 3.4% | 9.2% | 172.6% | 85 | 714 | 11.9% |

(1) Item appears also on other WCH Section Agreement Addenda. Data for 3A and 3B are not presented due to limitation in recent trend data.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI >=30.0); Adapted from the National Academy of Sciences. Weight Gain During Pregnancy: Reexamining the Guidelines. Washington, DC : National Academy Press, 2009.

SFY 2014-2015 AGREEMENT ADDENDA DATA
WIC QUALITY ASSURANCE DELIVERABLES

Gates

| Process Outcome Objectives | 2004 to 2006 | 2007 to 2009 | 2010 to 2012 | % Change from 2004 to 2012 | 2012 Data | | |
|---|--------------------|--------------------|--------------------|----------------------------------|-----------|-----|---------|
| | | | | | Num | Den | Percent |
| 1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2) | 84.2% | 78.3% | 48.1% | -42.9% | 28 | 55 | 50.9% |
| 1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services. | 70.6% | 74.9% | 81.7% | 15.6% | 65 | 78 | 83.3% |
| 1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services. | 59.2% | 61.9% | 63.8% | 7.7% | 236 | 353 | 66.9% |
| 2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy. | 31.7% | 25.9% | 25.3% | -20.0% | 32 | 134 | 23.9% |
| 2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum. | 33.5% | 67.7% | 70.8% | 111.1% | 45 | 67 | 67.2% |
| 3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1) | | | | | | | |
| 3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1) | | | | | | | |
| 4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were: | | | | | | | |
| a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1) | 65.5% | 64.3% | 58.4% | -10.9% | 199 | 350 | 56.9% |
| b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile) | 15.8% | 15.5% | 12.6% | -20.3% | 43 | 350 | 12.3% |
| c. Obese (BMI at or greater than the 95th percentile). | 15.5% | 16.5% | 17.8% | 14.8% | 68 | 350 | 19.4% |
| d. Underweight (BMI less than the 5th percentile). | 3.2% | 3.7% | 11.3% | 250.7% | 40 | 350 | 11.4% |
| 4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3) | | | | | | | |
| a. Recommended prenatal weight gain | 28.0% | 40.0% | 32.1% | 14.6% | 10 | 34 | 29.4% |
| b. Excessive prenatal weight gain | 43.9% | 40.0% | 39.3% | -10.5% | 17 | 34 | 50.0% |
| c. Inadequate prenatal weight gain | 28.0% | 20.0% | 28.6% | 1.9% | 7 | 34 | 20.6% |
| 5A. Percentage of women who participated in WIC who initiated breastfeeding. | 32.4% | 23.8% | 19.2% | -40.8% | 23 | 81 | 28.4% |
| 5B. Percentage of infants who participated in WIC who were: | | | | | | | |
| a. Breastfeeding at 6 weeks of age (1) | 9.8% | 12.9% | 9.7% | -1.2% | 9 | 96 | 9.4% |
| b. Breastfeeding at 6 months of age | 4.9% | 2.7% | 2.7% | -44.7% | 3 | 96 | 3.1% |

(1) Item appears also on other WCH Section Agreement Addenda. Data for 3A and 3B are not presented due to limitation in recent trend data.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI >=30.0); Adapted from the National Academy of Sciences. Weight Gain During Pregnancy: Reexamining the Guidelines. Washington, DC : National Academy Press, 2009.

SFY 2014-2015 AGREEMENT ADDENDA DATA

WIC QUALITY ASSURANCE DELIVERABLES

Granville

| Process Outcome Objectives | 2004 to 2006 | 2007 to 2009 | 2010 to 2012 | % Change from 2004 to 2012 | 2012 Data | | |
|---|--------------------|--------------------|--------------------|----------------------------------|-----------|-------|---------|
| | | | | | Num | Den | Percent |
| 1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2) | 75.8% | 75.8% | 74.4% | -1.9% | 221 | 301 | 73.4% |
| 1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services. | 77.2% | 76.0% | 75.1% | -2.8% | 269 | 362 | 74.3% |
| 1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services. | 59.1% | 60.0% | 59.5% | 0.6% | 1,116 | 1,874 | 59.6% |
| 2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy. | 31.0% | 33.2% | 27.7% | -10.5% | 430 | 1,554 | 27.7% |
| 2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum. | 80.2% | 75.2% | 76.7% | -4.4% | 556 | 707 | 78.6% |
| 3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1) | | | | | | | |
| 3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1) | | | | | | | |
| 4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were: | | | | | | | |
| a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1) | 62.9% | 62.3% | 65.8% | 4.6% | 436 | 662 | 65.9% |
| b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile) | 17.3% | 17.8% | 14.0% | -19.1% | 91 | 662 | 13.7% |
| c. Obese (BMI at or greater than the 95th percentile). | 16.3% | 17.1% | 15.8% | -2.9% | 104 | 662 | 15.7% |
| d. Underweight (BMI less than the 5th percentile). | 3.5% | 2.7% | 4.4% | 26.2% | 31 | 662 | 4.7% |
| 4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3) | | | | | | | |
| a. Recommended prenatal weight gain | 31.3% | 31.1% | 30.0% | -4.1% | 83 | 276 | 30.1% |
| b. Excessive prenatal weight gain | 38.7% | 45.6% | 49.6% | 28.0% | 138 | 276 | 50.0% |
| c. Inadequate prenatal weight gain | 29.9% | 23.3% | 20.4% | -31.9% | 55 | 276 | 19.9% |
| 5A. Percentage of women who participated in WIC who initiated breastfeeding. | 40.4% | 52.0% | 45.6% | 13.0% | 438 | 893 | 49.0% |
| 5B. Percentage of infants who participated in WIC who were: | | | | | | | |
| a. Breastfeeding at 6 weeks of age (1) | 28.0% | 38.1% | 35.4% | 26.2% | 337 | 928 | 36.3% |
| b. Breastfeeding at 6 months of age | 19.5% | 26.5% | 25.8% | 32.3% | 256 | 928 | 27.6% |

(1) Item appears also on other WCH Section Agreement Addenda. Data for 3A and 3B are not presented due to limitation in recent trend data.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI >=30.0); Adapted from the National Academy of Sciences. Weight Gain During Pregnancy: Reexamining the Guidelines. Washington, DC : National Academy Press, 2009.

SFY 2014-2015 AGREEMENT ADDENDA DATA

WIC QUALITY ASSURANCE DELIVERABLES

Martin

| Process Outcome Objectives | 2004 to 2006 | 2007 to 2009 | 2010 to 2012 | % Change from 2004 to 2012 | 2012 Data | | |
|---|--------------------|--------------------|--------------------|----------------------------------|-----------|-------|---------|
| | | | | | Num | Den | Percent |
| 1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2) | 75.5% | 84.0% | 82.8% | 9.7% | 141 | 173 | 81.5% |
| 1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services. | 73.3% | 68.5% | 82.2% | 12.2% | 159 | 185 | 85.9% |
| 1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services. | 58.0% | 59.4% | 60.0% | 3.4% | 649 | 1,052 | 61.7% |
| 2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy. | 35.3% | 33.0% | 28.0% | -20.7% | 269 | 995 | 27.0% |
| 2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum. | 81.3% | 82.0% | 80.5% | -1.0% | 268 | 320 | 83.8% |
| 3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1) | | | | | | | |
| 3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1) | | | | | | | |
| 4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were: | | | | | | | |
| a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1) | 66.0% | 63.2% | 62.7% | -5.0% | 261 | 458 | 57.0% |
| b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile) | 15.3% | 15.0% | 14.6% | -4.1% | 74 | 458 | 16.2% |
| c. Obese (BMI at or greater than the 95th percentile). | 12.9% | 17.1% | 17.1% | 32.7% | 93 | 458 | 20.3% |
| d. Underweight (BMI less than the 5th percentile). | 5.9% | 4.7% | 5.6% | -5.2% | 30 | 458 | 6.6% |
| 4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3) | | | | | | | |
| a. Recommended prenatal weight gain | 32.1% | 31.0% | 25.0% | -22.2% | 29 | 133 | 21.8% |
| b. Excessive prenatal weight gain | 43.3% | 41.7% | 46.0% | 6.1% | 64 | 133 | 48.1% |
| c. Inadequate prenatal weight gain | 24.5% | 27.3% | 29.0% | 18.3% | 40 | 133 | 30.1% |
| 5A. Percentage of women who participated in WIC who initiated breastfeeding. | 40.3% | 36.6% | 23.0% | -43.0% | 90 | 388 | 23.2% |
| 5B. Percentage of infants who participated in WIC who were: | | | | | | | |
| a. Breastfeeding at 6 weeks of age (1) | 25.5% | 21.6% | 11.2% | -56.3% | 32 | 351 | 9.1% |
| b. Breastfeeding at 6 months of age | 12.0% | 9.5% | 4.3% | -64.4% | 17 | 351 | 4.8% |

(1) Item appears also on other WCH Section Agreement Addenda. Data for 3A and 3B are not presented due to limitation in recent trend data.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI >=30.0); Adapted from the National Academy of Sciences. Weight Gain During Pregnancy: Reexamining the Guidelines. Washington, DC : National Academy Press, 2009.

SFY 2014-2015 AGREEMENT ADDENDA DATA

WIC QUALITY ASSURANCE DELIVERABLES

McDowell

| Process Outcome Objectives | 2004 to 2006 | 2007 to 2009 | 2010 to 2012 | % Change from 2004 to 2012 | 2012 Data | | |
|---|--------------------|--------------------|--------------------|----------------------------------|-----------|-------|---------|
| | | | | | Num | Den | Percent |
| 1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2) | 66.5% | 74.6% | 74.1% | 11.5% | 247 | 300 | 82.3% |
| 1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services. | 70.6% | 75.1% | 73.9% | 4.7% | 266 | 368 | 72.3% |
| 1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services. | 63.7% | 63.5% | 62.1% | -2.5% | 1,240 | 1,987 | 62.4% |
| 2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy. | 32.9% | 35.4% | 33.2% | 1.0% | 847 | 2,654 | 31.9% |
| 2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum. | 71.0% | 73.2% | 78.0% | 9.8% | 785 | 990 | 79.3% |
| 3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1) | | | | | | | |
| 3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1) | | | | | | | |
| 4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were: | | | | | | | |
| a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1) | 68.3% | 68.4% | 66.8% | -2.1% | 410 | 637 | 64.4% |
| b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile) | 14.5% | 15.2% | 14.2% | -2.1% | 94 | 637 | 14.8% |
| c. Obese (BMI at or greater than the 95th percentile). | 12.7% | 12.6% | 12.9% | 2.0% | 104 | 637 | 16.3% |
| d. Underweight (BMI less than the 5th percentile). | 4.6% | 3.8% | 6.1% | 32.2% | 29 | 637 | 4.6% |
| 4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3) | | | | | | | |
| a. Recommended prenatal weight gain | 34.2% | 29.4% | 28.8% | -15.9% | 70 | 259 | 27.0% |
| b. Excessive prenatal weight gain | 34.9% | 43.1% | 49.7% | 42.5% | 132 | 259 | 51.0% |
| c. Inadequate prenatal weight gain | 30.9% | 27.5% | 21.5% | -30.3% | 57 | 259 | 22.0% |
| 5A. Percentage of women who participated in WIC who initiated breastfeeding. | 49.2% | 56.9% | 59.1% | 20.2% | 670 | 1,190 | 56.3% |
| 5B. Percentage of infants who participated in WIC who were: | | | | | | | |
| a. Breastfeeding at 6 weeks of age (1) | 30.0% | 29.7% | 32.2% | 7.2% | 374 | 1,081 | 34.6% |
| b. Breastfeeding at 6 months of age | 15.2% | 14.0% | 17.3% | 13.9% | 185 | 1,081 | 17.1% |

(1) Item appears also on other WCH Section Agreement Addenda. Data for 3A and 3B are not presented due to limitation in recent trend data.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI >=30.0); Adapted from the National Academy of Sciences. Weight Gain During Pregnancy: Reexamining the Guidelines. Washington, DC : National Academy Press, 2009.

SFY 2014-2015 AGREEMENT ADDENDA DATA

WIC QUALITY ASSURANCE DELIVERABLES

Mitchell

| Process Outcome Objectives | 2004 to 2006 | 2007 to 2009 | 2010 to 2012 | % Change from 2004 to 2012 | 2012 Data | | |
|---|--------------------|--------------------|--------------------|----------------------------------|-----------|-----|---------|
| | | | | | Num | Den | Percent |
| 1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2) | 79.3% | 74.1% | 75.9% | -4.2% | 70 | 93 | 75.3% |
| 1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services. | 64.7% | 66.2% | 72.6% | 12.2% | 87 | 109 | 79.8% |
| 1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services. | 67.4% | 61.7% | 66.1% | -1.9% | 393 | 607 | 64.7% |
| 2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy. | 38.8% | 41.8% | 37.4% | -3.5% | 78 | 221 | 35.3% |
| 2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum. | 69.1% | 70.8% | 65.9% | -4.6% | 59 | 87 | 67.8% |
| 3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1) | | | | | | | |
| 3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1) | | | | | | | |
| 4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were: | | | | | | | |
| a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1) | 69.9% | 69.6% | 64.2% | -8.1% | 163 | 259 | 62.9% |
| b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile) | 11.0% | 13.0% | 16.5% | 49.3% | 40 | 259 | 15.4% |
| c. Obese (BMI at or greater than the 95th percentile). | 14.5% | 13.6% | 13.5% | -6.8% | 31 | 259 | 12.0% |
| d. Underweight (BMI less than the 5th percentile). | 4.6% | 3.8% | 5.8% | 26.2% | 25 | 259 | 9.7% |
| 4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3) | | | | | | | |
| a. Recommended prenatal weight gain | 41.6% | 29.2% | 27.7% | -33.5% | 21 | 77 | 27.3% |
| b. Excessive prenatal weight gain | 37.6% | 49.6% | 47.4% | 26.3% | 42 | 77 | 54.5% |
| c. Inadequate prenatal weight gain | 20.8% | 21.3% | 24.9% | 19.6% | 14 | 77 | 18.2% |
| 5A. Percentage of women who participated in WIC who initiated breastfeeding. | 54.0% | 65.4% | 65.0% | 20.5% | 77 | 109 | 70.6% |
| 5B. Percentage of infants who participated in WIC who were: | | | | | | | |
| a. Breastfeeding at 6 weeks of age (1) | 40.3% | 48.4% | 43.0% | 6.7% | 61 | 138 | 44.2% |
| b. Breastfeeding at 6 months of age | 25.2% | 27.0% | 25.5% | 1.2% | 30 | 138 | 21.7% |

(1) Item appears also on other WCH Section Agreement Addenda. Data for 3A and 3B are not presented due to limitation in recent trend data.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI >=30.0); Adapted from the National Academy of Sciences. Weight Gain During Pregnancy: Reexamining the Guidelines. Washington, DC : National Academy Press, 2009.

SFY 2014-2015 AGREEMENT ADDENDA DATA
WIC QUALITY ASSURANCE DELIVERABLES

Pasquotank

| Process Outcome Objectives | 2004 to 2006 | 2007 to 2009 | 2010 to 2012 | % Change from 2004 to 2012 | 2012 Data | | |
|---|--------------------|--------------------|--------------------|----------------------------------|-----------|-------|---------|
| | | | | | Num | Den | Percent |
| 1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2) | 74.9% | 59.1% | 61.6% | -17.7% | 146 | 216 | 67.6% |
| 1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services. | 72.0% | 72.6% | 72.9% | 1.2% | 210 | 288 | 72.9% |
| 1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services. | 54.3% | 54.1% | 54.8% | 1.0% | 954 | 1,690 | 56.4% |
| 2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy. | 36.3% | 35.1% | 27.0% | -25.6% | 463 | 1,750 | 26.5% |
| 2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum. | 33.5% | 41.9% | 58.0% | 73.3% | 509 | 681 | 74.7% |
| 3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1) | | | | | | | |
| 3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1) | | | | | | | |
| 4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were: | | | | | | | |
| a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1) | 66.4% | 65.5% | 62.2% | -6.3% | 470 | 757 | 62.1% |
| b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile) | 14.8% | 15.9% | 14.7% | -0.3% | 101 | 757 | 13.3% |
| c. Obese (BMI at or greater than the 95th percentile). | 14.5% | 15.6% | 15.8% | 8.7% | 120 | 757 | 15.9% |
| d. Underweight (BMI less than the 5th percentile). | 4.4% | 2.9% | 7.3% | 67.1% | 66 | 757 | 8.7% |
| 4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3) | | | | | | | |
| a. Recommended prenatal weight gain | 33.9% | 31.9% | 27.2% | -19.7% | 54 | 197 | 27.4% |
| b. Excessive prenatal weight gain | 42.6% | 42.7% | 38.7% | -9.3% | 74 | 197 | 37.6% |
| c. Inadequate prenatal weight gain | 23.5% | 25.4% | 34.1% | 45.2% | 69 | 197 | 35.0% |
| 5A. Percentage of women who participated in WIC who initiated breastfeeding. | 20.8% | 23.9% | 32.6% | 56.9% | 275 | 857 | 32.1% |
| 5B. Percentage of infants who participated in WIC who were: | | | | | | | |
| a. Breastfeeding at 6 weeks of age (1) | 8.7% | 10.8% | 17.3% | 97.5% | 169 | 714 | 23.7% |
| b. Breastfeeding at 6 months of age | 3.4% | 3.4% | 9.2% | 172.6% | 85 | 714 | 11.9% |

(1) Item appears also on other WCH Section Agreement Addenda. Data for 3A and 3B are not presented due to limitation in recent trend data.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI >=30.0); Adapted from the National Academy of Sciences. Weight Gain During Pregnancy: Reexamining the Guidelines. Washington, DC : National Academy Press, 2009.

SFY 2014-2015 AGREEMENT ADDENDA DATA

WIC QUALITY ASSURANCE DELIVERABLES

Perquimans

| Process Outcome Objectives | 2004 to 2006 | 2007 to 2009 | 2010 to 2012 | % Change from 2004 to 2012 | 2012 Data | | |
|---|--------------------|--------------------|--------------------|----------------------------------|-----------|-------|---------|
| | | | | | Num | Den | Percent |
| 1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2) | 75.2% | 70.6% | 74.8% | -0.6% | 60 | 78 | 76.9% |
| 1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services. | 72.5% | 74.7% | 74.3% | 2.4% | 73 | 90 | 81.1% |
| 1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services. | 59.0% | 55.8% | 56.4% | -4.4% | 286 | 476 | 60.1% |
| 2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy. | 36.3% | 35.1% | 27.0% | -25.6% | 463 | 1,750 | 26.5% |
| 2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum. | 33.5% | 41.9% | 58.0% | 73.3% | 509 | 681 | 74.7% |
| 3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1) | | | | | | | |
| 3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1) | | | | | | | |
| 4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were: | | | | | | | |
| a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1) | 61.5% | 65.0% | 58.6% | -4.7% | 268 | 437 | 61.3% |
| b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile) | 16.9% | 16.4% | 15.4% | -9.3% | 58 | 437 | 13.3% |
| c. Obese (BMI at or greater than the 95th percentile). | 19.2% | 17.9% | 18.5% | -3.5% | 73 | 437 | 16.7% |
| d. Underweight (BMI less than the 5th percentile). | 2.4% | 0.7% | 7.6% | 210.6% | 38 | 437 | 8.7% |
| 4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3) | | | | | | | |
| a. Recommended prenatal weight gain | 33.8% | 40.5% | 24.7% | -26.8% | 21 | 69 | 30.4% |
| b. Excessive prenatal weight gain | 38.5% | 37.2% | 47.4% | 23.0% | 30 | 69 | 43.5% |
| c. Inadequate prenatal weight gain | 27.7% | 22.3% | 27.9% | 0.7% | 18 | 69 | 26.1% |
| 5A. Percentage of women who participated in WIC who initiated breastfeeding. | 20.8% | 23.9% | 32.6% | 56.9% | 275 | 857 | 32.1% |
| 5B. Percentage of infants who participated in WIC who were: | | | | | | | |
| a. Breastfeeding at 6 weeks of age (1) | 8.7% | 10.8% | 17.3% | 97.5% | 169 | 714 | 23.7% |
| b. Breastfeeding at 6 months of age | 3.4% | 3.4% | 9.2% | 172.6% | 85 | 714 | 11.9% |

(1) Item appears also on other WCH Section Agreement Addenda. Data for 3A and 3B are not presented due to limitation in recent trend data.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI >=30.0); Adapted from the National Academy of Sciences. Weight Gain During Pregnancy: Reexamining the Guidelines. Washington, DC : National Academy Press, 2009.

SFY 2014-2015 AGREEMENT ADDENDA DATA

WIC QUALITY ASSURANCE DELIVERABLES

Polk

| Process Outcome Objectives | 2004 to 2006 | 2007 to 2009 | 2010 to 2012 | % Change from 2004 to 2012 | 2012 Data | | |
|---|--------------------|--------------------|--------------------|----------------------------------|-----------|-------|---------|
| | | | | | Num | Den | Percent |
| 1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2) | 61.2% | 64.5% | 67.6% | 10.3% | 53 | 72 | 73.6% |
| 1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services. | 67.7% | 65.2% | 67.1% | -0.9% | 66 | 91 | 72.5% |
| 1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services. | 55.6% | 55.0% | 55.8% | 0.3% | 314 | 561 | 56.0% |
| 2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy. | 32.9% | 35.4% | 33.2% | 1.0% | 847 | 2,654 | 31.9% |
| 2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum. | 71.0% | 73.2% | 78.0% | 9.8% | 785 | 990 | 79.3% |
| 3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1) | | | | | | | |
| 3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1) | | | | | | | |
| 4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were: | | | | | | | |
| a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1) | 66.0% | 68.4% | 64.3% | -2.6% | 130 | 206 | 63.1% |
| b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile) | 16.5% | 16.9% | 15.4% | -6.2% | 28 | 206 | 13.6% |
| c. Obese (BMI at or greater than the 95th percentile). | 15.4% | 12.6% | 18.0% | 17.0% | 43 | 206 | 20.9% |
| d. Underweight (BMI less than the 5th percentile). | 2.1% | 2.1% | 2.2% | 4.6% | 5 | 206 | 2.4% |
| 4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3) | | | | | | | |
| a. Recommended prenatal weight gain | 28.5% | 29.6% | 31.5% | 10.5% | 24 | 66 | 36.4% |
| b. Excessive prenatal weight gain | 38.9% | 46.4% | 49.3% | 26.8% | 33 | 66 | 50.0% |
| c. Inadequate prenatal weight gain | 32.6% | 24.0% | 19.2% | -41.0% | 9 | 66 | 13.6% |
| 5A. Percentage of women who participated in WIC who initiated breastfeeding. | 49.2% | 56.9% | 59.1% | 20.2% | 670 | 1,190 | 56.3% |
| 5B. Percentage of infants who participated in WIC who were: | | | | | | | |
| a. Breastfeeding at 6 weeks of age (1) | 30.0% | 29.7% | 32.2% | 7.2% | 374 | 1,081 | 34.6% |
| b. Breastfeeding at 6 months of age | 15.2% | 14.0% | 17.3% | 13.9% | 185 | 1,081 | 17.1% |

(1) Item appears also on other WCH Section Agreement Addenda. Data for 3A and 3B are not presented due to limitation in recent trend data.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI >=30.0); Adapted from the National Academy of Sciences. Weight Gain During Pregnancy: Reexamining the Guidelines. Washington, DC : National Academy Press, 2009.

SFY 2014-2015 AGREEMENT ADDENDA DATA

WIC QUALITY ASSURANCE DELIVERABLES

Rutherford

| Process Outcome Objectives | 2004 to 2006 | 2007 to 2009 | 2010 to 2012 | % Change from 2004 to 2012 | 2012 Data | | |
|---|--------------------|--------------------|--------------------|----------------------------------|-----------|-------|---------|
| | | | | | Num | Den | Percent |
| 1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2) | 63.7% | 71.1% | 83.7% | 31.3% | 419 | 475 | 88.2% |
| 1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services. | 78.9% | 79.8% | 77.4% | -1.9% | 439 | 567 | 77.4% |
| 1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services. | 59.9% | 61.5% | 60.1% | 0.3% | 1,767 | 2,888 | 61.2% |
| 2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy. | 32.9% | 35.4% | 33.2% | 1.0% | 847 | 2,654 | 31.9% |
| 2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum. | 71.0% | 73.2% | 78.0% | 9.8% | 785 | 990 | 79.3% |
| 3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1) | | | | | | | |
| 3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1) | | | | | | | |
| 4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were: | | | | | | | |
| a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1) | 67.5% | 66.0% | 65.1% | -3.6% | 516 | 787 | 65.6% |
| b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile) | 14.8% | 16.3% | 15.1% | 1.7% | 114 | 787 | 14.5% |
| c. Obese (BMI at or greater than the 95th percentile). | 13.0% | 13.6% | 14.4% | 10.2% | 114 | 787 | 14.5% |
| d. Underweight (BMI less than the 5th percentile). | 4.6% | 4.0% | 5.5% | 17.6% | 43 | 787 | 5.5% |
| 4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3) | | | | | | | |
| a. Recommended prenatal weight gain | 30.6% | 31.7% | 26.4% | -13.7% | 124 | 467 | 26.6% |
| b. Excessive prenatal weight gain | 38.9% | 40.1% | 50.6% | 30.0% | 231 | 467 | 49.5% |
| c. Inadequate prenatal weight gain | 30.5% | 28.2% | 23.0% | -24.5% | 112 | 467 | 24.0% |
| 5A. Percentage of women who participated in WIC who initiated breastfeeding. | 49.2% | 56.9% | 59.1% | 20.2% | 670 | 1,190 | 56.3% |
| 5B. Percentage of infants who participated in WIC who were: | | | | | | | |
| a. Breastfeeding at 6 weeks of age (1) | 30.0% | 29.7% | 32.2% | 7.2% | 374 | 1,081 | 34.6% |
| b. Breastfeeding at 6 months of age | 15.2% | 14.0% | 17.3% | 13.9% | 185 | 1,081 | 17.1% |

(1) Item appears also on other WCH Section Agreement Addenda. Data for 3A and 3B are not presented due to limitation in recent trend data.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI >=30.0); Adapted from the National Academy of Sciences. Weight Gain During Pregnancy: Reexamining the Guidelines. Washington, DC : National Academy Press, 2009.

SFY 2014-2015 AGREEMENT ADDENDA DATA

WIC QUALITY ASSURANCE DELIVERABLES

Tyrrell

| Process Outcome Objectives | 2004 to 2006 | 2007 to 2009 | 2010 to 2012 | % Change from 2004 to 2012 | 2012 Data | | |
|---|--------------------|--------------------|--------------------|----------------------------------|-----------|-----|---------|
| | | | | | Num | Den | Percent |
| 1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2) | 78.9% | 90.9% | 81.9% | 3.8% | 18 | 23 | 78.3% |
| 1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services. | 67.7% | 79.5% | 82.8% | 22.3% | 27 | 29 | 93.1% |
| 1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services. | 65.6% | 68.9% | 73.1% | 11.5% | 130 | 166 | 78.3% |
| 2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy. | 35.3% | 33.0% | 28.0% | -20.7% | 269 | 995 | 27.0% |
| 2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum. | 81.3% | 82.0% | 80.5% | -1.0% | 268 | 320 | 83.8% |
| 3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1) | | | | | | | |
| 3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1) | | | | | | | |
| 4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were: | | | | | | | |
| a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1) | 62.0% | 65.2% | 63.6% | 2.6% | 130 | 235 | 55.3% |
| b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile) | 15.6% | 13.5% | 12.5% | -19.9% | 31 | 235 | 13.2% |
| c. Obese (BMI at or greater than the 95th percentile). | 19.3% | 18.3% | 17.3% | -10.3% | 54 | 235 | 23.0% |
| d. Underweight (BMI less than the 5th percentile). | 3.1% | 3.1% | 6.6% | 113.5% | 20 | 235 | 8.5% |
| 4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3) | | | | | | | |
| a. Recommended prenatal weight gain | 29.6% | 35.9% | 23.5% | -20.8% | 6 | 28 | 21.4% |
| b. Excessive prenatal weight gain | 42.6% | 44.4% | 45.7% | 7.2% | 12 | 28 | 42.9% |
| c. Inadequate prenatal weight gain | 27.8% | 19.7% | 30.9% | 11.1% | 10 | 28 | 35.7% |
| 5A. Percentage of women who participated in WIC who initiated breastfeeding. | 40.3% | 36.6% | 23.0% | -43.0% | 90 | 388 | 23.2% |
| 5B. Percentage of infants who participated in WIC who were: | | | | | | | |
| a. Breastfeeding at 6 weeks of age (1) | 25.5% | 21.6% | 11.2% | -56.3% | 32 | 351 | 9.1% |
| b. Breastfeeding at 6 months of age | 12.0% | 9.5% | 4.3% | -64.4% | 17 | 351 | 4.8% |

(1) Item appears also on other WCH Section Agreement Addenda. Data for 3A and 3B are not presented due to limitation in recent trend data.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI >=30.0); Adapted from the National Academy of Sciences. Weight Gain During Pregnancy: Reexamining the Guidelines. Washington, DC : National Academy Press, 2009.

SFY 2014-2015 AGREEMENT ADDENDA DATA

WIC QUALITY ASSURANCE DELIVERABLES

Vance

| Process Outcome Objectives | 2004 to 2006 | 2007 to 2009 | 2010 to 2012 | % Change from 2004 to 2012 | 2012 Data | | |
|---|--------------------|--------------------|--------------------|----------------------------------|-----------|-------|---------|
| | | | | | Num | Den | Percent |
| 1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2) | 69.9% | 75.9% | 71.8% | 2.6% | 299 | 413 | 72.4% |
| 1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services. | 76.2% | 74.8% | 76.2% | -0.1% | 405 | 539 | 75.1% |
| 1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services. | 55.3% | 57.2% | 56.1% | 1.5% | 1,585 | 2,764 | 57.3% |
| 2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy. | 31.0% | 33.2% | 27.7% | -10.5% | 430 | 1,554 | 27.7% |
| 2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum. | 80.2% | 75.2% | 76.7% | -4.4% | 556 | 707 | 78.6% |
| 3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1) | | | | | | | |
| 3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1) | | | | | | | |
| 4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were: | | | | | | | |
| a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1) | 67.1% | 67.9% | 66.2% | -1.4% | 455 | 681 | 66.8% |
| b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile) | 14.6% | 12.7% | 11.6% | -21.0% | 81 | 681 | 11.9% |
| c. Obese (BMI at or greater than the 95th percentile). | 13.5% | 11.3% | 12.0% | -11.1% | 77 | 681 | 11.3% |
| d. Underweight (BMI less than the 5th percentile). | 4.7% | 8.1% | 10.2% | 116.8% | 68 | 681 | 10.0% |
| 4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3) | | | | | | | |
| a. Recommended prenatal weight gain | 30.6% | 31.9% | 29.7% | -3.2% | 90 | 358 | 25.1% |
| b. Excessive prenatal weight gain | 44.8% | 40.4% | 38.9% | -13.2% | 157 | 358 | 43.9% |
| c. Inadequate prenatal weight gain | 24.6% | 27.7% | 31.5% | 28.0% | 111 | 358 | 31.0% |
| 5A. Percentage of women who participated in WIC who initiated breastfeeding. | 40.4% | 52.0% | 45.6% | 13.0% | 438 | 893 | 49.0% |
| 5B. Percentage of infants who participated in WIC who were: | | | | | | | |
| a. Breastfeeding at 6 weeks of age (1) | 28.0% | 38.1% | 35.4% | 26.2% | 337 | 928 | 36.3% |
| b. Breastfeeding at 6 months of age | 19.5% | 26.5% | 25.8% | 32.3% | 256 | 928 | 27.6% |

(1) Item appears also on other WCH Section Agreement Addenda. Data for 3A and 3B are not presented due to limitation in recent trend data.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI >=30.0); Adapted from the National Academy of Sciences. Weight Gain During Pregnancy: Reexamining the Guidelines. Washington, DC : National Academy Press, 2009.

SFY 2014-2015 AGREEMENT ADDENDA DATA
WIC QUALITY ASSURANCE DELIVERABLES

Washington

| Process Outcome Objectives | 2004 to 2006 | 2007 to 2009 | 2010 to 2012 | % Change from 2004 to 2012 | 2012 Data | | |
|---|--------------------|--------------------|--------------------|----------------------------------|-----------|-----|---------|
| | | | | | Num | Den | Percent |
| 1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2) | 85.7% | 89.6% | 84.2% | -1.7% | 88 | 103 | 85.4% |
| 1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services. | 74.7% | 74.5% | 83.7% | 11.9% | 99 | 116 | 85.3% |
| 1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services. | 69.6% | 65.7% | 63.0% | -9.4% | 441 | 662 | 66.6% |
| 2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy. | 35.3% | 33.0% | 28.0% | -20.7% | 269 | 995 | 27.0% |
| 2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum. | 81.3% | 82.0% | 80.5% | -1.0% | 268 | 320 | 83.8% |
| 3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1) | | | | | | | |
| 3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1) | | | | | | | |
| 4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were: | | | | | | | |
| a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1) | 68.9% | 66.2% | 63.8% | -7.5% | 196 | 338 | 58.0% |
| b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile) | 9.9% | 9.8% | 12.0% | 21.4% | 50 | 338 | 14.8% |
| c. Obese (BMI at or greater than the 95th percentile). | 9.3% | 12.0% | 14.9% | 59.6% | 64 | 338 | 18.9% |
| d. Underweight (BMI less than the 5th percentile). | 11.8% | 12.0% | 9.3% | -21.6% | 28 | 338 | 8.3% |
| 4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3) | | | | | | | |
| a. Recommended prenatal weight gain | 25.0% | 29.1% | 29.4% | 17.5% | 26 | 91 | 28.6% |
| b. Excessive prenatal weight gain | 44.8% | 45.7% | 40.6% | -9.3% | 37 | 91 | 40.7% |
| c. Inadequate prenatal weight gain | 30.2% | 25.2% | 30.0% | -0.7% | 28 | 91 | 30.8% |
| 5A. Percentage of women who participated in WIC who initiated breastfeeding. | 40.3% | 36.6% | 23.0% | -43.0% | 90 | 388 | 23.2% |
| 5B. Percentage of infants who participated in WIC who were: | | | | | | | |
| a. Breastfeeding at 6 weeks of age (1) | 25.5% | 21.6% | 11.2% | -56.3% | 32 | 351 | 9.1% |
| b. Breastfeeding at 6 months of age | 12.0% | 9.5% | 4.3% | -64.4% | 17 | 351 | 4.8% |

(1) Item appears also on other WCH Section Agreement Addenda. Data for 3A and 3B are not presented due to limitation in recent trend data.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI >=30.0); Adapted from the National Academy of Sciences. Weight Gain During Pregnancy: Reexamining the Guidelines. Washington, DC : National Academy Press, 2009.

SFY 2014-2015 AGREEMENT ADDENDA DATA

WIC QUALITY ASSURANCE DELIVERABLES

Watauga

| Process Outcome Objectives | 2004 to 2006 | 2007 to 2009 | 2010 to 2012 | % Change from 2004 to 2012 | 2012 Data | | |
|---|--------------------|--------------------|--------------------|----------------------------------|-----------|-------|---------|
| | | | | | Num | Den | Percent |
| 1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2) | 70.4% | 49.0% | 73.5% | 4.4% | 122 | 155 | 78.7% |
| 1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services. | 63.8% | 66.0% | 57.5% | -9.9% | 118 | 203 | 58.1% |
| 1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services. | 62.0% | 62.1% | 60.6% | -2.2% | 631 | 1,047 | 60.3% |
| 2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy. | 48.9% | 44.6% | 37.2% | -24.0% | 172 | 489 | 35.2% |
| 2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum. | 59.9% | 69.4% | 69.8% | 16.6% | 139 | 192 | 72.4% |
| 3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1) | | | | | | | |
| 3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1) | | | | | | | |
| 4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were: | | | | | | | |
| a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1) | 68.9% | 73.7% | 64.8% | -5.9% | 235 | 371 | 63.3% |
| b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile) | 14.6% | 13.5% | 17.3% | 18.4% | 59 | 371 | 15.9% |
| c. Obese (BMI at or greater than the 95th percentile). | 12.7% | 10.0% | 13.4% | 5.7% | 58 | 371 | 15.6% |
| d. Underweight (BMI less than the 5th percentile). | 3.8% | 2.8% | 4.4% | 16.3% | 19 | 371 | 5.1% |
| 4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3) | | | | | | | |
| a. Recommended prenatal weight gain | 30.2% | 31.5% | 33.3% | 10.1% | 48 | 153 | 31.4% |
| b. Excessive prenatal weight gain | 51.5% | 49.9% | 52.1% | 1.1% | 85 | 153 | 55.6% |
| c. Inadequate prenatal weight gain | 18.3% | 18.6% | 14.7% | -19.9% | 20 | 153 | 13.1% |
| 5A. Percentage of women who participated in WIC who initiated breastfeeding. | 56.2% | 66.5% | 70.6% | 25.6% | 149 | 221 | 67.4% |
| 5B. Percentage of infants who participated in WIC who were: | | | | | | | |
| a. Breastfeeding at 6 weeks of age (1) | 43.7% | 48.2% | 53.1% | 21.5% | 120 | 218 | 55.0% |
| b. Breastfeeding at 6 months of age | 27.0% | 28.9% | 33.3% | 23.5% | 88 | 218 | 40.4% |

(1) Item appears also on other WCH Section Agreement Addenda. Data for 3A and 3B are not presented due to limitation in recent trend data.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI >=30.0); Adapted from the National Academy of Sciences. Weight Gain During Pregnancy: Reexamining the Guidelines. Washington, DC : National Academy Press, 2009.

SFY 2014-2015 AGREEMENT ADDENDA DATA

WIC QUALITY ASSURANCE DELIVERABLES

Yancey

| Process Outcome Objectives | 2004 to 2006 | 2007 to 2009 | 2010 to 2012 | % Change from 2004 to 2012 | 2012 Data | | |
|---|--------------------|--------------------|--------------------|----------------------------------|-----------|-----|---------|
| | | | | | Num | Den | Percent |
| 1A. Percentage of pregnant women enrolled in Medicaid who received prenatal WIC Program services. (1) (2) | 68.9% | 68.6% | 72.4% | 5.2% | 88 | 113 | 77.9% |
| 1B. Percentage of children less than 12 months of age enrolled in Medicaid who received WIC Program services. | 60.6% | 61.4% | 67.1% | 10.6% | 88 | 130 | 67.7% |
| 1C. Percentage of children 1 to 5 years of age enrolled in Medicaid who received WIC Program Services. | 60.0% | 60.2% | 65.0% | 8.2% | 410 | 627 | 65.4% |
| 2A. Percentage of pregnant women who participated in WIC and received WIC Program services during the first trimester of pregnancy. | 39.5% | 42.6% | 35.7% | -9.6% | 125 | 374 | 33.4% |
| 2B. Percentage of postpartum women who participated in WIC during pregnancy and were recertified for WIC by 6 weeks postpartum. | 66.4% | 66.3% | 68.1% | 2.5% | 87 | 127 | 68.5% |
| 3A. Percentage of children less than 12 months of age who were served in the Child Health Clinic and received WIC Program services. (1) | | | | | | | |
| 3B. Percentage of children 1 to 5 years of age who were served in the Child Health Clinic and received WIC Program Services. (1) | | | | | | | |
| 4A. Percentage of children 2 to 4 years of age who received services from the health department and who, according to their Body Mass Index (BMI) for age and gender were: | | | | | | | |
| a. At a healthy weight (BMI less than the 85th percentile but at or greater than the 5th percentile) (1) | 66.3% | 67.6% | 64.4% | -2.9% | 204 | 301 | 67.8% |
| b. Overweight (BMI at or above the 85th percentile but less than the 95th percentile) | 17.3% | 15.7% | 16.2% | -6.0% | 38 | 301 | 12.6% |
| c. Obese (BMI at or greater than the 95th percentile). | 13.8% | 12.6% | 14.3% | 3.2% | 33 | 301 | 11.0% |
| d. Underweight (BMI less than the 5th percentile). | 2.6% | 4.1% | 5.1% | 98.6% | 26 | 301 | 8.6% |
| 4B. Percentage of women with live term singleton births who received WIC Program services during pregnancy and who gained recommended/excessive/inadequate weight according to the National Academy of Sciences - Institute of Medicine 2009 recommended total weight gain ranges during pregnancy. (3) | | | | | | | |
| a. Recommended prenatal weight gain | 32.4% | 32.7% | 29.5% | -8.9% | 22 | 90 | 24.4% |
| b. Excessive prenatal weight gain | 44.9% | 43.8% | 43.9% | -2.2% | 48 | 90 | 53.3% |
| c. Inadequate prenatal weight gain | 22.8% | 23.6% | 26.7% | 17.0% | 20 | 90 | 22.2% |
| 5A. Percentage of women who participated in WIC who initiated breastfeeding. | 72.0% | 75.1% | 63.4% | -11.9% | 99 | 153 | 64.7% |
| 5B. Percentage of infants who participated in WIC who were: | | | | | | | |
| a. Breastfeeding at 6 weeks of age (1) | 35.5% | 36.2% | 41.6% | 17.2% | 64 | 162 | 39.5% |
| b. Breastfeeding at 6 months of age | 22.4% | 21.9% | 30.9% | 37.9% | 46 | 162 | 28.4% |

(1) Item appears also on other WCH Section Agreement Addenda. Data for 3A and 3B are not presented due to limitation in recent trend data.

(2) These data exclude women enrolled in Medicaid only for emergency delivery.

(3) Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI >=30.0); Adapted from the National Academy of Sciences. Weight Gain During Pregnancy: Reexamining the Guidelines. Washington, DC : National Academy Press, 2009.